

A single dose of hydrocortisone and an albumin infusion were given initially. She was subsequently treated with warmed intravenous fluids for three days and antibiotics for 10 days. She recovered completely and continues to enjoy good health.

Profound hypothermia is extremely rare in children over 5 years of age. Results of investigations excluded infective and endocrine causes. A normal MRI brain scan showed there was no lesion of the hypothalamus or corpus callosum.

Ibuprofen is commonly prescribed for a raised temperature and is well tolerated in children. Side effects are not common, even in overdose.<sup>1</sup> Nevertheless we postulate that ibuprofen was responsible for hypothermia in this case. We are not aware of any published evidence documenting hypothermia after a single therapeutic dose of ibuprofen, but it has been recorded in a few cases of accidental and deliberate overdosage. Although patients may sometimes receive ibuprofen in toxic quantities, hypothermia is not a consistent feature.<sup>2,3</sup> Hypothermia in overdosage is attributed to central nervous system depression.<sup>4</sup>

#### P R Desai

Southend Hospital, Room 2, Doctor's Quarters, Chelmsford Accommodation, St John's Hospital, Chelmsford CM2 9BG, UK; prpravin@yahoo.com

#### S Sriskandan

Department of Paediatrics, Southend Hospital, Prittlewell Chase, Southend on Sea SSO 0RY, UK

#### References

- 1 **Court H**, Streete P, Volaus GN. Acute poisoning with ibuprofen. *Hum Toxicol* 1983;**2**:381.
- 2 **Ritter A**, Barnett E. Ibuprofen overdose presenting with severe agitation and hypothermia [letter]. *Am J Emerg Med* 1998;**16**:549–50.
- 3 **Chelluri L**, Jastremski M. Coma caused by ibuprofen overdose. *Crit Care Med* 1986;**14**:1078–9.
- 4 **Le HT**, Bosse GM, Tsai Y. Ibuprofen overdose complicated by renal failure, adult respiratory distress syndrome and metabolic acidosis. *Clin Toxicol* 1994;**32**:315–20.

### Vagal overactivity: a risk factor of sudden infant death syndrome?

Since early 1990, the incidence of sudden infant death syndrome (SIDS) has dropped sharply because of public health campaigns decrying the dangers of the prone sleep position. The other known risk factors, such as preterm birth and young maternal age, are less susceptible to prevention campaigns.<sup>1</sup>

Disordered autonomic function, including cardiorespiratory control, has been suggested to be involved in SIDS.<sup>2,3</sup> Vagal overactivity (VO), characterised by breath holding spells and repeated syncope in specific circumstances, has been described as a manifestation of autonomic dysfunction.<sup>4</sup> To investigate a possible relation between VO and SIDS, we investigated 65 children presenting documented VO; for example, clinical characteristics and a positive test for eyeball compression and/or electrocardiographic monitoring. Parents of these children were interviewed about their family history, especially with respect to the occurrence of SIDS among their other children.

Among their siblings, five of 126 had died of SIDS. All five children were full term infants. The average maternal age, birth weight, and age at death were respectively 27.4 (3.5) years, 3.3 (0.3) kg, and 3.5 (1.1) months. The rates of SIDS in siblings of children with VO were compared to those in the general population using the standardised incidence ratio (SIR), which is the ratio of the observed number to the expected number of cases of SIDS calculated by French incidence rates. The expected number of SIDS was 0.17 and hence the SIR was 29.4 (95% CI 9.5 to 68.6;  $p < 0.00011$ ). Our result showed an overall significant excess of SIDS among siblings of children with VO. We verified that recruited children had not come to the centre because of a family history of SIDS. Since children with a positive family history of SIDS could be followed up more regularly than others, we estimated the SIR separately among siblings of children recruited during their follow up and those of children recruited during their first visit, and verified that there was no significant difference in SIR between these cases.

Despite the marked decline in SIDS, it is still the leading cause of postneonatal mortality. Better knowledge of other risk factors may allow identification of populations at high risk and a possible decline in infant mortality from SIDS through the implementation of appropriate prevention measures. Our findings suggest that VO may be involved in SIDS and that children with VO or a family history of VO may be a population at potential high risk of SIDS.

#### T Shojaei-Brosseau, C Bonaiti-Pellie

Unité de Recherche en Épidémiologie des Cancers, INSERM U521, Villejuif, France

#### S Lyonnet, J Feingold

Unité de recherche sur les Handicaps Génétiques de l'Enfant, INSERM, U393, Paris, France

#### V Lucet

Centre de Cardiologie Infantile du Château des Côtes, Les Loges-en-Josas, France

Correspondence to: Dr T Shojaei-Brosseau, Service de Biostatistiques, Institut Curie, 70 rue Mouffetard, 75005 Paris, France; taraneh.shojaei@curie.net

#### References

- 1 **Irgens LM**, Markestad T, Baste V, et al. Sleeping position and sudden infant death syndrome in Norway 1967–91. *Arch Dis Child* 1995;**72**:478–82.
- 2 **Lucet V**, de Bethman O, Denjoy I. Paroxysmal over-activity, apparent life-threatening event and sudden infant death. *Biol Neonate* 2000;**78**:1–7.
- 3 **Fox GP**, Matthews TG. Autonomic dysfunction at different ambient temperatures in infants at risk of sudden infant death. *Lancet* 1989;**2**:1065–7.
- 4 **Lurie KG**, Benditt D. Syncope and the autonomic nervous system. *J Cardiovasc Electrophysiol* 1996;**7**:760–6.

### Perforated duodenal ulcer disclosing medium chain acyl-CoA dehydrogenase deficiency

Medium chain acyl-CoA dehydrogenase deficiency (MCADD; McKusick 201450) typically presents in the first two years of life with recurrent episodes of hypoketotic hypoglycaemia, lethargy, coma, or sudden infant death. The trigger may be fasting, intercurrent infections, anaesthesia, or surgery. Incidence in the

UK is estimated at 0.45–1/10 000 live births.<sup>1</sup> We describe the case of a child who presented with marked encephalopathy unexplained by perforated duodenal ulcer, which led to the diagnosis of MCADD.

A 2 year old girl presented with a three week history of coryzal symptoms and three day history of frequent coffee ground vomiting. She was shocked, and had hepatomegaly and decreased conscious level. Blood glucose was 3.9 mmol/l (reference interval 3.3–5.5), plasma sodium 129 mmol/l (135–147), potassium 5.2 mmol/l (3.5–5.0), urea 17.8 mmol/l (3.3–6.6), creatinine 36 mmol/l (30–74), bicarbonate 15 mmol/l (21–28), base excess –5.4 mmol/l (–4 to +2) and C reactive protein 4 mg/l (0–5). Liver function tests and clotting were normal. She was resuscitated with a total of 50 ml/kg of colloid and crystalloid. The following day she relapsed with abdominal distension, shock, and deteriorating conscious level. Investigations showed glucose 14.2 mmol/l, amylase 20 IU/l (8–85), AST 186 IU/l (10–45), and ALT 129 IU/l (10–40). An x ray examination of the abdomen showed free air under the right hemidiaphragm. Emergency laparotomy revealed a single, 1 cm × 1 cm acute perforation in the second part of the duodenum. Histology and rapid urease test (CLO) of the duodenal biopsy for *Helicobacter pylori* were negative. Fasting blood gastrin was 20 mU/l (10–100). She was discharged home taking omeprazole. Upper gastrointestinal endoscopic biopsy (eight weeks later) for histopathology and CLO test from oesophagus, stomach, antrum, and duodenum were normal.

Analysis of urinary organic acids by gas chromatography and mass spectrometry, obtained a day after clinical presentation, revealed a marked increase in 5-hydroxyhexanoic acid (21% of total organic acids); a modest dicarboxylicaciduria (suberic accounted for 8% and adipic 6% of total organic acids); and a small but significant quantity of hexanoyl glycine (2% total organic acids) in the absence of ketonuria.

Blood obtained a week after clinical presentation, when analysed by tandem mass spectrometry, showed octanoylcarnitine 2.91 μmol/l (≤0.19), hexanoylcarnitine 0.67 μmol/l (≤0.29), and decenoylcarnitine 0.63 μmol/l (≤0.10), with a subnormal concentration of acetylcarnitine 4.0 μmol/l (6.2–27.5). This profile was consistent with MCADD. Polymerase chain reaction/restriction digests based method revealed two mutations in the MCAD gene.

The clinical details coupled with the absence of ketones and the increased 5-hydroxyhexanoic acid led us to look for an abnormality in the oxidation of fatty acids, and resulted in identification of the minor constituent, hexanoylglycine that is recognised as an indicative marker of MCADD. Increases in urinary hexanoylglycine and 5-hydroxyhexanoic acids in the absence of ketonuria have been reported previously in MCAD patients during clinical attack,<sup>2</sup> and also in a boy who died.<sup>3</sup> Our case was unusual in that the amount of 5-hydroxyhexanoic acid was greater than even the sum of the individual dicarboxylic acids present, although high levels of 5-hydroxyhexanoic acids are reported in acute episodes.<sup>4</sup> The increased concentration of octonoyl carnitine in blood was also consistent with a diagnosis of MCADD.

We believe that this is the first report of MCADD presenting with duodenal ulcer. It could be argued that the ulcer was the primary problem and that the decompensation was caused by the subsequent illness.