

Sequencing of the Gene Encoding the Major Pilin of Pilus Colonization Factor Antigen III (CFA/III) of Human Enterotoxigenic *Escherichia coli* and Evidence that CFA/III Is Related to Type IV Pili

TOORU TANIGUCHI,¹ YUJI FUJINO,¹ KOICHIRO YAMAMOTO,¹ TOSHIO MIWATANI,^{1,2}
AND TAKESHI HONDA^{1*}

Department of Bacteriology and Serology, Research Institute for Microbial Diseases, Osaka University, Yamadaoka, Suita, Osaka 565,¹ and Department of Nutritional Science, Faculty of Health and Welfare Science, Okayama Prefectural University, Kuboki, Sojya, Okayama,² Japan

Received 8 August 1994/Returned for modification 7 September 1994/Accepted 16 November 1994

The plasmid-encoded structural gene *cofA* necessary for the production of the major pilin subunit of pilus colonization factor antigen III (CFA/III) of human enterotoxigenic *Escherichia coli* was identified, and the nucleotide sequence of the gene was determined. *cofA* consists of 714 nucleotides encoding a 238-amino-acid protein (molecular weight of 25,309). *CofA* seems to be a precursor of CFA/III pilin, because the first 23 residues of the N-terminal amino acid sequence of the purified CFA/III pili coincided with the deduced amino acid sequence for residues 32 to 54 of *CofA*. Western blot (immunoblot) analysis of *CofA* also indicated its processing to form mature pilin in the presence of the downstream region of *cofA*. These results suggest that the major pilin of CFA/III pili is produced as a precursor form which is posttranslationally modified to the mature pilin and forms morphological pili after cleavage of the Gly-30–Met-31 junction, probably by a protease encoded by an as-yet-unknown gene located downstream of *cofA*. Interestingly, the N-terminal 30-amino-acid sequence of mature CFA/III shows the highest identity (76.7%) to TcpA pilin of *Vibrio cholerae*, which is a type IV class B pilin.

The ability of enterotoxigenic *Escherichia coli* (ETEC), an important cause of diarrhea with a worldwide distribution, to adhere to and colonize intestinal epithelium is an essential step for pathogenicity in addition to the ability to produce heat-labile and/or heat-stable enterotoxins. The colonizing ability of human ETEC depends on the presence of colonization factor antigens (CFAs) on the surface of the cells, which form pili (or fimbriae). Distinct types of CFAs, such as CFA/I, CFA/II, CFA/III, and CFA/IV, have been described for human ETEC strains. CFA/I consists of a single antigen (10), whereas some types such as CFA/II (5, 23, 27, 37) and CFA/IV (25, 40) consist of a complex of different antigens named *E. coli* surface (CS) antigens. Cloning and sequencing of genes encoding CFA/I (15, 21), CS1 and CS3 of CFA/II (2, 31), and CS5 of CFA/IV (4, 20) have been reported previously. Although CFA/I- and CFA/II-carrying ETEC seem to be the most prevalent, a wide variation in the prevalence of ETEC strains harboring CFAs in different parts of the world has been reported (5, 9, 10, 14, 19). According to our survey (19), 8% of ETEC strains isolated from patients with traveler's diarrhea in Japan were found to carry CFA/III pili. However, no information on CFA/III genetic determinants has yet been reported. We and colleagues have previously cloned the gene necessary for expressing the CFA/III pili (35). In the present study, we report the sequencing of the gene encoding CFA/III pilin and evidence that the CFA/III pilus is related to type IV pili, especially the class B pili as defined in reference 13.

A 55-kb plasmid controlling the expression of CFA/III was isolated from *E. coli* 260-1 (18, 19) after it was marked with ampicillin-resistant transposon Tn3 (35). The 17.4-kb region of

the Tn3-marked plasmid pSH1001 responsible for CFA/III formation was determined as described previously (35). The partially overlapping 11.5- and 12.4-kb fragments of the region have been separately cloned to compatible plasmid vectors, resulting in pTT202 and pTT206, respectively (35). The simultaneous presence of pTT202 and pTT206 in *E. coli* was necessary for cell agglutination with anti-CFA/III antiserum (19) and pilus formation. Cells were grown on CFA agar plates at 37°C for CFA/III production (18). Long, straight pili were observed only on *E. coli* harboring both pTT202 and pTT206 but not on *E. coli* harboring two cloning vectors (pACYC 184 and pMW119) by electron microscopic examination (19). To determine which region is responsible for production of the major pilin, Western blot (immunoblot) analysis of the cell extracts was carried out (Fig. 1). *E. coli* HB101 harboring both pTT202 and pTT206 produced CFA/III with a molecular mass of 20.5 kDa which was identical to the purified CFA/III. No CFA/III was observed in *E. coli* harboring only pTT206. *E. coli* carrying only pTT202, however, produced a cross-reacting material with an apparent molecular mass of 26.5 kDa although no pilus formation was observed on the cells. These results suggest that the structural gene for CFA/III pilin is located on pTT202 and that CFA/III pilin may be produced as a 26.5-kDa precursor and then proteolytically processed to form the 20.5-kDa mature pilin.

Thus, we further subcloned the fragments of pTT202 into vector plasmid pMW119 (42) and the products of *E. coli* harboring a series of clones were analyzed by Western blotting (41) with CFA/III antiserum (19). As shown in Fig. 2, the cell extracts of pTT201, pTT217, and pTT213, all of which shared the 1.1-kb *EcoRI-SalI* fragment, produced 26.5-kDa antigens but the clones (pTT210 and pTT220) which did not have the fragment produced no antigen. This suggests that the struc-

* Corresponding author. Phone: 81-6-879-8276. Fax: 81-6-879-8277.




FIG. 1. Western blotting analysis of *E. coli* extracts using anti-CFA/III rabbit antiserum. Lane 1, purified CFA/III; lane 2, wild-type strain (260-1); lane 3, HB101 harboring pTT206 and pTT202; lane 4, HB101 harboring pTT202; lane 5, HB101 harboring pTT206; lane 6, HB101. The symbols below the gel (CFA/III) represent the results of bacterial slide agglutination tests with CFA/III antiserum. The 20.5- and 26.5-kDa bands are indicated (arrowheads).

tural gene for the pilin subunit is located in the 1.1-kb *EcoRI-SalI* region.

From these findings, we determined the DNA sequence of the 1.4-kb *AvaI-SalI* fragment of pTT213 (Fig. 2). The DNA fragment obtained after subcloning was finally cloned into the M13mp18 vector (43) and then digested by exonuclease III to generate DNA fragments of various lengths (16). The nucleotide sequences of cloned fragments were determined by the dideoxy chain termination method using a commercial DNA Sequencing Kit (Takara Shuzo Co., Kyoto, Japan) (33). Sequence analysis revealed a 714-bp open reading frame (ORF), ORF1 (Fig. 2). In ORF1, a Shine-Dalgarno sequence (36) in the 10-bp region upstream of the start codon and a transcriptional terminator-like inverted repeat sequence (32) in the 10-bp region downstream of the stop codon were observed (Fig. 3). ORF1 encodes 238 amino acids with a calculated molecular weight of 25,309, which agrees well with the size of




FIG. 2. Subcloning of the gene encoding CFA/III pilin from pTT202 and strategy for DNA sequencing of pTT213. General cloning techniques, digestion of DNA with restriction enzyme, gel electrophoresis, ligation, and transformation were performed as previously described (3, 35). Results of Western blotting analysis are shown on the right. The arrows represent sequenced fragments and their direction of reading. S, *SalI*; E, *EcoRI*; B, *BamHI*; H, *HindIII*; P, *PstI*; A, *AvaI*; C, *Clal*.

the precursor-like protein (26.5 kDa) identified by Western blot analysis (Fig. 1). Furthermore, the sequence of the 23 amino acids beginning at Ser-32 was completely identical to that obtained sequencing data (the N-terminal 23 amino acids) of the purified 20.5-kDa mature pilin protein (39), indicating that ORF1 is a structural gene (designated *cofA*) of the major pilin of CFA/III. We found that the sequence of the 30 amino acids from the N terminus of the mature pilin had the highest identity (76.7%) to the sequence of TcpA pili of *Vibrio cholerae* (11, 34) by a sequence homology search and the pili were highly homologous to other type IV pili expressed in several other organisms (Fig. 4) (8, 11, 12, 17, 24, 26, 28). No significant homology of the amino acid sequence in the residual region of CFA/III to those of type IV pili was observed. Similar observations of various type IV pili have been reported previously (8, 11, 12, 17, 24, 26, 28).

Several distinct types of pilus colonization factors for human and animal ETEC have been described. Among these, K88 and K99 found in animal ETEC have been intensively characterized at the molecular level (1, 6, 29, 30). Genes for K88 and K99 have been demonstrated to consist of operons, and each gene product has also been characterized in detail (1, 6, 29, 30). On the other hand, genetic information regarding CFAs of human ETEC is still limited. Only the structural genes for the major pilin subunit of CFA/I and for CS1, CS3, and CS5 have been reported (2, 4, 15, 20, 21, 31).

In the present paper, we describe the complete nucleotide sequence of the structural gene for CFA/III pilin. In our experience with subcloning, construction of the mature pili (morphological pilus formation on *E. coli*) required a large (21-kb) DNA fragment (*Clal*₁-*Bam*HI₂) (35), suggesting the requirement of a complex of genes, probably operons. Judging from our present results, there may be a gene for a protease which cleaves (or processes) a precursor of CFA/III pilin (238 amino acids, 26.5 kDa) to the mature pilin (20.5 kDa). Comparison of the amino acid sequences of the precursor CFA/III pilin and mature pilin (39) suggests a cleavage of the precursor between Gly-30 and Met-31.

Interestingly, the N-terminal 30-amino-acid sequence of the mature CFA/III pilin is highly hydrophobic and has homology to those of type IV pili such as TcpA of *V. cholerae* O395 (11, 34) and the pili of *Neisseria gonorrhoeae* (MS11) (26), *Pseudomonas aeruginosa* PAK (28), *Moraxella bovis* EPP63 (24) and *Bacteroides nodosus* 265 (8). Overall identity between amino acid sequences of CFA/III and sequences of these type 4 pilins was about 30%. Among these, the highest identity (76.7%) was found between CFA/III pilin and TcpA pilin, which is believed to be an important colonization factor of *V. cholerae* (22). Not only amino acid sequence homology but also morphological similarities (long and straight pili) and functional similarities (attachment to intestinal cells [18]) between CFA/III and type IV pili (TcpA) were observed. In addition to these similarities CofA was processed to form the mature pilin (CFA/III) only in the presence of the downstream region of *cofA*. Processing of the TcpA precursor to the mature pilin is believed to be carried out by a protease encoded in the downstream region of *tcpA* (22), suggesting that the genes for CFA/III and TcpA pili may be evolutionarily closely related.

Along with ORF1 (*cofA*), another ORF (ORF2) was observed in the 61-bp region downstream of ORF1, although the entire sequence of ORF2 has not yet been cloned (Fig. 2). A Shine-Dalgarno sequence (GGAG) was observed 7 bp upstream of ORF2, and some similarity (about 20 to 30%) of the deduced amino acid sequence of ORF2 to the accessory fimbrial proteins of *V. cholerae* (11, 34), *B. nodosus* (17), and *M. bovis* (12) was observed at least in the N-terminal 113-amino-

	<u>AvaII</u>		
1	GGTCCTATTTTAAATAATTATTGAGCCATCGGTGATGCTCCTTGATGGTTTCAAATTGTAAGATATTGTCATTGGTATGT		79
80	TTTGATGGAAATATTACCAATGCAGCATCGAAAACAACGGAGGGCATCAGGTAGTACTGGATGTGGCTTTTCTCATAGGA		158
159	GTGATATATATCATGGTCTCTCTGGCTGTATCCGGTTGTTTATGATGAATACATATTATGGTTTCATGACCTATTTA		237
	<u>EcoRI</u>		
238	ATTTAATTATCGTAATTAATTGTAGATGAATTCAACAGGAGGGAAGTTTCA	ATG CTT TCG GTT TAT AAC AGA	309
	S.D.	Met Leu Ser Val Tyr Asn Arg	
310	ACG CAA AAA ATG AAA GAA GAG GCA AGA AAA AAA CTG GCC AAG TAT CAT GAA TTA CGT AAA		369
	Thr Gln Lys Met Lys Glu Glu Ala Arg Lys Lys Leu Ala Lys Tyr His Glu Leu Arg Lys		
370	CAG CGA GGT ATG AGC CTT CTG GAA GTC ATC ATC GTT CTG GGG ATT ATC GGA ACA ATT GCT		429
	Gln Arg Gly Met Ser Leu Leu Glu Val Ile Ile Val Leu Gly Ile Ile Gly Thr Ile Ala		
430	GCG GGT GTG GTG ATT CTG GCT CAA CGA GCA TTT GAC TCA CGT ACT GTT TCT GAA TTG GTC		489
	Ala Gly Val Val Ile Leu Ala Gln Arg Ala Phe Asp Ser Arg Thr Val Ser Glu Leu Val		
490	ACT AAT ACG AAT ACT ATT CGT GTT GCG ATG AAA GAT GCT TAT CAG CGT GAC GGT AAG TAT		549
	Thr Asn Thr Asn Thr Ile Arg Val Ala Met Lys Asp Ala Tyr Gln Arg Asp Gly Lys Tyr		
550	CCG GAT TAT CAA GCT CCA TTA AGT CTT ACT GCT GAT TCA ATT AAA ACA GAT TCA ACA GGT		609
	Pro Asp Tyr Gln Ala Pro Leu Ser Leu Thr Ala Asp Ser Ile Lys Thr Asp Ser Thr Gly		
610	ATA GCG GTT GCG CAG TTA GTC CAA TTA GGG AAA CTA ACC CCT GAT GAA GCC CGA AAT GGT		669
	Ile Ala Val Ala Gln Leu Val Gln Leu Gly Lys Leu Thr Pro Asp Glu Ala Arg Asn Gly		
670	ATT TCT GGG GAC TAT ATT GGT ATT GGT GGT GCA ATA ACA TCT TCA GGT TCT ACA ATC AAC		729
	Ile Ser Gly Asp Tyr Ile Gly Ile Gly Gly Ala Ile Thr Ser Ser Gly Ser Thr Ile Asn		
730	AAG GGA TTT GCA ATG GAA CTG AAC GGA CTT AGC CAA GAG CAA TGT CGT TCA ATT CTT GGA		789
	Lys Gly Phe Ala Met Glu Leu Asn Gly Leu Ser Gln Glu Gln Cys Arg Ser Ile Leu Gly		
790	CAA GTT GGT GAT AAC TGG GAG TAT GTG GCA GTT GGT ACT AGT CCT TCT GGT TCT TAT GAT		849
	Gln Val Gly Asp Asn Trp Glu Tyr Val Ala Val Gly Thr Ser Pro Ser Gly Ser Tyr Asp		
	<u>PstI</u>		
850	GCT CTG TCT GCA GGC GCA GTA AAC ATG CTG GCT GCT ACT GAT AAT ACT ACA ATA TTA CGT		909
	Ala Leu Ser Ala Gly Ala Val Asn Met Leu Ala Ala Thr Asp Asn Thr Thr Ile Leu Arg		
910	AGC CTG GCG GCT AAT GGT CAA GTA TCA CTG ACA GCT GAG AAA ATT TTA AAA ACC TGC ACA		969
	Ser Leu Ala Ala Asn Gly Gln Val Ser Leu Thr Ala Glu Lys Ile Leu Lys Thr Cys Thr		
970	GCC ACA GTT AAC TCT ATT ACT TTG GCG AGC CGT TAA TAAGATATTTAAATACAGGTCCTATTCATG		1036
	Ala Thr Val Asn Ser Ile Thr Leu Ala Ser Arg ***		
	<u>AvaII</u>		
1037	GACCTGTATTTACGTGCCGGGAGTTCTTT	ATG AAT ATG AGG GGT TTC ACG CTT CTG GAA ATG ATT	1101
	S.D.	Met Asn Met Arg Gly Phe Thr Leu Leu Glu Met Ile	
1102	GTT ACT CTG GCT GTT ATG GGA GTT GCA ATG TTA TCT GTC ATT AAA TAT AAA GAG AAA GAA		1161
	Val Thr Leu Ala Val Met Gly Val Ala Met Leu Ser Val Ile Lys Tyr Lys Glu Lys Glu		
1162	GCA GAT GAA GCC AGA CGA CAA ATT GTA TCT AAT GCT CTG ATT TCA GAA ATC GCC GGC ATT		1221
	Ala Asp Glu Ala Arg Arg Gln Ile Val Ser Asn Ala Leu Ile Ser Glu Ile Ala Gly Ile		
1222	GTG GAT TTT GTC GCA GAG GAA CAA ATA ACC GTT ATA GAA CAG GGA ATA GAA AAA GAA ATT		1281
	Val Asp Phe Val Ala Glu Glu Gln Ile Thr Val Ile Glu Gln Gly Ile Glu Lys Glu Ile		
1282	ACG AAT CCA CTT TAT GAG CAG AGC TCT GGG ATT CCA TAT ATA AAT CGA ACT ACA AAT AAA		1341
	Thr Asn Pro Leu Tyr Glu Gln Ser Ser Gly Ile Pro Tyr Ile Asn Arg Thr Thr Asn Lys		
1342	GAT TTA AAC TCA ACT ATG TCA ACA AAT GCC TCT GAG TTT ATT AAT TGG GGG GCT GGT ACG		1401
	Asp Leu Asn Ser Thr Met Ser Thr Asn Ala Ser Glu Phe Ile Asn Trp Gly Ala Gly Thr		
	<u>SalI</u>		
1402	TCG AC		1406
	Ser		

FIG. 3. Nucleotide and amino acid sequencing of the gene encoding CFA/III pilin. The underlined Shine-Dalgarno sequences (S.D.) and the proteolytic site (vertical arrow) are indicated. Inverted repeat sequences and restriction enzyme-recognizing sequences are represented by horizontal arrows under and lines above the nucleotides, respectively. The nucleotide sequence numbers are shown in the margins.

8. **Elleman, T. C., P. A. Hoyne, N. M. McKern, and D. J. Stewart.** 1986. Nucleotide sequence of the gene encoding the two-subunit pilin of *Bacteroides nodosus* 265. *J. Bacteriol.* **167**:243–250.
9. **Evans, D. G., D. J. Evans, Jr., and W. Tjoa.** 1977. Hemagglutination of human group A erythrocytes by enterotoxigenic *Escherichia coli* isolated from adults with diarrhea: correlation with colonization factor. *Infect. Immun.* **18**:330–337.
10. **Evans, D. G., R. P. Silver, D. J. Evans, Jr., D. G. Chase, and S. L. Gorbach.** 1975. Plasmid-controlled colonization factor associated with virulence in *Escherichia coli* enterotoxigenic for humans. *Infect. Immun.* **12**:656–667.
11. **Faast, R., M. A. Ogierman, U. H. Strocher, and P. A. Manning.** 1989. Nucleotide sequence of the structural gene, *tcpA*, for a major pilin subunit of *Vibrio cholerae*. *Gene* **85**:227–231.
12. **Fulks, K. A., C. F. Marrs, S. P. Stevens, and M. R. Green.** 1990. Sequence analysis of the inversion region containing the pilin genes of *Moraxella bovis*. *J. Bacteriol.* **172**:310–316.
13. **Giron, J. A., M. M. Levine, and J. B. Kaper.** 1994. Longus: a long pilus ultrastructure produced by human enterotoxigenic *Escherichia coli*. *Mol. Microbiol.* **12**:71–82.
14. **Guth, B. E. C., E. G. Aguiar, P. M. Griffin, S. R. T. S. Ramos, and T. A. T. Gomes.** 1994. Prevalence of colonization factor antigens (CFAs) and adherence to HeLa cells in enterotoxigenic *Escherichia coli* isolated from feces of children in Sao Paulo. *Microbiol. Immunol.* **38**:695–701.
15. **Hamers, A. M., H. J. Pel, G. A. Willshaw, J. G. Kusters, B. A. M. van der Zeist, and W. Gaastra.** 1989. The nucleotide sequence of the first two genes of the CFA/I fimbrial operon of human enterotoxigenic *Escherichia coli*. *Microb. Pathog.* **6**:297–309.
16. **Henikoff, S.** 1984. Unidirectional digestion with exonuclease III creates targeted breakpoints for DNA sequencing. *Gene* **28**:351–359.
17. **Hobbs, M., B. P. Dalrymple, P. T. Cox, S. P. Livingstone, S. F. Delaney, and J. S. Mattick.** 1991. Organization of the fimbrial gene region of *Bacteroides nodosus*: class I and class II strains. *Mol. Microbiol.* **5**:543–560.
18. **Honda, T., M. Arita, and T. Miwatani.** 1984. Characterization of new hydrophobic pili of human enterotoxigenic *Escherichia coli*: a possible new colonization factor. *Infect. Immun.* **43**:959–965.
19. **Honda, T., N. Wetprasit, M. Arita, and T. Miwatani.** 1989. Production and characterization of monoclonal antibodies to a pilus colonization factor (colonization factor antigen III) of human enterotoxigenic *Escherichia coli*. *Infect. Immun.* **57**:3452–3457.
20. **Heuzenroeder, M. W., B. L. Neal, C. J. Thomas, R. Halter, and P. A. Manning.** 1989. Characterization and molecular cloning of the PCF8775 CS5 antigen from an enterotoxigenic *Escherichia coli* O115:H40 isolated in Central Australia. *Mol. Microbiol.* **3**:303–310.
21. **Karjalainen, T. K., D. G. Evans, M. So, and C.-H. Lee.** 1989. Molecular cloning and nucleotide sequence of the colonization factor antigen I gene of *Escherichia coli*. *Infect. Immun.* **57**:1126–1130.
22. **Kaufman, M. R., and R. K. Taylor.** 1994. The toxin-coregulated pilus: biogenesis and function, p. 187–202. *In* I. K. Wachsmuth, P. A. Blake, and Ø. Olsvik (ed.), *Vibrio cholerae* and cholera. American Society for Microbiology, Washington, D.C.
23. **Levine, M. M., P. Ristaino, G. Marley, C. Smyth, S. Knutton, E. Boedeker, R. Black, C. Young, M. L. Clements, C. Cheney, and R. Patnaik.** 1984. Coli surface antigens 1 and 3 of colonization factor antigen II-positive enterotoxigenic *Escherichia coli*: morphology, purification, and immune responses in humans. *Infect. Immun.* **44**:409–420.
24. **Marrs, C. F., G. Schoolnik, J. M. Koomey, J. Hardy, J. Rothbard, and S. Falkow.** 1985. Cloning and sequencing of a *Moraxella bovis* pilin gene. *J. Bacteriol.* **163**:132–139.
25. **McConnell, M. M., L. V. Thomas, G. A. Willshaw, H. R. Smith, and B. Rowe.** 1988. Genetic control and properties of coli surface antigens of colonization factor antigen IV (PCF8775) of enterotoxigenic *Escherichia coli*. *Infect. Immun.* **56**:1974–1980.
26. **Meyer, T. F., E. Billyard, R. Haas, S. Strozbach, and M. So.** 1984. Pilus genes of *Neisseria gonorrhoeae*: chromosomal organization and DNA sequence. *Proc. Natl. Acad. Sci. USA* **81**:6110–6114.
27. **Mullany, P., A. M. Field, M. M. McConnell, S. M. Scotland, H. R. Smith, and B. Rowe.** 1983. Expression of plasmids coding for colonization factor antigen II (CFA/II) and enterotoxin production in *Escherichia coli*. *J. Gen. Microbiol.* **129**:3591–3601.
28. **Pasloske, B. L., B. B. Finlay, and W. Paranchych.** 1985. Cloning and sequencing of the *Pseudomonas aeruginosa* PAK pilin gene. *FEBS Lett.* **183**:408–412.
29. **Pedersen, P. A.** 1991. Structure-function analysis of the K88ab fimbrial subunit protein from porcine enterotoxigenic *Escherichia coli*. *Mol. Microbiol.* **5**:1073–1080.
30. **Pedersen, P. A., and L. N. Andersen.** 1991. Deletion and duplication of specific sequences in the K88ab fimbrial subunit protein from porcine enterotoxigenic *Escherichia coli*. *Mol. Gen. Genet.* **229**:285–291.
31. **Perez-Casal, J., J. S. Swartley, and J. P. Scott.** 1990. Gene encoding the major subunit of CS1 pili of human enterotoxigenic *Escherichia coli*. *Infect. Immun.* **58**:3594–3600.
32. **Rosenberg, M., and D. Court.** 1979. Regulatory sequences involved in the promotion and termination of RNA transcription. *Annu. Rev. Genet.* **13**:319–353.
33. **Sanger, F., S. Nicklen, and A. R. Coulson.** 1977. DNA sequencing with chain-terminating inhibitors. *Proc. Natl. Acad. Sci. USA* **74**:5463–5467.
34. **Shaw, C. E., and R. K. Taylor.** 1990. *Vibrio cholerae* O395 *tcpA* pilin gene sequence and comparison of predicted protein structural features to those of type 4 pilins. *Infect. Immun.* **58**:3042–3049.
35. **Shinagawa, H., T. Taniguchi, O. Yamaguchi, K. Yamamoto, and T. Honda.** 1993. Cloning of the genes that control formation of the fimbrial colonization factor antigen III (CFA/III) from an enterotoxigenic *Escherichia coli*. *Microbiol. Immunol.* **37**:689–694.
36. **Shine, J., and L. Dalgarno.** 1974. The 3'-terminal sequence of *Escherichia coli* 16S ribosomal RNA: complementarity to nonsense triplets and ribosome binding sites. *Proc. Natl. Acad. Sci. USA* **71**:1342–1346.
37. **Smith, C. L.** 1982. Two mannose-resistant haemagglutinins on enterotoxigenic *Escherichia coli* of serotype O6:K15:H16 or H⁻ isolated from traveler's and infantile diarrhoea. *J. Gen. Microbiol.* **128**:2081–2096.
38. **Sohel, I., J. L. Puente, W. J. Murray, J. Vuopio-Varkika, and G. K. Schoolnik.** 1993. Cloning and characterization of the bundle-forming pilin gene of enteropathogenic *Escherichia coli* and its distribution in *Salmonella* serotypes. *Mol. Microbiol.* **7**:563–575.
39. **Taniguchi, T., M. Arita, M. Sato, K. Yamamoto, T. Miwatani, and T. Honda.** 1994. Evidence that N-terminal amino acid sequence of pilus colonization factor antigen III produced by human enterotoxigenic *Escherichia coli* is similar to that of *TcpA* pilin of *Vibrio cholerae*. *J. Infect. Dis.* **170**:1049–1050.
40. **Thomas, L. V., M. M. McConnell, B. Rowe, and A. M. Field.** 1985. The possession of three novel coli surface antigens by enterotoxigenic *Escherichia coli* strains positive for the putative colonization factor PCF8775. *J. Gen. Microbiol.* **131**:2319–2326.
41. **Towbin, H., T. Staehelin, and J. Gordon.** 1979. Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. *Proc. Natl. Acad. Sci. USA* **76**:4350–4354.
42. **Yamaguchi, K., and Y. Masamune.** 1985. Autogenous regulation of synthesis of the replication protein in plasmid pSC101. *Mol. Gen. Genet.* **200**:362–367.
43. **Yanisch-Perron, C., J. Vieira, and J. Messing.** 1985. Improved M13 phage cloning vectors and host strains: nucleotide sequences of the M13mp18 and pUC19 vectors. *Gene* **33**:103–119.