

- 17 **Lau KF**, Miller CC, Anderton BH, Shaw PC. Molecular cloning and characterization of the human glycogen synthase kinase-3beta promoter. *Genomics* 1999;60(2):121–8.
- 18 **Zambrano N**, Minopoli G, de Candia P, Russo T. The Fe65 adaptor protein interacts through its PID1 domain with the transcription factor CP2/LSF/LBP1. *J Biol Chem* 1998;273(32):20128–33.
- 19 **Cao X**, Sudhof TC. A transcriptionally [correction of transcriptionally] active complex of APP with Fe65 and histone acetyltransferase Tip60. *Science* 2001;293(5527):115–20.
- 20 **Blacker D**, Albert MS, Bassett SS, Go RC, Harrell LE, Folstein MF. Reliability and validity of NINCDS-ADRDA criteria for Alzheimer's disease. The National Institute of Mental Health Genetics Initiative. *Arch Neurol* 1994;51(12):1198–204.
- 21 **Blacker D**, Bertram L, Saunders AJ, Moscarillo TJ, Albert MS, Wiener H, Perry RT, Collins JS, Harrell LE, Go RC, Mahoney A, Beaty T, Fallin MD, Avramopoulos D, Chase GA, Folstein MF, McNinis MG, Bassett SS, Doheny KJ, Pugh EW, Tanzi RE. Results of a high-resolution genome screen of 437 Alzheimer's disease families. *Hum Mol Genet* 2003;12(1):23–32.
- 22 **Mullin K**, Bertram L, Moscarillo TJ, Becker KD, Wang C, Growdon J, Blacker D, Tanzi RE. Genetic association within the IDE region on chromosome 10 - The Consortium on Alzheimer's Genetics (CAG). Poster [abstract no. 202.8] presented at the 33rd Annual Meeting of the Society for Neuroscience, New Orleans, LA, 2003.
- 23 **Rabinowitz D**, Laird N. A unified approach to adjusting association tests for population admixture with arbitrary pedigree structure and arbitrary missing marker information. *Hum Hered* 2000;50(4):211–23.
- 24 **Lake SL**, Blacker D, Laird NM. Family-based tests of association in the presence of linkage. *Am J Hum Genet* 2000;67(6):1515–25.
- 25 **Martin ER**, Monks SA, Warren LL, Kaplan NL. A test for linkage and association in general pedigrees: the pedigree disequilibrium test. *Am J Hum Genet* 2000;67(1):146–54.
- 26 **Martin ER**, Bass MP, Kaplan NL. Correcting for a potential bias in the pedigree disequilibrium test. *Am J Hum Genet* 2001;68(4):1065–7.
- 27 **Witte JS**, Gauderman WJ, Thomas DC. Asymptotic bias and efficiency in case-control studies of candidate genes and gene-environment interactions: basic family designs. *Am J Epidemiol* 1999;149(8):693–705.
- 28 **Siegmund KD**, Langholz B, Kraft P, Thomas DC. Testing linkage disequilibrium in sibships. *Am J Hum Genet* 2000;67(1):244–8.
- 29 **Laird NM**, Mosteller F. Some statistical methods for combining experimental results. *Int J Technol Assess Health Care* 1990;6(1):5–30.
- 30 **DerSimonian R**, Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986;7(3):177–88.
- 31 **Ioannidis JP**, Ntzani EE, Trikalinos TA, Contopoulos-Ioannidis DG. Replication validity of genetic association studies. *Nat Genet* 2001;29(3):306–9.
- 32 **Sanchez-Guerra M**, Combarros O, Infante J, Llorca J, Berciano J, Fontalba A, Fernandez-Luna JL, Pena N, Fernandez-Viadero C. Case-control study and meta-analysis of low density lipoprotein receptor-related protein gene exon 3 polymorphism in Alzheimer's disease. *Neurosci Lett* 2001;316(1):17–20.
- 33 **Kolsch H**, Piok U, Mohamed I, Schmitz S, Rao ML, Maier W, Heun R. Association of the C766T polymorphism of the low-density lipoprotein receptor-related protein gene with Alzheimer's disease. *Am J Med Genet* 2003;121B(1):128–30.
- 34 **Perry RT**, Collins JS, Wiener H, Acton R, Go RC. The role of TNF and its receptors in Alzheimer's disease. *Neurobiol Aging* 2001;22(6):873–83.
- 35 **Alvarez V**, Mata IF, Gonzalez P, Lahoz CH, Martinez C, Pena J, Guisasola LM, Coto E. Association between the TNFalpha-308 A/G polymorphism and the onset-age of Alzheimer disease. *Am J Med Genet* 2002;114(5):574–7.
- 36 **Lehmann DJ**, Johnston C, Smith AD. Synergy between the genes for butyrylcholinesterase K variant and apolipoprotein E4 in late-onset confirmed Alzheimer's disease. *Hum Mol Genet* 1997;6(11):1933–6.
- 37 **Hiltunen M**, Mannermaa A, Helisalmi S, Koivisto A, Lehtovirta M, Ryyanen M, Riekkonen P Sr, Soiniinen H. Butyrylcholinesterase K variant and apolipoprotein E4 genes do not act in synergy in Finnish late-onset Alzheimer's disease patients. *Neurosci Lett* 1998;250(1):69–71.

## CORRECTION

doi: 10.1136/jmg.2005.26138corr1

In the paper titled, A new syndrome, congenital extraocular muscle fibrosis with ulnar hand anomalies, maps to chromosome 21qter (*J Med Genet* 2004;42:408–15) there are a number of errors. The affiliations and correspondence details were incorrect, the correct details have been listed below:

**T Tukel, R J Desnick\***, Department of Human Genetics, Mount Sinai School of Medicine of New York University, New York, USA

**A Uzumcu, H Kayserili, M Y Apak, O Uyguner, Bernd Wollnik\***, Child Health Institute, Division of Medical Genetics, Istanbul University, Istanbul, Turkey

**A Gezer**, Department of Ophthalmology, Istanbul Faculty of Medicine, Istanbul University

**S H Gultekin**, Department of Pathology, Section of Neuropathology, Mount Sinai School of Medicine of New York University

**H-C Hennies, P Nurnberg**, Gene Mapping Center and Department of Molecular Genetics, Max Delbrück Center for Molecular Genetics, Berlin, Germany

\* Co-senior authors who contributed equally to this research

Correspondence to:

Professor Robert J Desnick, PhD, MD, Department of Human Genetics, Mount Sinai School of Medicine, 1425 Madison Avenue, Box 1498, New York, NY 10029, USA; rjdesnick@mssm.edu

Bernd Wollnik, MD, Institute of Child Health, Division of Medical Genetics, Istanbul University, Millet Caddesi, Capa, 34390 Istanbul, Turkey; wollnik@superonline.com

In figure 3 the marker D21S1400 should be D21S1440.

On page 413 in the second paragraph, the marker D21S1446 should be D21S1446.