


## LETTERS TO THE EDITOR

### Facial diplegia with paraesthesias: facial nerve enhancement in three dimensional MRI

Facial diplegia (bilateral facial paralysis) is a rare clinical finding that can be the presenting symptom in a wide range of diseases.<sup>1</sup> It occurs in about 50% of patients with Guillain-Barré syndrome (GBS). Guillain-Barré syndrome causes regional and functional variants with unusual features. Ropper described four patients with facial diplegia and distal limb paraesthesias, and he defined them as having a rare variant form of GBS because of shared clinical, electrophysiological, and CSF features.<sup>2</sup> The aetiology and nosological position of facial diplegia presenting in this variant form is still controversial. We experienced a patient who had bilateral facial paralysis, distal limb paraesthesias, and diminished reflexes whose contrast enhanced three dimensional MRI (3-D MRI) showed enhancing lesions in the bilateral facial nerves.

A 27 year old woman had nasal discharge and coughing. One week later she noticed paraesthesias in her fingers and toes. Nine days after the onset of her neurological symptom, she developed bilateral facial weakness. On admission (day 12) she showed moderate, bilateral facial paralysis that caused her difficulty in moving her forehead, in approximating her eyelids, and in lifting the corners of her mouth. Tearing and taste were preserved. The other cranial nerves were normal. A motor examination showed normal strength in her limbs. Superficial and deep senses were normal even though she had distal limb paraesthesias. Deep tendon reflexes were absent in all her limbs and her plantar responses were flexor type. Cerebellar ataxia and autonomic nervous dysfunction were excluded. Chest radiography was normal. Laboratory studies of the identifiable causes of facial diplegia (sarcoidosis, Lyme disease, syphilis, infectious mononucleosis, herpes simplex virus, diabetes mellitus, and connective tissue disease) were all negative. On day 12, the CSF examination detected mild increases in protein concentration (57 mg/dl) without pleocytosis. The blink reflex was elicited and both the R1 and R2 components were reduced, but their latencies were not prolonged on day 15. Motor and sensory nerve conduction velocities, and median and tibial nerve responses were all normal on day 18. Auditory brainstem responses were normal. On Day 19, when her facial diplegia was moderate, conventional brain MRI detected no abnormality. A contrast enhanced 3-D MRI, which was obtained by spoiled gradient recalled acquisition in the steady state sequence using a 1.5 tesla system after injection of gadolinium-diethylenetriamine pentaacetic acid (Gd-DTPA), was performed. The intracranial segments of the bilateral facial nerves were remarkably enhanced by Gd-DTPA (figure). On Day 45, when her symptoms were no longer present, there were no abnormal enhancements of her facial nerves.

After showing symptoms of upper respiratory infection, the patient experienced the



(A) Before and (B) after contrast enhanced transverse reformatted three dimensional MRI. The lateral intracranial segments of both facial nerves are enhanced (arrows) in B.

acute onset of facial diplegia, distal limb paraesthesias, and areflexia but no other neurological deficits. The CSF examination showed albuminocytological dissociation, and clinical and laboratory examinations excluded the possibility of viral or bacterial infection, Lyme disease, tumour, sarcoidosis, cerebrovascular disease, diabetes mellitus, bilateral Bell's palsy, and congenital and familial disorders. The patient's illness followed a monophasic course. We therefore diagnosed this case as having "facial diplegia with paraesthesias", which should be included for the differential diagnosis whenever sudden bilateral facial paresis occurs. Routine brain MRI showed no abnormalities, whereas contrast enhanced 3-D MRI showed Gd enhancement of the bilateral facial nerves. The MRI findings indicate the involvement of the peripheral facial nerves in our patient. Fulbright *et al*<sup>3</sup> reported an additional case of GBS with multiple cranial nerve enhancements seen on Gd enhanced MRI. The mechanism of abnormal enhancement of the cranial nerves in the patients with GBS is not entirely understood; however, it is widely regarded as disruption of the blood nerve barrier by the inflammatory infiltrate. Ramsey *et al*<sup>4</sup> evaluated the MRI findings obtained with Gd contrast enhancement in five patients who had acute bilateral facial paralysis: GBS (n=1), herpes simplex polyneuritis (n=1), meningeal lymphoma (n=1), and bilateral Bell's palsy (n=2). Gd enhanced MRI has been shown to be the procedure of choice for demonstrating inflammatory lesions of the facial nerves. Nagaoka *et al*<sup>5</sup> showed oculomotor nerve enhancement on 3-D MRI in Fisher's syndrome, the best known variant of GBS. Ours is the first report of facial nerve enhancement in "facial diplegia with paraesthesias". These findings suggest that 3-D MRI with Gd-DTPA can be used to identify inflammatory conditions that produce peripheral lesion of the cranial nerves in GBS variants.

AKINORI HOZUMI  
NOBUHIRO YUKI  
KAORU YAMAZAKI  
KOICHI HIRATA

Department of Neurology, Dokkyo University School of Medicine, Tochigi, Japan

Correspondence to: Dr Akinori Hozumi, Department of Neurology, Dokkyo University School of Medicine, Kitakobayashi 880, Mibu, Shimotsuga, Tochigi 321-0293, Japan. Telephone +81 282 86 1111 ext 2720; fax +81 282 86 5884; email ahozumi@ibm.net

- 1 Keane JR. Bilateral seventh nerve palsy: analysis of 43 cases and review of the literature. *Neurology* 1994;44:1198-202.
- 2 Ropper AH. Further regional variants of acute immune polyneuropathy. *Arch Neurol* 1994;51:671-5.
- 3 Fulbright RK, Erdum E, Sze G, *et al*. Cranial nerve enhancement in the Guillain-Barré syndrome. *Am J Neuroradiol* 1995;16:923-5.
- 4 Ramsey KL, Kaseff LG. Role of magnetic resonance imaging in the diagnosis of bilateral facial paralysis. *Am J Otol* 1993;14:605-9.
- 5 Nagaoka U, Kato T, Kurita K, *et al*. Cranial nerve enhancement on three-dimensional MRI in Miller Fisher syndrome. *Neurology* 1996;47:1601-2.

### Restless legs syndrome associated with spinal cord lesions

Restless legs syndrome may be either a primary or a secondary disorder. The primary form of the syndrome is often familial whereas the secondary form is mainly associated with uraemia, iron deficiency, or pregnancy. Almost all patients with restless legs syndrome show periodic leg movements during sleep.<sup>1</sup> The pathogenesis of both restless legs syndrome and periodic leg movements is still speculative. Yokota *et al* have reported an association of periodic leg movements with spinal cord lesions.<sup>2</sup> However, none of these patients had the typical clinical features of restless legs syndrome. Restless legs syndrome associated with myelopathy is documented in one patient with a *Borrelia* induced myelitis.<sup>3</sup> We report three patients who developed a restless legs syndrome in close temporal association with spinal cord lesions.

Case 1 was a 35 year old woman who presented with a 3 week history of painless restlessness of her left lower leg that occurred only at rest, particularly in the evening and at night. The restlessness was partially relieved by walking. With the onset of these symptoms, the patient had noticed a numbness of her left hemibody below the breast. Nine months previously, a numbness of the right leg had subsided spontaneously within 1 week. On examination, the patient had decreased senses for touch, pain, and temperature over the left hemibody below the T6 dermatome. Examination of CSF disclosed a normal cell count, normal protein content, increased CNS synthesis of IgG, and positive oligoclonal bands. In MRI studies of the spinal cord, no abnormality was detected. Brain MRI was not performed. Transcranial magnetic stimulation showed a slightly prolonged central conduction time of motor evoked potentials recorded over the left abductor hallucis muscle. Otherwise, multimodality evoked potentials were normal. A myelitis due to multiple sclerosis accompanied by a symptomatic unilateral restless legs syndrome was diagnosed. The patient was treated with 500 mg prednisolone intravenously over 5 days without any clinical effect. However, a single dose of 100 mg levodopa plus benserazide led to a dramatic improvement of the restless legs syndrome. The levodopa treatment was continued and resulted in complete relief.

Case 2 was a 49 year old man who had a traumatic atlantoaxial dislocation that necessitated operative stabilisation of the cervical spine. Preoperative MRI studies had shown a compression of the medulla and the cervical cord. When we saw the patient 3 years later, he complained of a sensation of cold, pain, and restlessness in both lower legs that was present only at rest, particularly in the

evening, and was relieved by walking around and rubbing the legs with cold water. The onset of these symptoms was only a few weeks after the accident. Clinical examination disclosed a mild dysarthrophonia and atrophic pareses of the left sternomastoid and the left trapezius muscle. Tendon reflexes were hyperactive and plantar responses were extensor bilaterally. The patient's gait was spastic, but he was able to walk unassisted. In both legs, pain and temperature sensation were markedly reduced. The diagnosis of a restless legs syndrome secondary to a traumatic lesion of the medulla and the cervical cord was made. Treatment with 100 mg levodopa plus benserazide and 100 mg tramadol resulted in a satisfactory relief of the restless legs syndrome.

Case 3 was a 65 year old man who developed slowly progressive spastic tetraparesis and ascending sensory disturbances in both legs. An MRI study showed a cervical spondylotic myelopathy at the level C3-C6 and the patient underwent spinal cord decompression. Five years later, he was referred to our hospital because of an intense sensation of restlessness of both legs located in the feet and calves. The restlessness occurred when sitting and lying for more than 20 minutes. It was pronounced at night and improved when he was walking around. These symptoms had started simultaneously with the motor and sensory disturbances due to the cervical spondylotic myelopathy and did not improve postoperatively. On examination, the patient was mildly impaired in carrying out fine motor tasks and his gait was moderately spastic. He had reduced touch and vibration senses in both upper limbs. A restless legs syndrome due to a cervical spondylotic myelopathy was diagnosed. Treatment with pergolide resulted in an excellent control of the restless legs syndrome.

Our patients meet the criteria for the diagnosis of restless legs syndrome.<sup>4</sup> Over a follow up period of at least 6 months, restless legs syndrome symptoms were sufficiently relieved by dopaminergic treatment. The association of myelopathy and restless legs syndrome may be merely coincidental. However, the close temporal relation between the onset of myelopathy and restless legs syndrome strongly suggests that restless legs syndrome was secondary to the spinal cord lesions.

The pathogenesis of restless legs syndrome and periodic leg movements is still speculative. In patients with myelopathy and periodic leg movements, it is hypothesised that a spinal cord lesion may permit the expression of a spinal periodic leg movements generator by interrupting descending inhibitory spinal pathways.<sup>2</sup> Our finding of restless legs syndrome in three patients with myelopathy provides evidence that disinhibition of spinal pathways may also be involved in its pathogenesis.

In patient 1, restless legs syndrome was strictly confined to the left leg. Preceding transitory sensory disturbances of the right leg and CSF findings support the diagnosis of multiple sclerosis in this patient. Clinical findings suggest a spinal lesion at the thoracic level. Involvement above the spinal level cannot be excluded. However, clinically and neurophysiologically no supraspinal lesion was detected. Yokota *et al* described three cases of periodic leg movements associated with spinal lesions due to multiple sclerosis.<sup>2</sup> Ferini-Strambi *et al* performed polysomnographic studies in 25 patients with multiple

sclerosis and in an age and sex matched control group.<sup>5</sup> The prevalence of periodic leg movements was significantly higher in the multiple sclerosis group (36% *v* 8%). Patients with multiple sclerosis with periodic leg movements had higher MRI lesion loads in infratentorial regions compared with patients with multiple sclerosis without periodic leg movements. However, spinal MRI was not done and clinical findings were not reported in detail. Thus, further studies are needed to elucidate the prevalence and the pathogenesis of restless legs syndrome and periodic leg movements in patients with multiple sclerosis.

In conclusion, our report suggests that restless legs syndrome may occur secondary to spinal cord lesions due to different causative diseases including multiple sclerosis, spinal cord injury, and cervical spondylotic myelopathy. Similar to idiopathic restless legs syndrome and other secondary forms, restless legs syndrome due to myelopathy may respond well to dopaminergic drugs.

MATHIAS HARTMANN  
ROBERT PFISTER  
KARL PFADENHAUER

Department of Neurology, Zentralklinikum Augsburg,  
Augsburg, Germany

Correspondence to: Dr M Hartmann, Department of Neurology, Zentralklinikum Augsburg, Stenglinstrasse 2, 86156 Augsburg, Germany. Telephone 0049 821 400 2991; fax 0049 821 400 2691.

- 1 Lugaresi E, Cirignotta F, Coccagna G, *et al*. Nocturnal myoclonus and restless legs syndrome. *Adv Neurol* 1986;43:295-307.
- 2 Yokota T, Hirose K, Tanabe H, *et al*. Sleep-related periodic leg movements (nocturnal myoclonus) due to spinal cord lesion. *J Neurol Sci* 1991;104:13-18.
- 3 Hemmer B, Riemann D, Glocker FX, *et al*. Restless legs syndrome after a Borrelia-induced myelitis. *Mov Disord* 1995;10:521-2.
- 4 Walters AS, The International Restless Legs Syndrome Study Group. Toward a better definition of the restless legs syndrome. *Mov Disord* 1995;10:634-42.
- 5 Ferini-Strambi L, Filippi M, Martinelli V, *et al*. Nocturnal sleep study in multiple sclerosis: correlations with clinical and brain magnetic resonance imaging findings. *J Neurol Sci* 1994; 125:194-7.

### Coma in thrombotic thrombocytopenic purpura

Patients with thrombotic thrombocytopenic purpura (TTP) can present with devastating neurological abnormalities.<sup>1</sup> Mortality may be as high as 95%, but current treatment has

reduced this to about 10% and early treatment improves the rate of recovery.<sup>2</sup> We describe two patients who presented with predominantly neurological symptoms and signs who, because of a delay in making a diagnosis of TTP, were referred for treatment at a late stage. Both patients were reviewed by neurological and haematological experts, who considered that the prognosis was poor.

The first case was a 49 year old woman with a longstanding diagnosis of schizophrenia and a previous left sided cerebrovascular accident. She was admitted to her local hospital with a 3 day history of drowsiness, confusion, epistaxes, and spontaneous bruising, having been noted to be increasingly agitated and disoriented over the preceding 6 weeks. Her only medication was trifluoperazine and paroxetine. The second case was a 58 year old man, previously fit and well, who presented to his local hospital with a 3 week history of confusion, drowsiness, jaundice, and right upper quadrant pain. He was taking no medication. The initial findings in both patients are summarised in the table. In both a diagnosis of TTP was made, although this was not until 5 days after admission in the first case, and both patients were transferred to the intensive care unit for plasma exchange and further management.

Treatment was started in both cases with five cycles of plasma exchange, each cycle using 3l cryodepleted fresh frozen plasma, and in the first patient this was followed by a course of oral prednisolone and azathioprine. Both made an excellent recovery, with an improvement in conscious level, a rise in platelet count, disappearance of red cell fragments, a fall in LDH and bilirubin concentrations, and normalisation of renal function. The first patient was self ventilating with no neurological deficit at time of transfer back to the referring hospital. The second patient had a Glasgow coma score of 15 by the fifth day of treatment, the only focal neurology being a bilateral internuclear ophthalmoplegia (INO). Three months later the ophthalmoplegia had resolved and the patient was self caring with minimal disability. Both patients were extensively investigated to look for an underlying cause for TTP, but none was found.

Thrombotic thrombocytopenic purpura is a syndrome comprising a pentad of features—fever, thrombocytopenia, microangiopathic haemolytic anaemia, neurological abnormalities, and renal dysfunction. Not all five features are required to make the

### Summary of patients

	Patient 1	Patient 2
Age (y)	49	58
Sex	F	M
Platelet count (x10 <sup>9</sup> /l)	19	17
Peripheral blood film	RBC fragments +++	RBC fragments +++
Clotting screen+fibrinogen	Normal	Normal
Bilirubin (μmol/l)	67	49
Lactate dehydrogenase (U/l) 1773	931	
Creatinine (μmol/l)	121	131
Glasgow coma score	3/15	3/15
Neurological findings	Pupils equal and reactive, corneal reflexes intact bilaterally; jaw jerk brisk, gag absent; no motor response to painful stimuli; reflexes absent; upgoing plantars	Midposition, non-reactive pupils, disconjugate gaze in all directions; extension of all four limbs to painful stimuli; plantars equivocal
Brain CT	Normal	Raised ICP, nil else.
Brain MRI	Multiple foci of high intensity on the T2 weighted images in white matter of both cerebral hemispheres and poles of lateral ventricles	Normal, although only a scanning MRI was done. A full MRI would have been required to exclude a midbrain lesion, but was not technically possible.