

CORRECTION

Table 2 Comparison of susceptibility and PPNG of *N gonorrhoeae* isolated in Guangzhou during 1996–2001

Year	No	Penicillin			Spectinomycin		Ceftriaxone		Ciprofloxacin			PPNG (%)	
		S (%)	I (%)	R (%)	S (%)	R (%)	S (%)	I (%)	R (%)	S (%)	I (%)		R (%)
1996	201*	17 (8.4)	69 (34.3)	115 (57.2)	199 (99)	2 (1)	120 (62.5)	72 (37.5)	0 (0)	37 (18.6)	127 (63.8)	35 (17.6)	4 (2)
1997	51	2 (3.9)	13 (25.5)	36 (70.6)	51 (100)	0 (0)	40 (78.4)	11 (21.6)	0 (0)	2 (3.9)	18 (35.3)	31 (60.8)	1 (2)
1998	203	1 (0.5)	81 (39.9)	121 (59.6)	203 (100)	0 (0)	122 (60.1)	81 (39.9)	0 (0)	16 (7.9)	64 (31.5)	123 (60.6)	7 (3.4)
1999	167	3 (1.6)	24 (14.4)	140 (83.8)	167 (100)	0 (0)	153 (91.6)	14 (8.4)	0 (0)	5 (3.0)	31 (18.6)	131 (78.4)	9 (5.4)
2000	61	0 (0)	17 (27.9)	44 (72.1)	61 (100)	0 (0)	50 (81.9)	11 (18.0)	0 (0)	0 (0)	15 (24.6)	46 (75.4)	10 (16.4)
2001	110	1 (0.9)	19 (17.3)	90 (81.8)	110 (100)	0 (0)	90 (81.8)	20 (18.2)	0 (0)	2 (1.8)	28 (25.5)	80 (72.7)	24 (21.8)

*192 strains were tested for ceftriaxone and 199 for ciprofloxacin.