

PostScript

MATTERS ARISING

If you have a burning desire to respond to a paper published in the *Annals of the Rheumatic Diseases*, why not make use of our "rapid response" option?

Log on to our website (www.annrheumdis.com), find the paper that interests you, and send your response via email by clicking on "eLetters" option in the box at the top right hand corner.

Providing it isn't libellous or obscene, it will be posted within seven days. You can retrieve it by clicking on "read eLetters" on our homepage.

The editor will decide as before whether also to publish it in a future paper issue.

Differences in the management of shoulder pain between primary and secondary care in Europe: time for a consensus

We read with great interest the articles of Van der Windt and Bouter¹ and Hay *et al.*² There is no doubt that the study of Hay *et al.* is well designed and has practical implications. They showed that physiotherapy or subacromial joint injection are equally effective for shoulder pain. This is new evidence as, so far, there has been little evidence to support the effectiveness of any common intervention for shoulder pain.³ However, the definition of "shoulder pain" illustrates the practical problem in diagnosis that general practitioners and hospital specialists face in routine clinical practice. We agree that the positive outcome for physiotherapy may reflect the increased contact time between physiotherapist and patient or the better understanding of the anatomical problem by the physiotherapist. The differences in management and in the effectiveness of physiotherapy by the British compared with the Dutch may also represent a cultural difference between the expectations and beliefs of patients in the two countries. It is likely that physiotherapy departments could be overloaded with referrals from primary care doctors if they are always the first next step in the pathway of managing shoulder problems. Hay *et al.* did not carry out a cost-benefit analysis of the different treatments for shoulder pain (that is, injection *v* physiotherapy). A course of physiotherapy would cost around £200–320 (€284–454), whereas an injection would cost around £60 (€85).

There is a lack of consensus in the UK about the exact role of the general practitioner in the treatment of shoulder disease.⁴ A survey among rheumatologists and physiotherapists practising in the Southeast Thames Region of London (47 rheumatologists and 9 physiotherapists) showed that the management of adhesive capsulitis in secondary care varied widely. Nearly all the rheumatologists

(98%) used intra-articular steroid injection, but the time, site, and frequency of injections were variable, with 72% believing that early injections are a priority. One of five rheumatologists (22%) believed that physiotherapy and mobilisation offered no benefit. Only a small number of rheumatologists (14%) believed physiotherapy to be the only means of treatment.⁵ Interestingly, 90% of physiotherapists working in secondary care wanted to see patients with a frozen shoulder as early as possible before or immediately after steroid injections. However their waiting time varied considerably (range of 3 days–3 months).

Similarly, across Europe treatment of shoulder pain varies considerably between primary and secondary care.^{6,7} Therefore we propose that European consensus guidelines on the management of the painful shoulder should be developed.^{8,9} This consensus may be weakened by the lack of an adequate evidence base. In addition, we would suggest a third and fourth arm to future studies—steroid injection with physiotherapy and a no intervention control group.

D G Kassimos

Rheumatology Department, Dudley Group of Hospitals, Dudley, West Midlands, UK

G Panayi

Rheumatology Department, Division of Medicine, GKT School of Medicine, Guy's Hospital, London UK

Correspondence to: Dr D G Kassimos; kassimos.d@dudleygoh-tr.wmids.nhs.uk

Dr D G Kassimos is on study leave from the Ministry of Defence of Greece.

References

- 1 Van der Windt DAWM, Bouter LM. Physiotherapy or corticosteroid injection for shoulder pain? *Ann Rheum Dis* 2003;62:385–7.
- 2 Hay EM, Thomas E, Paterson SM, Dzieczic K, Croft PR. A pragmatic randomized controlled trial of local corticosteroid injection and physiotherapy for the treatment of new episodes of unilateral shoulder pain in primary care. *Ann Rheum Dis* 2003;62:394–9.
- 3 Green S, Buchbinder R, Glazier R, Forbes A. Systematic review of randomized controlled trials of interventions for painful shoulder: selection criteria, outcome assessment, and efficacy. *BMJ* 1998;316:354–60.
- 4 Huston GJ. An offer of rheumatology training: failure to influence clinic referrals. *BMJ* 1988;296:1773–4.
- 5 Kassimos DG, Panayi G, Choy E, Yanni G. The current practice on management of frozen shoulder in secondary care. *BMJ*, (e-letter) May, 2002.
- 6 Van der Windt DAWN, Koes BW, Boeke AJP, Deville W, De Jong BA, Bouter LM. Corticosteroid injections versus physiotherapy for painful stiff shoulder in primary care: randomised trial. *BMJ* 1998;317:1292–6.
- 7 Winters JC, Sobel JS, Groenier KH, Arendzen HJ, Meyboom-de Jong B. Comparison of physiotherapy, manipulation, and corticosteroid injection for treating shoulder complaints in general practice: randomised, single blind study. *BMJ* 1997;314:1320–5.
- 8 Winters JC, De Jongh AC, Van der Windt DAWM, Jonquiere M, De Winter AF, Van der Heijden GJMG, *et al.* General practice guidelines on shoulder complaints [NHG-Standaard Schouderklachten]. *Huisarts&Wetenschap* 1999:222–31.

- 9 AAOS. Clinical guideline on shoulder pain. Rosemont, IL: American Academy of Orthopedic Surgeons, 2001, (www.aaos.org).

Author's reply

Kassimos and Panayi deal with several important issues about the management of shoulder pain in their comments on the article by Hay *et al.*¹ and our leader.² We agree that differences in the effect of treatment between the Netherlands and England may, at least partly, reflect differences in the organisation of care, as well as differences in expectations and beliefs between the two countries. We are also aware of the lack of consensus among general practitioners, physiotherapists, and rheumatologists about the management of shoulder pain. Between primary and secondary care, especially, the differences are large. This can partly be explained by the fact that the primary care doctor is confronted with an entirely different spectrum of disease than the specialist.³ Many patients in primary care present with signs and symptoms that are troublesome and cause worry, but are relatively benign and have a favourable prognosis. Patients referred to secondary care have been pre-selected by the nature and severity of symptoms, and have another prognosis, resulting in different treatment requirements.

The lack of consensus among health professionals, indeed, emphasises the need for multidisciplinary guidelines for the management of shoulder pain. Regardless of the quality of the evidence base, multidisciplinary guidelines will facilitate communication among health professionals and may optimise diagnosis and treatment of patients with shoulder pain. We suggest that the AGREE Instrument (Appraisal of Guidelines for Research and Evaluation)⁴ is used in the development of any guideline for shoulder pain. This instrument includes recommendations for the description of the scope and purpose of a guideline, stakeholder involvement, rigour of development, clarity and presentation, applicability, and editorial independence.

The development of a European guideline for shoulder pain will be quite an undertaking. The authors of the EULAR guideline for the management of knee osteoarthritis indicated that there was often discordance between research evidence and the opinion of experts.⁵ In this international guideline, variation across countries in healthcare delivery systems, access to health professionals, ways of funding, and attitudes towards the disease, all contributed to this discordance. The use of a Delphi system permitted consensus agreement on difficult issues, but still the applicability in individual countries may be limited. In the case of shoulder pain, it may be wise to start out with the development of national (multidisciplinary) guidelines. As yet, only a few European countries or professional organisations have developed such guidelines.

Finally, regarding the closing point by Kassimos and Panayi, we agree that there is a need for additional research comparing physiotherapy or corticosteroid injections with a no treatment control. It might be difficult or undesirable to carry out such a

trial in patients with severe pain and limitations in daily activities, but controlled trials will certainly help to establish the effectiveness and cost effectiveness of physiotherapy and injections in patients with mild to moderate shoulder pain. Future trials may also evaluate the effectiveness of combined treatment (injections plus physiotherapy).

D A W M van der Windt

Correspondence to: Dr D A W M van der Windt;
dawm.van_der_windt.emgo@med.vu.nl

References

- 1 **Van der Windt DAWM**, Bouter LM. Physiotherapy or corticosteroid injection for shoulder pain. *Ann Rheum Dis* 2003;**62**:385–7.
- 2 **Hay EM**, Thomas E, Paterson SM, Dzieczic K, Croft PR. A pragmatic randomized controlled trial of local corticosteroid injection and physiotherapy for the treatment of new episodes of unilateral shoulder pain in primary care. *Ann Rheum Dis* 2003;**62**:394–9.
- 3 **Knotterus JA**. Medical decision making by general practitioners and specialists. *Fam Pract* 1991;**8**:305–7.
- 4 **The AGREE Collaboration**. Appraisal of Guidelines for Research & Evaluation (AGREE) Instrument. <http://www.agreecollaboration.org>.
- 5 **Pendleton A**, Arden N, Dougados M, Doherty M, Bannwarth B, Bijlsma JWJ, et al. EULAR Recommendations for the management of knee osteoarthritis: report of a Task Force of the Standing Committee for International Clinical Studies Including Therapeutic Trials (ESCSIT). *Ann Rheum Dis* 2000;**59**:936–44.

Exercise in juvenile idiopathic arthritis: promise or passé

We were interested in the recently published article in the *Annals* by Takken *et al.*¹ Notwithstanding their substantial work, we have a few comments pertaining to the exercise regimens in children with juvenile idiopathic arthritis (JIA).

Firstly, we did not see any information about whether the patients had ever been following an exercise protocol before they were included in the study and also whether they were prescribed a protocol afterwards. Information about these two points is important for an interpretation of the patients' results and for providing evidence about the practical implications of the study.

Secondly, when mentioning the diminished loadbearing capacity of these subjects owing to their inflammatory disease and the immune suppressive drugs, they drew attention to a study in which weightbearing exercises were shown to improve the aerobic endurance of such patients.² At this point, it is noteworthy to add that the myopathic effects of corticosteroids should also be remembered when exercise is prescribed. It is known that eccentric muscle contractions in normal subjects are responsible for a much greater efflux of muscle enzymes into the circulation than is caused by concentric contractions, and are associated with ultrastructural indications of damage to the muscle.^{3,4} Thus in patients with JIA—where steroid use is prevalent—concentric types of exercise should preferably be prescribed. These may include simply walking, cycling, or running. However, the list of sports which can be played is endless and there is an excess of activities these—otherwise sedentary—children can be encouraged to take part in to obtain exercise.⁵ In this way not only will there be an increase in their aerobic

capacities but also they will encounter fewer disabilities related to muscle anaerobiosis—much more common in children who use much more energy than adults during daily activities.

L Özçakar

Hacettepe University Medical School, Department of Physical Medicine and Rehabilitation, Ankara, Turkey

Z B Özçakar

Ankara University Medical School, Department of Pediatric Nephrology, Ankara, Turkey

Correspondence to: Dr L Özçakar, Yeni Ankara sokak 27/1, Cebeci, Ankara, Turkey; lozcar@yahoo.com

References

- 1 **Takken T**, van der Net J, Kuis W, Helders PJM. Physical activity and health related physical fitness in children with juvenile idiopathic arthritis. *Ann Rheum Dis* 2003;**62**:885–9.
- 2 **Klepper SE**. Effects of an eight-week physical conditioning program on disease signs and symptoms in children with chronic arthritis. *Arthritis Care Res* 1999;**12**:52–60.
- 3 **Newham DJ**, Jones DA, Edwards RH. Plasma creatine kinase changes after eccentric and concentric contractions. *Muscle Nerve* 1986;**9**:59–63.
- 4 **Friden J**, Sjöström M, Ekblom B. A morphological study of delayed muscle soreness. *Experientia* 1981;**37**:506–7.
- 5 **Ozçakar L**, Topaloglu R. An amateur badminton player with juvenile dermatomyositis: courage and questions. *Br J Sports Med* (in press).

Authors' reply

We would sincerely like to thank Özçakar and Özçakar for their response.

Firstly, the patients studied did not actively participate in endurance sports activities at the time of measurement. However, some of the patients had taken part in some sports activities in the period before the disease onset, but not in the six months before our study was performed. It is known from the literature that there is a rapid diminution in fitness once training stops.¹

We did not prescribe exercises based on the current findings. The Caltrac is a portable electronic activity monitor that measures movements in the vertical plane. It sums and integrates the absolute value of the acceleration versus time curve and derives a numerical count that is displayed on the monitor. There are no normal values for this instrument. The described data were baseline data from a randomised controlled trial for the effectiveness of aquatic exercise therapy.² Secondly, we did not discuss the effects of corticosteroid treatment on aerobic fitness, because only a small minority of our patients (four) had systemic juvenile idiopathic arthritis (JIA), in which steroids are the preferred treatment. In other JIA subgroups, non-steroidal anti-inflammatory drugs and methotrexate are the common treatment in our country nowadays. A discussion on the effects of drugs and inflammation on exercise capacity can be found elsewhere.^{3,4}

We could not comment on the paper cited by the authors because it had not yet been published when we wrote this letter. Furthermore, we would like to add that JIA and juvenile dermatomyositis (JDM) are distinct diseases and that the exercise capacities of these patients do differ significantly, with patients with JDM being more affected than patients with JIA.⁵ Therefore, the exercise prescription for patients with JIA and JDM

should be different, and adapted to the individual patients needs and capacity.

Moreover, we are not aware of studies showing an anaerobiosis in muscles of patients with JIA during activities of daily living.

T Takken, J van der Net, W Kuis, P J M Helders

Department of Paediatric Physical Therapy, Wilhelmina Children's Hospital, University Medical Centre Utrecht, Room KB2.056.0, PO Box 85090, NL 3508 AB Utrecht, The Netherlands

Correspondence to: Dr T Takken, t.takken@wku.az.u.nl

References

- 1 **Rowland TW**. Effect of prolonged inactivity on aerobic fitness of children. *J Sports Med Phys Fitness* 1994;**34**:147–55.
- 2 **Takken T**, Van der Net J, Kuis W, Helders PJ. Aquatic fitness training for children with juvenile idiopathic arthritis. *Rheumatology (Oxford)* 2003;**42**:1408–14.
- 3 **Walsmith J**, Roubenoff R. Cachexia in rheumatoid arthritis. *Int J Cardiol* 2002;**85**:89–99.
- 4 **Takken T**. Studies on physical performance and functional ability in juvenile idiopathic arthritis. Utrecht, the Netherlands: Utrecht University, 2003;(PhD thesis).
- 5 **Takken T**, Spermon N, Helders PJ, Prakken AB, Van Der Net J. Aerobic exercise capacity in patients with juvenile dermatomyositis. *J Rheumatol* 2003;**30**:1075–80.

Progressive multifocal leucoencephalopathy and immunosuppression

We report an immunocompromised patient with progressive multifocal leucoencephalopathy (PML), who demonstrates the usefulness and limitation of the algorithm of Warnatz *et al.*¹ for investigation of patients with pre-existing autoimmune diseases and new onset neuropsychiatric abnormalities. A prerequisite for the use of this algorithm requires a high degree of awareness for infection to prevent misclassification of the underlying problem.

This 61 year old white woman had had dermatomyositis since 1996 as manifest by Gottron's papules, heliotrope rash, proximal muscle weakness, and antinuclear antibody (ANA) titre 1/1280 speckled pattern. Previous management included azathioprine, methotrexate, hydroxychloroquine, and intravenous immunoglobulin; the disease was controlled for the previous 20 months while receiving cyclophosphamide 100 mg and prednisone 5 mg daily.

One week before admission the patient developed dizziness, weakness, and left sided hearing loss. Meclizine was prescribed for possible Ménière's disease. Facial weakness and dysarthria developed. A physical examination showed left sided hearing loss, left facial droop, left hemiparesis with concomitant graphaesthesia, and impaired stereognosis; left patella hyperreflexia was also present. Magnetic resonance imaging (MRI) of the brain was performed at an outlying facility and was felt to demonstrate a subacute infarct. There was increased signal intensity in the right posterior temporal lobe measuring 4 cm in diameter without mass effect or haemorrhage, and an additional temporo-parietal lesion. Punctate areas of increased signal were seen in the mid-portion of the pons (fig 1A). She was admitted for further evaluation of stroke. Laboratory data included normal complete blood counts,