

EXTENDED REPORT

Effect of treatment of rheumatoid arthritis with infliximab on IFN γ , IL4, T-bet, and GATA-3 expression: link with improvement of systemic inflammation and disease activity

M Kawashima, P Miossec



Ann Rheum Dis 2005;64:415–418. doi: 10.1136/ard.2004.022731

See end of article for authors' affiliations

Correspondence to: Professor P Miossec, Clinical Immunology Unit, Department of Immunology and Rheumatology, Hospital Edouard Herriot, 69437 Lyon Cedex 03, France; miossec@univ-lyon1.fr

Accepted 9 July 2004
Published Online First
29 July 2004

Objective: To study interferon γ (IFN γ) production and the expression of T-bet and GATA-3, the transcription factors associated with Th1 and Th2, in peripheral blood mononuclear cells (PBMC) from patients with rheumatoid arthritis before and during infliximab treatment, so as to distinguish between a disease specific and a disease activity dependent defect.

Methods: Rheumatoid PBMC were obtained at weeks 0 and 6 of infliximab treatment and cultured for seven days with or without interleukin (IL)12 or the combination of IL12 and IL18. IFN γ concentrations in supernatants were determined by ELISA. mRNA expression of IFN γ , IL4, T-bet, and GATA-3 was determined by real time RT-PCR in whole blood at weeks 0 and 22.

Results: A reduction in spontaneous IFN γ production and in the response to Th1 inducing cytokines occurred in rheumatoid PBMC. Reduction of systemic inflammation with infliximab treatment increased IFN γ production in response to IL12 or IL12+IL18. The IFN γ /IL4 expression ratio of rheumatoid blood before treatment was lower than in healthy controls but was increased by infliximab treatment. T-bet expression or T-bet/GATA-3 ratio of rheumatoid blood was less than in controls. The T-bet/GATA-3 ratio was not influenced by infliximab treatment.

Conclusions: Regulation of T-bet and GATA-3 or IFN γ and IL4 expression appeared different. The IFN γ /IL4 ratio might express the blood Th1/Th2 balance better than the T-bet/GATA-3 ratio. Reduced IFN γ production by rheumatoid PBMC and levels of IFN γ and IL4 mRNA expression in blood were linked to disease improvement, indicating an association between this systemic Th1 feature and disease activity.

Anti-tumour necrosis α (TNF α) treatment has been shown to decrease disease activity in patients with rheumatoid arthritis, but cases of severe tuberculosis have been reported during the early stage of treatment.¹ However, an increased incidence of tuberculosis appears to be associated with rheumatoid arthritis even in the absence of anti-TNF treatment.² Although the role of TNF α in the human immune response to mycobacteria is incompletely understood, TNF α has a pivotal role in granuloma formation and in the control of the disease in animals.³

Considering the critical role of interferon γ (IFN γ)—the prototypic Th1 cytokine—in tubercular infection, an underlying defect of cell mediated immunity in rheumatoid blood may be related to the development of tuberculosis. Indeed, a disturbed balance between T helper (Th)1 and Th2 subsets has been reported in the pathogenesis of rheumatoid arthritis.⁴ With others, we have shown the selective accumulation of Th1 and Th0 cells in the synovium by intracellular cytokine staining of peripheral blood and synovial tissue T cells from rheumatoid patients.⁵ On the other hand, the Th1/Th2 balance in the peripheral circulation remains controversial in rheumatoid arthritis. At the protein level, rheumatoid blood cells produced less IFN γ and more interleukin (IL)4 in response to non-specific stimulation than controls.⁶ We recently found that peripheral blood mononuclear cells (PBMC) from rheumatoid patients were less sensitive to the Th1 inducing cytokines IL12 and IL18 than those from controls.⁷ Furthermore, mRNA expression of IFN γ and IL4, as well as T-bet and GATA-3—the transcription factors involved in Th1 and Th2 development, respectively—showed a

decreased Th1 pattern in rheumatoid blood compared with control.⁷

As anti-TNF treatment for rheumatoid arthritis can often improve disease activity rather rapidly, we studied whether such treatment might affect the underlying systemic Th1 pattern. We looked at mRNA expression of IFN γ , IL4, T-bet, and GATA-3 in the circulation of rheumatoid patients at week 0 and week 22 of infliximab treatment. In addition, we studied IFN γ production by PBMC from rheumatoid patients in response to IL12 and IL18 at week 0 and 6. Responders, but not non-responders, augmented the IFN γ /IL4 ratio in whole blood. Infliximab treatment augmented the levels of IFN γ production by rheumatoid PBMC cultures with or without IL12 or IL12+IL18. The systemic Th1 immune pattern of rheumatoid arthritis indicated by the impaired IFN γ production by rheumatoid PBMC appears to be related to disease activity.

METHODS

Whole blood sample collection and storage

Blood samples were obtained from 12 healthy controls (three men, nine women) and from 15 patients with rheumatoid arthritis (four men, 11 women), who fulfilled the 1987 revised criteria of the American College of Rheumatology (ACR),⁸ before the first (week 0) and the fifth (week 22) infusion of infliximab (given in a dose of 3 mg/kg according

Abbreviations: ACR, American College of Rheumatology; IFN γ , interferon γ ; IL, interleukin; PBMC, peripheral blood mononuclear cells; Th, T helper cell; TNF α , tumour necrosis factor α

to the ATTRACT protocol). The mean age of the healthy controls was 45.4 years (range 31 to 64), and of the rheumatoid patients, 48.2 years (range 28 to 68). Mean disease duration was 8.2 years (range 3 to 18). The majority of patients were being treated with non-steroidal anti-inflammatory drugs and methotrexate, alone or combined with prednisone. Patient response was quantified using the ACR response criteria.⁹ Patients with an ACR score of 20 or more were defined as responders and those with a score of less than 20 as non-responders. Peripheral blood samples were collected in PAXgene tubes containing a cationic detergent and additive salts (PreAnalytiX/QIAGEN GmbH, Hilden, Germany), and then stored at -20°C .

RNA isolation from whole blood sample and RT-PCR

PAXgene blood RNA kit (PreAnalytiX) was used for purification of intracellular RNA from whole blood samples. The ThermoScriptTM reverse transcriptase polymerase chain reaction (RT-PCR) system (Invitrogen SARL, Cergy Pontoise, France) was used to synthesise cDNA.

Parameter specific primer sets for IFN γ , IL4, T-bet, GATA-3, and β actin optimised for the LightCycler instrument (Roche Applied Science, Roche Diagnostics Corporation, Indianapolis, Indiana, USA) were developed and purchased from Search-LC (Heidelberg, Germany). PCR was carried out as recently described,¹⁰ using the LightCycler FastStart DNA Sybr Green I kit (Roche Applied Science) according to the protocol provided with the parameter specific kits (45 amplification cycles, denaturation at 96°C , primer annealing at 68°C with touchdown to 58°C , amplicon extension at 72°C). The copy number of IFN γ , IL4, T-bet, and GATA-3 was normalised by the housekeeping gene β actin and is presented as the number of transcripts per copy of β actin.




Figure 1 Increased type 1 response in whole blood from patients with rheumatoid arthritis following anti-TNF α treatment. Whole blood samples were obtained from 12 healthy controls (HC) and from 15 patients with rheumatoid arthritis (RA) at week 0 (w0) and 22 (w22) of infliximab treatment. IFN γ , IL4, T-bet, GATA-3, and β actin mRNA expression levels were determined by quantitative reverse transcriptase polymerase chain reaction. * $p < 0.05$, ** $p < 0.01$

Cytokines and reagents

Recombinant human IL12 was purchased from R&D systems, Abingdon, Berkshire, United Kingdom. Recombinant human IL18 was from MBL, Nagoya, Japan. RPMI 1640 culture media was purchased from Invitrogen SARL, Cergy Pontoise, France and supplemented with 100 units/ml penicillin, 100 $\mu\text{g}/\text{ml}$ streptomycin, and 10% fetal calf serum (Invitrogen).

Preparation of PBMC culture

Peripheral blood samples were obtained from six controls (two men, four women), and from six rheumatoid patients (two men, four women) before the first (week 0) and the third (week 6) infusion of infliximab. The mean age of the controls was 48.3 years (range 44 to 61) and of the rheumatoid patients, 51.7 years (range 44 to 65), and mean disease duration was 9.6 years (range 4 to 38). PBMC were isolated from heparinised blood by Ficoll density gradient centrifugation, washed twice with phosphate buffered saline, and resuspended in RPMI1640 medium. PBMC were cultured in 96-well plates (Nunc, Roskilde, Denmark) at a density of 1×10^6 cells/ml in 200 μl of RPMI medium.

Determination of levels of IFN γ and serum C reactive protein

IFN γ levels were measured by quantitative sandwich enzyme linked immunosorbent assay (ELISA), using a commercially available ELISA kit (DuoSet ELISA Development System human IFN γ , R&D Systems). The detection limit of the assay was 20 pg/ml. C reactive protein concentrations were measured by the hospital laboratory, and the limit of assay was 1.5 mg/l. Undetectable levels were regarded as 0.

Statistical analysis

Results were expressed as mean (SD) of the indicated number of experiments. The statistical significance of differences between two groups was determined by the Mann-Whitney U test. Wilcoxon's signed rank test was used to analyse matched pairs.

RESULTS

Improvement of Th1 defect in rheumatoid blood by infliximab treatment

Paired peripheral blood samples from 15 rheumatoid patients at week 0 and 22 of infliximab treatment and from the healthy controls were collected directly in PAXgene tubes. IFN γ and IL4 mRNA expression levels were then determined




Figure 2 Improvement in impaired interferon γ (IFN γ) production of peripheral blood mononuclear cells (PBMC) from patients with rheumatoid arthritis resulting from anti-tumour necrosis α (TNF α) treatment. PBMC from healthy controls (HC) ($n = 6$) and rheumatoid patients (RA) ($n = 6$) obtained at weeks 0 and 6 of infliximab treatment were cultured with or without interleukin (IL)12 (1 ng/ml) or IL12+IL18 (5 ng/ml) for seven days. IFN γ levels in the supernatants were determined by a specific sandwich enzyme linked immunosorbent assay. * $p < 0.05$

by quantitative RT-PCR. As shown in fig 1A, IFN γ expression level in rheumatoid blood at week 0 was almost identical to that of the controls, but the IFN γ /IL4 expression ratio was significantly lower in the rheumatoid samples (fig 1B). However, infliximab augmented IFN γ expression and the IFN γ /IL4 ratio in rheumatoid blood following 22 weeks of treatment. Eleven responders of 15 patients who had an ACR response score of ≥ 20 at week 22 had an increased IFN γ /IL4 ratio in their blood (from 1.26 (0.76) to 3.78 (2.59), $p = 0.003$). However, four non-responders who had an ACR response score of less than 20 did not show an increased IFN γ /IL4 ratio (from 1.30 (0.83) to 0.82 (0.22), $p = 0.61$). Thus rheumatoid blood expressed markers of a decreased Th1/Th2 balance. This defect was modified by infliximab treatment only in the responders.

Impaired T-bet/GATA-3 ratio in rheumatoid blood

We next extended the results to the transcription factors T-bet and GATA-3, which have been described as Th1 and Th2 markers, respectively. Rheumatoid blood showed a decreased T-bet expression or T-bet/GATA-3 ratio compared with the controls at both week 0 and week 22 of infliximab treatment (fig 1C). Infliximab upregulated not only T-bet but also GATA-3 expression; therefore the T-bet/GATA-3 ratio was not influenced by infliximab treatment (fig 1D). Accordingly, the expression of IFN γ and T-bet mRNA in rheumatoid blood appeared to be regulated differently. The IFN γ /IL4 ratio could be more sensitive than the T-bet/GATA-3 ratio in evaluating the Th1/Th2 balance in blood samples.

Improvement in impaired IFN γ production by rheumatoid PBMC following infliximab

PBMC samples from six rheumatoid patients at week 0 and week 6 of infliximab treatment and from six healthy controls were cultured for seven days with or without IL12 (1 ng/ml) or the combination of IL12 and IL18 (5 ng/ml). IFN γ levels in the supernatants were determined by ELISA. All patients had a decrease in C reactive protein concentrations (from 72.2 (47.9) to 19.4 (20.5) mg/l, $p = 0.02$) following six weeks of infliximab treatment. Rheumatoid PBMC at week 0 produced significantly lower levels of IFN γ than control. However, rheumatoid PBMC at week 6 produced higher levels of IFN γ than the week 0 samples, either in response to IL12 alone ($p = 0.04$) or to IL12+IL18 ($p = 0.04$) (fig 2), indicating improvement in the impaired IFN γ production by rheumatoid PBMC following to anti-TNF treatment in parallel with amelioration of the disease.

DISCUSSION

Several lines of evidence have shown the predominance of IFN γ producing Th1 cells at the site of chronic inflammation in rheumatoid arthritis. An increased IFN γ /IL4 ratio in synovial fluid compared with peripheral blood has also been reported. However, the Th1/Th2 balance in the peripheral circulation in rheumatoid arthritis remains to be clarified. We and others have shown that rheumatoid blood cells produce less IFN γ and more IL4 than cells from healthy controls,⁶ indicating a decreased Th1 response in the peripheral circulation in rheumatoid arthritis. Recently we found that PBMC from rheumatoid patients produced lower levels of IFN γ in response to IL12 and IL18 than controls,¹¹ suggesting that rheumatoid patients have a systemic Th1 defect.

In the present study, we looked at the systemic Th1/Th2 balance in patients with rheumatoid arthritis before and after anti-TNF α treatment. Responders to anti-TNF α clearly upregulated their blood expression of IFN γ , but non-responders did not. This suggests that the decreased systemic Th1 pattern seen in rheumatoid arthritis may be related to disease activity rather than being disease specific. After

anti-TNF α treatment PBMC from rheumatoid patients also produced higher levels of IFN γ than before treatment in response to IL12 and IL18. Our results showed that patients with active rheumatoid arthritis may have decreased systemic Th1, and improvement of this defect appears to be associated with amelioration of rheumatoid activity. The T-bet/GATA-3 ratio was also lower in rheumatoid blood than in controls, and this remained unchanged by infliximab treatment. T-bet is expressed by various blood cells such as T cells, B cells, and monocytes. Thus the T-bet/GATA-3 ratio may not be a sensitive Th1/Th2 marker in blood samples.

Accumulation of Th2 cells rather than Th1 cells in the blood may result from changes in T cell migration. Indeed, the numbers of IFN γ producing T cells were significantly increased in the peripheral blood of rheumatoid patients shortly after anti-TNF α treatment, resulting in a shift of the Th1:Th2 ratio towards Th1 in the blood.¹² Adhesion molecules such as P- and E-selectin or VCAM-1 are considered to be important for the selective homing of Th1 cells but not of Th2 cells.¹³ Expression of E-selectin and VCAM-1 was significantly reduced by anti-TNF treatment.¹⁴ Thus anti-TNF treatment may suppress the selective migration of Th1 cells into rheumatoid synovium through the rapid downregulation of adhesion molecules. It remains to be determined whether this is associated with an increased migration of Th2 cells with anti-inflammatory properties.

The contrast between the improvement described above and the reactivation of tuberculosis at an early stage of infliximab treatment needs to be explored. Our understanding of the increased risk based on these results is that at an early stage of treatment rheumatoid patients express a systemic immune defect related to disease activity. This defect is increased when anti-TNF is used because it affects cell interactions and granuloma formation. This has a beneficial effect on joint inflammation, leading to reduced cellularity of the synovium. However, the effect could also be deleterious, as tubercular granulomas may dissociate leading to massive spread of the disease. Later the risk is reduced following the improvement in disease activity and the associated immune defect.

In conclusion, our data show that successful anti-TNF α treatment not only improves systemic inflammation and rheumatoid disease activity but also improves the impairment of systemic Th1 immune function.

Authors' affiliations

M Kawashima, P Miossec, Department of Immunology and Rheumatology and INSERM U-403, Hospital Edouard Herriot, Lyon, France

REFERENCES

- Keane J, Gershon S, Wise RP, Mirabile-Levens E, Kasznica J, Schwietzman WD, et al. Tuberculosis associated with infliximab, a tumor necrosis factor alpha-neutralizing agent. *N Engl J Med* 2001;**345**:1098-104.
- Carmona L, Hernandez-Garcia C, Vadillo C, Pato E, Balsa A, Gonzalez-Alvaro I, et al. Increased risk of tuberculosis in patients with rheumatoid arthritis. *J Rheumatol* 2003;**30**:1436-9.
- Mohan VP, Scanga CA, Yu K, Scott HM, Tanaka KE, Tsang E, et al. Effects of tumor necrosis factor alpha on host immune response in chronic persistent tuberculosis: possible role for limiting pathology. *Infect Immun* 2001;**69**:1847-55.
- Miossec P, van den Berg W. Th1/Th2 cytokine balance in arthritis. *Arthritis Rheum* 1997;**40**:2105-15.
- Morita Y, Yamamura M, Kawashima M, Harada S, Tsuji K, Shibuya K, et al. Flow cytometric single-cell analysis of cytokine production by CD4+ T cells in synovial tissue and peripheral blood from patients with rheumatoid arthritis. *Arthritis Rheum* 1998;**41**:1669-76.
- Haddad A, Bienvenu J, Miossec P. Increased production of a Th2 cytokine profile by activated whole blood cells from rheumatoid arthritis patients. *J Clin Immunol* 1998;**18**:399-403.
- Kawashima M, Miossec P. Decreased response to IL-12 and IL-18 of peripheral blood cells in rheumatoid arthritis. *Arthritis Res Ther* 2004;**6**:R39-45.

- 8 **Arnett FC**, Edworthy SM, Bloch DA, McShane DJ, Fries JF, Cooper NS, *et al.* The American Rheumatism Association 1987 revised criteria for the classification of rheumatoid arthritis. *Arthritis Rheum* 1988;**31**:315–24.
- 9 **Felson D**, Anderson J, Boers M. American college of rheumatology preliminary definition of improvement in rheumatoid arthritis. *Arthritis Rheum* 1995;**38**:727–35.
- 10 **Krug A**, Towarowski A, Britsch S, Rothenfusser S, Hornung V, Bals R, *et al.* Toll-like receptor expression reveals CpG DNA as a unique microbial stimulus for plasmacytoid dendritic cells which synergizes with CD40 ligand to induce high amounts of IL-12. *Eur J Immunol* 2001;**31**:3026–37.
- 11 **Kawashima M**, Miossec P. Decreased response to IL-12 and IL-18 of blood versus synovial RA cells: linkage with T-bet and modulation with IL-18BP [abstract]. *Arthritis Rheum* 2002;**46**(suppl 9):S252–3.
- 12 **Maurice MM**, van der Graaff WL, Leow A, Breedveld FC, van Lier RA, Verweij CL. Treatment with monoclonal anti-tumor necrosis factor alpha antibody results in an accumulation of Th1 CD4+ T cells in the peripheral blood of patients with rheumatoid arthritis. *Arthritis Rheum* 1999;**42**:2166–73.
- 13 **Austrup F**, Vestweber D, Borges E, Lohning M, Brauer R, Herz U, *et al.* P- and E-selectin mediate recruitment of T-helper-1 but not T-helper-2 cells into inflamed tissues. *Nature* 1997;**385**:81–3.
- 14 **Tak PP**, Taylor PC, Breedveld FC, Smeets TJ, Daha MR, Kluin PM, *et al.* Decrease in cellularity and expression of adhesion molecules by anti-tumor necrosis factor alpha monoclonal antibody treatment in patients with rheumatoid arthritis. *Arthritis Rheum* 1996;**39**:1077–81.

Get published within days of acceptance with ARD

We are delighted to announce that the *Annals of the Rheumatic Diseases* launched a “publish ahead of print” programme in February 2004. Selected papers are fast tracked and published online months before they appear in the print journal.

Papers of major significance to the international rheumatology community are **published within days of acceptance**. The first published article is the raw accepted manuscript; edited and typeset versions are also published as soon as they are available.

In addition to being available on *ARD Online*, the publish ahead of print articles are searchable through PubMed/Medline—establishing primacy for your work. They are linked from the *ARD Online* home page.

ARD's publish ahead of print programme is unique among the major rheumatology journals—to take advantage of this service submit your papers to *Annals of the Rheumatic Diseases* using our online submission and review system Bench>Press (<http://submit-ard.bmjournals.com>). For further information contact ARD@bmjgroup.com.