

Assessment of gonadal maturation by evaluation of spermaturia

Franz Schaefer, Joachim Marr, Christoffer Seidel, Wolfgang Tilgen, Karl Schärer

Abstract

The urinary excretion of spermatozoa (spermaturia) reflects the achievement of exocrine testis function during male puberty. In order to test the sensitivity and practicability of repetitive urine sampling, we analysed the sediments of 1160 first morning urine specimens obtained on successive days from 129 healthy schoolboys aged 10.1 to 17.8 years for the presence of spermatozoa. The proportion of subjects with sperm positive urines increased from pubic hair stage (PH) 1 (6%) to PH 5 (92%) with a steep rise between PH 2 and 3. Estimated median age of first positive spermaturia ('spermarche') was 14.1 years. While at PH 1 to 4 all positive samples were found within the first five days of collection, at PH 5 cumulative frequency of spermaturia increased up to day 8. We conclude that repetitive morning urine sampling is a useful tool in assessing spermaturia and may be helpful to screen for testicular damage in epidemiological surveys.

The detection of spermatozoa in the urine has been proposed as a quick, non-invasive method to assess the functional state of the maturing gonad and may be useful in screening for spermatogenic damage in male adolescents.¹⁻⁶ The evidence derived from longitudinal^{2, 4, 5} and cross sectional³ studies that spermaturia is a discontinuous phenomenon seems, however, to limit its value as a routine screening procedure. In those earlier reports either a single 24 hour urine^{2, 5} or variable numbers of morning urine specimens^{4, 6} were analysed. Considering the intermittent appearance of spermaturia, which probably occurs as a urethral wash out phenomenon after involuntary or masturbatory sperm emission,⁷ the probability of urinary sperm excretion should rise according to the number of specimens collected. No systematic investigations have been performed, however, in study populations of adequate size over extended

periods of time so far. Therefore, we attempted to determine the sensitivity and practicability of repetitive morning urine collections in a group of 129 healthy peripubertal boys, and to relate spermaturia to their state of sexual maturation.

Subjects and methods

One hundred and twenty nine healthy boys from five secondary schools in the city of Heidelberg agreed to participate in the study. The distribution of age and stages of puberty are given in tables 1 and 2. The children and their parents were informed that the purpose of the study was to evaluate normal sexual maturation by clinical assessment and urinary hormone excretion; however, they were not aware that spermaturia was investigated.

The boys were requested to collect their first morning void for two consecutive weeks from Monday to Friday respectively. In case of night time micturition they were advised to collect the first nocturnal urine. The specimens were brought to school in 100 ml plastic vessels and collected daily by one of the investigators. The bottles contained 3 ml of a 0.7% saponin solution in 0.4% aqueous formalin. The urine was stored at 4°C and analysed within 10 days after collection.

Urine analysis was performed according to the method described by Hirsch *et al.*³ In brief, a 10 ml aliquot of the well shaken urine was centrifuged at 1800 g for 15 minutes. The pellet was resuspended in 0.5 ml of the supernatant after decantation. The resuspended sediment was examined microscopically. Forty visual fields were checked at a magnification of $\times 400$ for the presence of spermatozoa. The same observer (JM) investigated all specimens. A specimen was considered sperm positive if one or more spermatozoa were detected.

At the end of the first week of urine collection, a clinical examination of the pubertal status was performed by a single observer (JM). Genital and pubic hair stage was recorded according to Tanner.⁸ The volume of each testicle was estimated by use of a Prader orchidometer, and the mean volume of the two testes was calculated.

ANALYSIS OF DATA

As not all boys delivered complete series of 10

Table 1 Age distribution of the 129 boys under study. Age ranged from 10.1 to 17.8 (median 12.8) years

	Age (years)						
	10	11	12	13	14	15	16-17
No of boys	14	26	31	14	20	18	6

Table 2 Distribution of Tanner stages of puberty (n=129). PH=pubic hair stage, G=genital stage

	PH1	G1	PH2	G2	PH3	G3	PH4	G4	PH5	G5
No of boys	34	41	45	27	16	14	17	14	17	33

Division of
Paediatric Nephrology,
University Children's
Hospital,
Heidelberg
Franz Schaefer
Joachim Marr
Christoffer Seidel
Karl Schärer

Department of
Dermatology,
University Hospital,
Heidelberg
Wolfgang Tilgen

Correspondence and
reprint requests to:
Dr Franz Schaefer,
Division of
Paediatric Nephrology,
University Children's
Hospital,
Im Neuenheimer Feld 150,
D-6900 Heidelberg,
Federal Republic of Germany.
Accepted 11 June 1990

Table 3 Distribution of number of collected urines

No of urine specimens collected	No (%) of subjects
4	2 (1.6)
5	6 (4.7)
6	2 (1.6)
8	5 (3.8)
9	10 (7.7)
10	104 (80.6)
	129 (100.0)

morning urine specimens (table 3), the cumulative probability to remain sperm negative was calculated by use of the Kaplan-Meier life table analysis.⁹ The subjects who dropped out of the study before spermaturia was detected were classified as lost to follow up at the last day of collection and treated as censored data. Spermaturia rates were estimated stratified by age, pubertal stage, and testicular volume.

The median age at spermarche, defined as the age at which spermaturia can be demonstrated in 50% of a given population, was estimated by probit analysis.¹⁰ In this procedure, the percentage of subjects exhibiting spermaturia are recorded at the successive age stages. To this percentage distribution a probit transformation is applied, and a regression line fitted to the probit values. Thus it is possible to calculate the median and variance of age at spermarche, assuming a Gaussian distribution of the timing of the event in the population.




Figure 1 Kaplan-Meier estimation of probability to remain sperm negative in up to 10 consecutive morning urine samples, stratified by Tanner's genital (G) stage.




Figure 2 Overall cumulative probability of spermaturia according to age, Tanner stages for pubic hair (1-5) and genitals (1-5), and testicular volume (ml) (n=129).

All calculations were performed by use of the respective procedures supplied in the SAS software package.¹¹

Results

As shown in table 3, the overall participation rate in the study was very high. Almost 90% of the boys collected nine or 10 morning urines. A total of 1120 specimens was analysed.

The Kaplan-Meier analysis of all patients' probability to remain sperm free with increasing numbers of examinations is shown in fig 1, stratified by Tanner's genital (G) stage. The proportion of sperm positive subjects increased with the state of genital maturity. After 10 urine collections, less than 20% of the boys in G5 had not exhibited spermaturia in any specimen, whereas in G1 spermatozoa were only found in one out of 40 boys. In presence of stages G1 to G4, no new sperm positive individuals were detected beyond day 5, whereas in G5 the cumulative frequency of spermaturia gradually increased up to day 8.

Similar analyses were performed according to the other maturational parameters under study. A survey of the probability of spermaturia as calculated by Kaplan-Meier estimation is given for age, pubertal stages, and testicular volume in fig 2. Age and testicular volume were subdivided into five classes. No subject was found sperm positive below 11 years of age; in the upper age groups, only about two thirds of the boys exhibited spermaturia. When related to pubic hair (PH) stage, a steep rise of the sperm positive fraction was seen between PH2 and PH3, with more than 90% being positive at PH5. The most discriminative parameter was testicular volume; when the mean testicular size exceeded 20 ml, 93.5% of the subjects excreted urinary sperm.

The probit analysis for the estimation of the age at spermarche is depicted in fig 3. The median age at spermarche was calculated as 14.1 years, with a 95% confidence interval of 13.5 to 14.9 years.

We also performed a quantitative analysis of spermaturia. The number of positive morning urine specimens per subject ranged from one to nine. A wide variation of sperm concentration (one to 458 spermatozoa/40 visual fields) was noted. In the sperm positive individuals no




Figure 3 Probit analysis of median age at spermarche. Age was grouped in six month intervals. Triangles mark unsmoothed percentages of sperm positive individuals per age group. Full line indicates smoothed sigmoid probability curve, dotted lines 95% confidence intervals.

significant correlation was found between the number of spermatozoa per 40 visual fields and age ($r=0.09$, $p=0.52$) or testicular volume ($r=0.23$, $p=0.12$). The correlations were not improved when the sperm count was related to urinary creatinine concentration to account for the varying dilution of the morning urines.

Discussion

The primary target of our study was to evaluate how regularly spermaturia can be picked up in healthy boys by the consecutive collection of 10 morning urines. In contrast to earlier studies^{3,4} the good compliance of the study participants allowed us to calculate overall probability rates from appropriate numbers of boys at all stages of puberty.

Our data demonstrate that spermaturia is better predicted by the clinical stage of puberty than by chronological age. The most powerful criterion to be sperm positive was a mean testicular volume of greater than 20 ml, but also in Tanner PH stage 5 more than 90% of the boys exhibited spermaturia (fig 2). Thus it is evident that spermaturia is detectable in the large majority of sexually mature subjects within a 10 day period of observation. Detailed analysis of cumulative spermaturia rates according to the number of investigated urines indicates that, however, once spermarche has occurred, spermaturia is a more regular event during early and mid puberty than in grown up individuals, where the first spermatozoa sometimes were not detected before the eighth day of collection (fig 1). This finding confirms an earlier suggestion by Hirsch *et al* who, analysing a maximum of five consecutive morning urine specimens, found a more slowly rising cumulative probability of spermaturia in 16 year olds compared with younger boys.³ We assume that this difference may reflect changes from predominantly spontaneous nocturnal sperm emission towards rather intermittent masturbatory ejaculations in late puberty.

The high sperm detection rate in mature boys with the sampling protocol used in this study also indicates that a 10 day repetitive morning urine collection is more suitable to identify reliably postspermarcheal subjects than a 24 hour or a short term morning urine sampling regime as proposed in earlier reports.^{2,3,5}

Despite the high sensitivity of our sampling technique it probably does not allow quantitative estimations of sperm production during puberty. In early studies of biopsy specimens a gradual increase of the density of mature spermatozoa in the tubular lumen during adolescence has been demonstrated,¹² and immediate postejaculatory urinary sperm concentrations correlated with ejaculate sperm counts.⁷ Thus one might expect that the sperm number

observed in the urine sediment should increase with advancing puberty. Yet no such correlation was found, possibly due to previous voiding between ejaculation and urine collection, variable urine volume and concentration, or other factors.

The median 'age at spermarche' observed in this cross sectional study was 14.1 years. In relation to pubertal stage, the 50% mark was between PH 2 and 3, although in a few cases spermaturia was found even in clinically prepubertal boys. Despite methodological and ethnic differences these results compare well with the findings of Richardson and Short who found a median age at spermarche of 13.3 years,² Hirsch *et al* (14.5 years),³ and Nielsen *et al* (13.4 years).⁵ It is also noteworthy that the median age at spermarche found in these studies corresponds to the age at first conscious ejaculation as evaluated by interviews.¹³

In summary, repetitive assessment of spermaturia is a sensitive, ethically acceptable, non-invasive procedure to determine the state of spermatogenic activity in pubertal subjects. The method cannot replace detailed quantitative sperm analysis, but may be used as a qualitative screening test to investigate potential damage to the maturing testis. It appears also suitable for epidemiological investigations in precocious or delayed puberty, for patients having chemotherapy or irradiation treatment, or for those with other chronic diseases potentially adversely affecting the gonads. Results of such studies can be analysed using the methods and be compared with the normal data presented in this study.

The authors thank Professor Michael Preece, Institute for Child Health, London, for his advice in the statistical analysis of the material.

- Baldwin B. The determination of sex maturation in boys by a laboratory method. *J Comp Psychol* 1928;8:39-43.
- Richardson D, Short R. Time of onset of sperm production in boys. *J Biosoc Sci* 1978;5:15-25.
- Hirsch M, Shemesh J, Modan M, Lunenfeld B. Emission of spermatozoa: age of onset. *Int J Androl* 1979;2:289-98.
- Hirsch M, Lunenfeld B, Modan M, Ovidia J, Shemesh J. Spermarche—the age of onset of sperm emission. *J Adolesc Health Care* 1985;6:35-9.
- Nielsen CT, Skakkebaek NS, Richardson DW, *et al*. Onset of the release of spermatozoa (spermarche) in boys in relation to age, testicular growth, pubic hair, and height. *J Clin Endocrinol Metab* 1986;62:532-5.
- Kulin HE, Frontera ME, Demers LD, Bartholomew MJ, Lloyd TA. The onset of sperm production in pubertal boys. *Am J Dis Child* 1989;143:190-3.
- Weissenberg R, Rozenman H, Hova M, Modan M, Lunenfeld B. The diagnostic value of sperm in post-ejaculatory urine. *Int J Androl* 1984;7:142-8.
- Tanner JM. *Growth at adolescence*. 2nd Ed. Oxford: Blackwell Scientific Publications, 1962.
- Kaplan EL, Meier P. Nonparametric estimation from incomplete observations. *Journal of the American Statistical Association* 1958;53:457-62.
- Finney DJ. *Probit analysis*. London: Cambridge University Press, 1971.
- SAS Institute Inc. *SAS user's guide: statistics version 5*. Cary, NC: SAS Institute, 1985.
- Charny CW, Conston AS, Meranze DR. Development of the testis. *Fertil Steril* 1952;3:461-79.
- Laron Z, Arad J, Gurewitz R, Grunebaum M, Dickerman Z. Age at first conscious ejaculation: a milestone in male puberty. *Helv Paediatr Acta* 1980;35:13-20.