

## Comparison of Radioimmunoassay and Enzyme Immunoassay Methods in Determining Gentamicin Pharmacokinetic Parameters and Dosages

J. C. ROTSCHAFFER, C. MORLOCK, L. STRAND, AND K. CROSSLEY

*Section of Clinical Pharmacology and Department of Internal Medicine, St. Paul-Ramsey Medical Center, St. Paul, Minnesota 55101, and College of Pharmacy and Department of Medicine, University of Minnesota Medical School, University of Minnesota, Minneapolis, Minnesota 55455*

Volume 22, no. 4, p. 650, column 1, Table 1: The line equations for groups I and III have been transposed. The correct equations are as follows: group I,  $y = 0.98x - 0.06$ ; group III,  $y = 0.86x - 0.19$ .

---

## Lysis of *Trypanosoma cruzi* by *Pseudomonas fluorescens*

TERESA I. MERCADO AND ALBA COLÓN-WHITT

*Laboratory of Parasitic Diseases and Laboratory of Streptococcal Diseases, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, Maryland 20205*

Volume 22, no. 6, p. 1051, column 2, lines 31–33: “. . . of the agar slants (optical density at 540 nm adjusted to 100 Klett units). . . .” should read “. . . of the agar slants and incubated for 24 h at 26°C with constant shaking. The 100-ml culture volumes were transferred to each of 8 liters of defined medium (optical density at 540 nm adjusted to 100 Klett units). . . .”

Page 1051, column 2, line 39: “. . . 200 ml of the same medium. . . .” should read “. . . 200 ml of the same medium (optical density at 540 nm adjusted to 600 Klett units). . . .”