

ERRATA

In Vitro Activity of a Group of Broad-Spectrum Cephalosporins and Other β -Lactam Antibiotics Against *Chlamydia trachomatis*

MARGARET R. HAMMERSCHLAG AND AIDA GLEYZER

Division of Infectious Diseases and Immunology, Department of Pediatrics, Downstate Medical Center, State University of New York, Brooklyn, New York 11203

Volume 23, no. 3, p. 493, abstract: "The activities of seven broad spectrum cephalosporins, four other β -lactam antibiotics, and one monobactam. . . ." should read "The activities of seven broad-spectrum cephalosporins and five other β -lactam antibiotics. . . ."

Page 493, column 1, lines 12-14: ". . . we tested seven of the new broad-spectrum cephalosporins, four other β -lactam antibiotics, and one monobactam. . . ." should read ". . . we tested seven of the new broad-spectrum cephalosporins and five other β -lactam antibiotics. . . ."

Page 493, column 2, lines 3-6: ". . . and the monobactam Sch 29,482. Sch 29,482, an oral penem highly resistant to β -lactamase, has been reported to be highly active. . . ." should read ". . . and Sch 29,482, an oral penem highly resistant to β -lactamase and reportedly highly active. . . ."

Page 494, column 1, lines 28-33: "The only other monobactam that has been tested against *C. trachomatis* is SQ 26,776 (5), which was reported to be inactive, with ≥ 100 $\mu\text{g/ml}$ needed to suppress normal inclusions. We found Sch 29,482 to be more active, with a mean MIC of 28 $\mu\text{g/ml}$ " should read "The mean MIC of Sch 29,482 was 28 $\mu\text{g/ml}$."

Incorporation of 5-Substituted Analogs of Deoxycytidine into DNA of Herpes Simplex Virus-Infected or -Transformed Cells Without Deamination to the Thymidine Analog

LAWRENCE FOX, MICHAEL J. DOBERSEN, AND SHELDON GREER

Departments of Microbiology, Biochemistry, and Oncology, University of Miami, School of Medicine, Miami, Florida 33101

Volume 23, no. 3, p. 465: The address should appear as shown above.

Page 474, column 1: Lines 18-27 should read "In the studies of Creasey (13), 5-methylcytidine and 5-methyl(dC) were shown to be moderate substrates for cytidine deaminase. Maley (29) has shown that 5-methyldeoxycytidylate has the lowest K_m value with respect to dCMP deaminase of chick embryo and monkey kidney cells among several analogs examined; 5-methyl(dCMP) had a lower K_m value than dCMP."