

		Size 0 (No Mutation)		Size 1 – 39		Size 40+	
		No.	%	No.	%	No.	%
Treatment	S9333	67	65%	9	38%	9	39%
Study	S9500	36	35%	15	63%	14	61%
Sex	Female	51	40%	13	52%	11	50%
	Male	52	50%	12	48%	11	50%
Race	Asian	2	2%	1	4%	1	5%
	Black	10	10%	2	8%	4	18%
	White	91	88%	21	84%	17	77%
	Unknown	0	0%	1	4%	0	0%
AML Onset (S9333 Only)	De novo	55	82%	10	100%	6	75%
	Secondary	12	18%	0	0%	2	25%
FAB (Local)	M1	20	19%	3	12%	7	32%
	M2	42	41%	6	24%	4	18%
	M4	27	26%	8	32%	6	27%
	M5	6	6%	5	20%	4	18%
	M6	0	0%	0	0%	0	0%
	M7	1	1%	0	0%	0	0%
	M0	6	6%	2	8%	1	5%
	Other AML	1	1%	1	4%	0	0%
	AML NOS	0	0%	0	0%	0	0%
Evaluable Cytogenetics?	Yes	87	84%	14	56%	13	59%
	No	16	16%	11	44%	9	41%
Normal Cytogenetics?	Yes	30	34%	12	86%	10	77%
	No	57	66%	2	14%	3	23%
Cytogenetic Risk Group	Favorable	11	13%	0	0%	1	8%
	Intermediate	39	45%	12	86%	11	85%
	Unfavorable	35	40%	2	14%	1	8%
	Indeterminate	2	2%	0	0%	0	0%
ITD/WT Ratio	< 0.836	--	--	11	46%	12	52%
	≥ 0.836	--	--	13	54%	11	48%

	Size 0 (No Mutation)		Size 1 – 39		Size 40+	
	Median	Range	Median	Range	Median	Range
Age	61	19 – 84	50	18 – 82	46	33 – 76
Marrow blasts (%)	70	19 – 99	80	32 – 99	79	54 – 95
WBC (1000/ μ L)	26.5	0.8 – 263.2	37.2	2.9 – 180.3	48.5	9.3 – 272.5
Peripheral blasts (%)	44	0 – 99	34	0 – 87	62	3 – 97
ITD/WT ratio	--	--	0.92	0.07 – 0.27	0.80	0.05 – 0.55