

Table 2. Regions showing simple effects of placebo during heat or anticipation

Region	x	y	z	Vol. (mm³)	Max T	SVC t- threshold	Correlation with RP
Placebo effects in heat ([P - C] Heat)							
<i>Increases</i>							
Ramy	-25	-2	-27	23	3.53	3.15	-0.61 *
Rmofc	-14	54	-20	11	3.90	3.22	-0.78 **
Lmofc	7	61	-20	46	3.42	3.22	0.30
LaOFC	27	56	-14	900	6.51	3.47	0.58 +
RaOFC	-29	58	-9	296	4.27	3.36	-0.44
RIOFC	-50	25	-14	34	2.97	2.60	-0.73 **
LIOFC	50	29	-14	34	3.21	2.74	-0.32
<i>Decreases</i>							
	-14	7	11	444	-2.93	2.12	-0.45
	38	11	56	376	-2.96	2.33	-0.53
Placebo effects in anticipation ([P - C] Warm)							
<i>Increases</i>							
No significant regions							
<i>Decreases</i>							
Lamy	29	0	-27	638	-3.68	2.73	0.33
LIOFC	36	25	-22	11	-3.05	2.75	0.32
Lvains	40	20	-20	11	-2.84	2.75	0.01
Lvains	43	18	-14	23	-2.85	2.75	-0.06
pag	4	-29	-4	57	-2.30	2.20	0.17
pgACC	-4	43	0	125	-2.70	2.57	0.11
Lcau	18	7	11	478	-2.31	1.71	0.41
Rifj	-43	2	25	159	-3.35	2.49	-0.47
RDLPFC	-47	4	50	11	-2.51	2.49	-0.28
RDLPFC	-29	9	56	11	-2.16	2.13	-0.51 +

Warm stimulation always preceded hot stimulation, and so is referred to as 'anticipation.' Column labels are as in Supporting Information Table 1. +, $P < .10$; *, $P < .05$; **, $P < .01$; ***, $p < .005$, all two-tailed. L, left; R, right.