

Variable	Log(Bayes Factor)	Log(Bayes Factor) average	Effect (OR)	Effect (OR)	Referent group
ACS	85.76	84.79	1.18 (1.14;1.24)		no ACS
Age	110.43	84.79	2.67 (2.44;2.91) [18-40]	7.61 (5.11;1.34) [>40]	2--18 years
Bilirubin	345.09	295.56	1.43 (1.28;1.60) [1.3-3.4]	1.59 (1.02;2.49) [>3.4]	normal
Blood Transfusion	47.00	39.61	2.01 (1.78;2.27)		no chronic BT
LDH	196.90	157.39	0.85 (0.76;0.94) [<300]	1.1 (0.99;1.24) [>600]	normal
MCV	561.19	474.22	0.54 (0.49;0.60) [<80]	1.98 (1.71; 2.29) [>98]	normal
Pain	245.70	205.31	1.61 (1.45;1.77)		no pain
Priapism	42.07	38.67	1.35 (1.09;1.67)		no priapism
Reticulocytes	324.12	276.23	0.5 (0.44;0.55) [<4.8]	1.51 (1.39;1.65) [>13]	normal
Sepsis	447.15	367.48	67.19(57.66;78.29)		no sepsis
Sex	2302.11	1957.89	1.16 (1.08;1.25)		females
Stroke	129.49	111.53	3.81 (3.20;4.54)		no stroke
Sys BP	88.49	79.95	2.84 (1.67;4.82) [low]	3.41 (2.71;4.27) [high]	normal
WBC	5.01	3.40	1.37 (1.24;1.51) [10.8-13.5]	1.92 (1.44;2.55) [>13.5]	normal