

Supplemental table A. Clinicopathologic features of the chromosome translocations positive and negative groups

Factors	t(11;18)/ <i>API2-MALT1</i>			<i>IGH</i> involved translocation		
	Positive (n=18) (%)	Negative (n=69) (%)	p Value	Positive (n=4) (%)	Negative (n=64) (%)	p Value
Age (yr)	16-80	18-80		40-62	16-80	
Mean,	53	59	0.064 ^a	53	58	NS ^a
Gender						
Male	12 (67)	33 (48)	NS ^b	2 (50)	32 (50)	NS ^b
Female	6 (33)	36 (52)		2 (50)	32 (50)	
Depth of tumour invasion						
Mucosa	7 (39)	28 (41)	NS ^b	2 (50)	26 (41)	NS ^b
Submucosa or beyond	11 (61)	41 (59)		2 (50)	38 (59)	
Clinical stage						
I	15 (83)	44 (64)	NS ^b	3 (75)	42 (66)	NS ^b
II ₁ or more	3 (17)	25 (36)		1 (25)	22 (34)	
<i>H pylori</i> status	(n=16)	(n=59)		(n=2)	(n=57)	
Positive	10 (63)	57 (97)	<0.001 ^b	2 (100)	49 (86)	NS ^b
Negative	6 (38)	2 (3)		0	8 (14)	
Response to eradication	(n=6)	(n=28)		(n=2)	(n=30)	
Complete remission (CR)	1 (17)	19 (68)	0.032 ^b	1 (50)	16 (53)	NS ^b
Non-CR	5 (83)	9 (32)		1 (50)	14 (47)	
Bcl10 nuclear expression	(n=14)	(n=66)			(n=58)	
Positive	10 (73)	26 (39)	0.029 ^b	2 (50)	29 (50)	NS ^b
Negative	4 (27)	40 (61)		2 (50)	29 (50)	

NS, not significant; CR, complete remission. ^aMann-Whitney *U* test. ^bFisher's exact probability test.