

Table S1. A summary of proteins tested for involvement in PC formation, siRNA target sequences, and methods used to monitor successful depletion

Protein	siRNA oligonucleotides	≥50% depletion monitored via
ALMS1	5'-CCCTATGATGACTGTCATAAA-3' (342), 5'-CAGAACTTTACCTGATGAA-3' (343)	IF
BBS4	5'-AAGGCACAAGACCAGTTGCAC-3'	Kim et al., 2004
CAP350	5'-ATGAACGATATCAGTGCTATA-3'	Yan et al., 2006
CCCAP	5'-GACCTGTTCATCTATTAAAT-3' (288), 5'-GAGGAACGACTTAGCTGAATA-3' (289)	qRT-PCR
centrin-2	5'-AAGAGCAAAAGCAGGAGATCC-3'	Salisbury et al., 2002
centrin-3	5'-CTGGTGACATTTAAGAATTA-3'	IF
Cep131	5'-CCCACTCAGCCCCGAAACAATA-3' (272), 5'-CAGCACGAGCTGGAGATAAA-3' (273)	qRT-PCR
Cep135	5'-AAAGCTTATTGCTCATTTAAA-3' (274), 5'-AAGCAGATTGAGCTAAGAGAA-3' (275)	qRT-PCR, IF
Cep152	5'-GCGGATCCAACGTGAAATCTA-3' (277)	qRT-PCR
Cep164	5'-CAGGTGACATTTACTATTTCATA-3' (278), 5'-ACCACTGGGAATAGAAGACAA-3' (279)	qRT-PCR, IF
Cep170	5'-GAAGGAATCCTCCAAGTCA-3'	Guarguaglini et al., 2005
Cep192	5'-AACAGTGAATGTGCAAGTAAA-3' (281)	qRT-PCR
Cep215	5'-GTGGAAGATCTCTAACTAA-3' (283)	qRT-PCR, IF
Cep27	5'-CAGCAGATCACAAATATTCAA-3' (254), 5'-CTGAAGAAAGTTATCTTATA-3' (255)	qRT-PCR
Cep290	5'-TAGCCTCGAAAGACTAGTAA-3' (285)	qRT-PCR, IF
Cep41	5'-TCCATGTTAGATGGACAGAAA-3' (256), 5'-CACTGGTAACAGTATGACTAA-3' (257)	qRT-PCR
Cep57	5'-AACCAAATAACTAAAGTCGA-3' (258), 5'-AGCCATCAAGGTCTAATGGAA-3' (259)	qRT-PCR
Cep63	5'-CAACGCTTGTATTATCAGCAA-3' (260), 5'-CAGCCTTAAAGTAGCCTCTTA-3' (261)	qRT-PCR
Cep68	5'-CACCCCTCAAATCACCTACTAA-3' (262), 5'-ACCGAAGATGATCCATCCCTA-3' (263)	qRT-PCR, IF
Cep70	5'-CCGGCGACTGAGTAAGATGAA-3' (264), 5'-GAAGATCGCATTGTCACTCAA-3' (265)	qRT-PCR
Cep72	5'-AGAGCTATGTATGATAATTAA-3' (266)	qRT-PCR
Cep76	5'-CTCGGTATTATAGGGCCAATA-3' (268), 5'-AAGGGTTTGAGGCCCTAGACAA-3' (269)	qRT-PCR
Cep78	5'-CAGTTGTGAAAGCTTAA-3' (270), 5'-TACACACTATTGGTGACCTA-3' (271)	qRT-PCR
chTOG	5'-AAGAGCCCAGAGTGGTCCAAA-3'	Cassimeris and Morabito, 2004
C-Nap1	5'-CTGGAAGAGCGCTAACTGAT-3'	Bahe et al., 2005
CP110	5'-CCCGAAATTATGCCAAAGTTA-3' (290), 5'-TAGACTTATGCAGACAGATAA-3' (291)	qRT-PCR
CPAP	5'-AAGGAAGATTGACCAAGTCAA-3' (250), 5'-CCCAATGGAACCTCGAAAGGAA-3' (251)	qRT-PCR, IF
Dynamin2	5'-AAGAGTCGGTGCTCGAGAAC-3'	Thompson et al., 2004
FOP*	5'-AAGTGATCAGGCGCTGTCAC-3'	Yan et al., 2006
hSAS6	5'-AAGCACGTTAACGCTACAA-3'	Leidel et al., 2005
NA14	5'-TGGCCTGTGATTATGAATAAA-3' (286), 5'-CTCCTTCACCACAGAACCAA-3' (287)	qRT-PCR
Nek2	5'-AACATGAGATGTGACATTCTA-3' (252), 5'-TTACGAGGATGTTAAACTTAA-3' (253)	qRT-PCR, IF
ninein	5'-GCGGAGCTCTGAAGTTAAA-3'	IF
Nlp	5'-CACACTGAAATTGACTATA-3' (249)	IF
Odf2	5'-AAGGATCTTATGTCGCTGAA-3'	IF
OFD-1	5'-CTCAGACAAGTCGACATTAA-3' (339)	IF
PCM-1	5'-AATCAGCTTCGTGATTCTCAG-3'	Dammermann and Merdes, 2002
pericentrin*	5'-AAGCAGCTGAGCTGAAGGAGA-3'	Dammermann and Merdes, 2002
Plk4	5'-CTGGTAGTACTAGTTCACCTA-3'	Habedanck et al., 2005
rootletin*	5'-AAGCCAGTCTAGACAAGGA-3'	Bahe et al., 2005
Sfi1p	5'-CTGCTGTGCAAATGTATCGAA-3'	qRT-PCR, IF

siRNA oligonucleotides were from QIAGEN unless marked by \*, in which case they were from Dharmacon. IF, immunofluorescence microscopy; qRT-PCR, quantitative RT-PCR.

## References

- Cassimeris, L., and J. Morabito. 2004. TOGp, the human homolog of XMAP215/Dis1, is required for centrosome integrity, spindle pole organization, and bipolar spindle assembly. *Mol. Biol. Cell.* 15:1580–1590.
- Dammermann, A., and A. Merdes. 2002. Assembly of centrosomal proteins and microtubule organization depends on PCM-1. *J. Cell Biol.* 159:255–266.
- Kim, J.C., J.L. Badano, S. Sibold, M.A. Esmail, J. Hill, B.E. Hoskins, C.C. Leitch, K. Venner, S.J. Ansley, A.J. Ross, et al. 2004. The Bardet-Biedl protein BBS4 targets cargo to the pericentriolar region and is required for microtubule anchoring and cell cycle progression. *Nat. Genet.* 36:462–470.
- Leidel, S., M. Delattre, L. Cerutti, K. Baumer, and P. Gonczy. 2005. SAS-6 defines a protein family required for centrosome duplication in *C. elegans* and in human cells. *Nat. Cell Biol.* 7:115–125.
- Thompson, H.M., H. Cao, J. Chen, U. Euteneuer, and M.A. McNiven. 2004. Dynamin 2 binds gamma-tubulin and participates in centrosome cohesion. *Nat. Cell Biol.* 6:335–342.
- Yan, X., R. Habedanck, and E.A. Nigg. 2006. A complex of two centrosomal proteins, CAP350 and FOP, cooperates with EB1 in microtubule anchoring. *Mol. Biol. Cell.* 17:634–644.