

# Monoclonal Antibodies to a Membrane Glycoprotein Induce the Phosphorylation of Histone H1 in Sea Urchin Spermatozoa

Victor D. Vacquier, Gary W. Moy, James S. Trimmer, Yukiko Ebina, and Donald C. Porter

Marine Biology Research Division, A-002, Scripps Institution of Oceanography, University of California, San Diego, La Jolla, California 92093

**Abstract.** Two groups of mAbs reacting with external domains of a major sea urchin sperm membrane glycoprotein of 210 kD were isolated. Previous studies have shown that group I mAbs inhibit the acrosome reaction induced by egg jelly and also cause large increases in intracellular  $\text{Ca}^{2+}$  ( $[\text{Ca}^{2+}]_i$ ). Group II mAbs, at comparable levels of cell surface binding, neither inhibit the egg jelly-induced acrosome reaction nor cause increases in  $[\text{Ca}^{2+}]_i$ . In this paper, we investigate the ability of these mAbs to induce the cAMP-dependent phosphorylation of sperm histone H1. Group I mAbs induce H1 phosphorylation to the same level and on the same peptide, as occurs upon treatment of sperm with egg jelly. These mAbs also activate

adenylate cyclase to the same extent as egg jelly. Group II mAbs do not induce H1 phosphorylation and are only poor activators of adenylate cyclase. Group I mAbs compete with each other, but not with group II mAbs, for binding to the cell surface. These data indicate that the activation of adenylate cyclase is an initial event in the pathway leading from the binding of mAbs to a specific domain of the 210-kD protein at the cell surface, to the discrete phosphorylation of histone H1 in highly condensed sperm chromatin. The domain on the 210-kD protein recognized by group I mAbs plays a critical role in signal transduction during the early events of fertilization.

SEA urchin eggs are covered by a jelly layer that can be solubilized by brief exposure to pH 5 seawater. The major macromolecule of egg jelly is a fucose sulfate-rich glycoconjugate (FSG)<sup>1</sup> that induces the acrosome reaction of sperm (SeGall and Lennarz, 1979, 1981; Garbers et al., 1983; DeAngelis and Glabe, 1987). This morphological change consists of the exocytosis of the acrosome granule and the polymerization of actin to form the acrosomal process, which is covered by the membrane destined to fuse with the egg (reviewed in Lopo, 1983; Tilney, 1985). FSG has profound physiological effects on sperm, including the opening of ion channels, resulting in the net influx of  $\text{Ca}^{2+}$  and  $\text{Na}^+$ , the net efflux of  $\text{K}^+$  and  $\text{H}^+$ , and the depolarization of the  $\text{K}^+$ -supported membrane potential (reviewed by Schackmann, 1986, 1988). In addition to these ionic events, FSG induces the activation of adenylate cyclase and protein kinase, the elevation of cAMP concentrations (reviewed in Garbers and Kopf, 1980), and the elevation of inositol

trisphosphate (Domino and Garbers, 1988). These effects of FSG are all dependent on the presence of external  $\text{Ca}^{2+}$ . The major substrate phosphorylated *in vivo* by the FSG-induced increases in cAMP-dependent protein kinase activity is sperm-specific histone H1 (Porter and Vacquier, 1986), which is phosphorylated on a single seryl site in the peptide RKGS(PO<sub>4</sub>)SNAR (Porter et al., 1988a). The phosphorylation of H1 on this single site in response to FSG may be a prerequisite for the hyperphosphorylation of H1 by egg kinases and its loss from sperm chromatin within 10 min after the sperm nucleus enters egg cytoplasm (Green and Poccia, 1985; Porter and Vacquier, 1988; Porter et al., 1988b).

These effects of FSG on sperm are complex, and the relationships between the changes in ionic permeability and the mobilization of the cyclic nucleotide cascade leading to H1 phosphorylation are not well understood. Although it has never been rigorously demonstrated, these effects of FSG on sperm must involve interaction with extracellular domains of cell surface proteins. In an attempt to identify sperm surface proteins involved in the recognition of FSG and the transduction of this signal across the plasma membrane, mAbs were isolated that react with extracellular domains of sperm membrane proteins (Trimmer, 1987). The major antigenic determinant on the sperm surface is a glycoprotein of 210–260 kD (the 210-kD protein). Two groups of mAbs reacting mono-

J. S. Trimmer's present address is Department of Physiology, Yale University School of Medicine, New Haven, CT 06510. Y. Ebina's present address is Central Research Laboratory, Fukushima Medical College, Fukushima, Japan. D. C. Porter's present address is Cancer Center, T-011, University of California, San Diego, La Jolla, CA 92093.

1. *Abbreviations used in this paper:* FSG, fucose sulfate glycoconjugate of egg jelly; WGA, wheat germ agglutinin.

specifically with this protein were isolated. Group I mAbs inhibit the FSG-induced acrosome reaction (Trimmer et al., 1985; Trimmer and Vacquier, 1986), yet induce large increases in intracellular  $\text{Ca}^{2+}$  (Trimmer et al., 1986). Group II mAbs do not exhibit these effects on sperm. Here we show that the group I mAbs, but not the group II mAbs, are potent inducers of adenylate cyclase and the phosphorylation of histone H1. This indicates that the 210-kD protein mediates the activation of sperm adenylate cyclase. The domain of the 210-kD protein recognized by group I mAbs must be critical for the interaction of FSG with sperm during fertilization.

## Materials and Methods

### mAbs

The production of mouse mAbs reacting with external domains of sea urchin sperm plasma membrane proteins has been described in detail (Trimmer et al., 1985, 1986, 1987; Trimmer, 1987; and Trimmer and Vacquier, 1988). This description includes preparation of the immunogen, screening by ELISA, labeling sperm with  $^{125}\text{I}$ , detergent extraction of labeled sperm, immunoprecipitation of labeled protein, gel electrophoresis of immunoprecipitates and autoradiography, production of ascites fluid, isotyping of immunoglobulins, Fab fragment production,  $^{125}\text{I}$  labeling of antibodies, and competitive binding experiments. In the experiments reported in this paper, mAb preparations used were IgG partially purified from ascites fluid by two cycles of ammonium sulfate precipitation by standard methods (Garvey et al., 1977) followed by dialysis into seawater and determination of protein concentration. mAb T8/40 reacts with an unknown human lymphocyte antigen and was a generous gift of Dr. I. S. Trowbridge of the Salk Institute, La Jolla, CA.

### Gametes and the Induction of H1 Phosphorylation by Egg Jelly and mAbs

Gametes of *Strongylocentrotus purpuratus* were obtained by injection of adults with 0.5 M KCl. We have previously shown that the macromolecular fraction of egg jelly containing FSG, and not the peptide-containing fraction, is the inducer of H1 phosphorylation (Porter and Vacquier, 1986). Preparation and quantitation of egg jelly, collection and storage of sperm at 0°C as undiluted semen (1  $\mu\text{l}$  = 100  $\mu\text{g}$  protein and  $4 \times 10^7$  cells), in vivo labeling of cells with  $^{32}\text{P}$ , dilution of the cells into test substances, preparation of the cells for PAGE, and autoradiography were as described (Porter and Vacquier, 1986). All filtered seawater was buffered at pH 8.0 with 10 mM Hepes. Briefly, carrier-free  $\text{H}_3^{32}\text{PO}_4$  (ICN Biochemicals, Irvine, CA) was added directly to undiluted semen to 2 mCi/ml and the tube was rotated at 1 rpm (15°C) for 2 h. Seawater was then added to achieve a 1:10 dilution of the semen and 25- $\mu\text{l}$  samples (250  $\mu\text{g}$  sperm protein) were added to 1 ml seawater containing mAb or other test substances (15°C). The final dilution of the semen was 1:400. At 10 min, the sample was made 10% in TCA and then prepared for PAGE in SDS and  $\beta$ -mercaptoethanol (Laemmli, 1970; Porter and Vacquier, 1986).

### Miscellaneous Procedures

Analysis of the site on H1 phosphorylated by the H1 kinase (Fig. 4) was performed exactly as described (Porter et al., 1988a). The determination of the binding of mAb to fixed cells was a modification of previously described methods (Trimmer et al., 1985, 1987). Sperm were fixed in 3% paraformaldehyde, 0.1% glutaraldehyde in seawater. Fixed cells were collected by centrifugation at 1,000 g for 5 min and washed twice in seawater containing 0.1 M glycine and 1 mg/ml BSA. A stock suspension of fixed cells was prepared in NaCl buffer (150 mM NaCl, 10 mM  $\text{NaN}_3$ , 10 mM Tris pH 7.5, 1 mg/ml BSA:Millipore filtered) at  $4 \times 10^9$  cells per ml. 100- $\mu\text{l}$  portions of various antibody dilutions (0–200  $\mu\text{g}/\text{ml}$ ) were placed in 12  $\times$  75 mm borosilicate tubes, 5  $\mu\text{l}$  stock sperm suspension was added, and slow rotary motion was applied for 45 min. 2 ml NaCl buffer was then added, the tubes were centrifuged at 1,000 g for 5 min and the supernatant was carefully aspirated away. After three such washes, 100  $\mu\text{l}$  of a 1:200 dilution of rhodamine-labeled, FC-specific goat anti-mouse IgG was added to the pellet. The second antibody had been absorbed overnight against an excess of fixed sperm and then centrifuged 30,000 g and the supernatant stored in 100- $\mu\text{l}$  aliquots at  $-70^\circ\text{C}$ . The cells were gently agitated for 45 min and then

washed as given above three times in 2-ml portions of NaCl buffer as before. The cells were quantitatively transferred to a new tube and pelleted. After the supernatant was removed, the cells were resuspended in 3 ml NaCl buffer containing 1% sodium dodecyl sulfate and 1%  $\beta$ -mercaptoethanol and heated at 60°C for 30 min. After centrifugation at 1,000 g for 20 min, 2 ml of the supernatant containing the denatured rhodamine-conjugated antibody was removed and the fluorescence was determined on a double-beam fluorescence spectrophotometer with excitation at 540 nm and emission at 570 nm. The percentage saturation of binding sites on sperm of the mAbs was calculated in arbitrary units of fluorescence. The competition binding experiment (Fig. 5) was done with living cells exactly as described previously (Trimmer et al., 1987). For the sequential treatment of cells with two mAbs (Fig. 6), 25- $\mu\text{l}$  portions of  $^{32}\text{P}$ -labeled sperm (a 1:10 dilution of semen) were added to tubes containing 0.5 ml seawater with 50  $\mu\text{g}$  J18/2 (15°C). At 5 min, 0.5 ml seawater containing either J10/14 or J16/5 IgG at concentrations between 0 and 200  $\mu\text{g}/\text{ml}$  was added and the tubes were incubated 10 min before being fixed in TCA. To determine the effects of mAbs on the egg jelly-induced phosphorylation of H1 (Fig. 7), sperm were treated with either seawater or seawater containing mAb J4/4 or J18/2 at 100  $\mu\text{g}/\text{ml}$  (the  $^{32}\text{P}$ -labeled semen was diluted 200-fold); after 5 min, 0.5-ml portions were added to 0.5 ml of egg jelly at concentrations ranging from 0 to 200  $\mu\text{g}$  fucose per milliliter. After 10 min, the cells were fixed in trichloroacetic acid. Quantitating the agglutination of fixed sperm by mAb and determination of mAb-induced inhibition of the egg jelly-induced acrosome reaction of living sperm were performed as described (Podell and Vacquier, 1984).


### Adenylate Cyclase

The method described by Schultz and Jakobs (1983) was modified as follows. Buffer A contained 90 mM Hepes, pH 7.5, 6.6 mM cAMP, 3 mM ATP, 15 mM  $\text{MnCl}_2$ , and 60 mM  $\text{NaN}_3$ . Buffer B contained 10 mM NaCl, 10 mM KCl, 2 mM Hepes (pH 7.5), 20 mM  $\text{NaN}_3$ , 300  $\mu\text{g}/\text{ml}$  creatine kinase, and 45 mM creatine phosphate (pH 7.5). Buffer C contained 10 mM NaCl, 10 mM KCl, 2 mM Hepes, 5  $\mu\text{M}$  EDTA, 0.1 mM DTT, 20 mM  $\text{NaN}_3$ , and 12 mg/ml BSA. 1-ml aliquots of all three buffers were stored at  $-70^\circ\text{C}$ . One part buffer A was mixed with one part buffer B and the isotope ( $[\alpha\text{-}^{32}\text{P}]\text{ATP}$ , [ICN Biochemicals]) was added (1  $\mu\text{Ci}/84 \mu\text{l}$  A+B). Usually, 750  $\mu\text{l}$  of this mixture was placed in a 1.5-ml Eppendorf tube in a 23°C water bath. The cell lysate was then prepared exactly as follows. A 1-ml portion of seawater, containing egg jelly (185  $\mu\text{g}$  fucose/ml) or mAb was placed in Eppendorf tubes at 23°C. A fresh 1:10 dilution of semen (dry sperm) was made, 60  $\mu\text{l}$  (600  $\mu\text{g}$  protein) was added, and the tube was capped and inverted. At 120 s, the tube was microfuged for 45 s, the supernatant was immediately removed by aspiration and 1 ml 0°C seawater was added, the cell pellet was resuspended with a 1-ml disposable pipette tip, the cells were repelleted by 45 s of centrifugation, the supernatant was carefully removed, and 500  $\mu\text{l}$  buffer C (0°C) was added to lyse the cells. The lysate was pulled up several times in the pipette tip, and at exactly 8 min after exposure of the cells to the test medium, 375  $\mu\text{l}$  of lysate was added to 750  $\mu\text{l}$  of reaction mixture. Every 3 min, 120  $\mu\text{l}$  of reaction mixture ( $\sim 36 \mu\text{g}$  sperm protein) was removed and added to an Eppendorf tube containing 480  $\mu\text{l}$  of 0.2 M zinc acetate, followed by 600  $\mu\text{l}$  0.2 M  $\text{Na}_2\text{CO}_3$ . The tubes were capped, vortexed, and either stored frozen or centrifuged 5 min at top speed in a microfuge. 1 ml of supernatant was put over 3-ml columns of neutral alumina equilibrated in 100 mM Tris pH 7.5. The columns were washed into scintillation vials with two portions of 3 ml Tris buffer and the Cerenkov radiation was determined in a scintillation counter.  $[\text{H}]\text{cAMP}$  was used to monitor recovery of  $[\text{H}]\text{cAMP}$  formed; recovery was virtually 100%. No  $[\text{H}]\text{ATP}$  substrate was detectable in the column eluates. Data points were linear from 3 to 18 min and activity was proportional to added sperm protein. Phosphodiesterase inhibitors were not included in the assay.

## Results

### Characterization of mAbs

The group I mAbs, J10/14 (IgG<sub>2a</sub>), J18/2 (IgG<sub>3</sub>), J16/5 (IgG<sub>1</sub>), and J17/10 (IgG<sub>2a</sub>); and group II mAbs, J4/4 (IgG<sub>2a</sub>) and J18/27 (IgG<sub>3</sub>), react monospecifically with the 210-kD protein, as shown by autoradiograms of SDS-PAGE analysis of immunoprecipitates of detergent extracts of  $^{125}\text{I}$ -labeled sperm proteins (Fig. 1). The lane of total detergent extract of radiolabeled sperm used as the starting material for the



**Figure 1.** Autoradiogram of immunoprecipitates of mAbs. All six mAbs react exclusively with the 210-kD protein. The total  $^{125}\text{I}$ -labeled sperm membrane protein used as the starting material for the immunoprecipitates appears in the far right lane with approximate relative mass given in kilodaltons (kD). Group I mAbs are J18/2, J10/14, J16/5, and J17/10. Group II mAbs are J4/4 and J18/27. T8/40 serves as a negative control.


immunoprecipitates shows the specificity of the mAbs for the 210-kD protein.

### ***mAb-induced Phosphorylation of H1***


To investigate the effects of these mAbs on the *in vivo* phosphorylation of H1, the ATP pools of sperm were labeled with  $^{32}\text{P}$  and the cells were diluted into seawater containing egg jelly or mAb. Group I mAbs (J10/14, J16/5, and J17/10) at 100  $\mu\text{g}/\text{ml}$  induced H1 phosphorylation to the same extent as did egg jelly (Fig. 2). Group II mAbs (J4/4 and J18/27), and the negative control mAb T8/40, were noninductive at concentrations from 5 to 400  $\mu\text{g}/\text{ml}$  (Fig. 2). Both J10/14 IgG and Fab fragments were inductive at concentrations as low as 5  $\mu\text{g}/\text{ml}$ . mAb J18/2 (group I) was unique in that at high concentrations (100–200  $\mu\text{g}/\text{ml}$ , Fig. 2) it was noninductive, but at lower concentrations it induced H1 phosphorylation to the same extent as seen upon treatment of sperm with egg jelly (Fig. 3). In this same experiment, mAb J17/10 (group I) was inductive to 200  $\mu\text{g}/\text{ml}$ , whereas J18/27 (group II) was noninductive at all concentrations tested.

### ***H1 Site Phosphorylated by mAb Treatment***


Previous chemical analysis for phosphate of purified H1 had



**Figure 2.** mAb-induced phosphorylation of histone H1. Total sperm protein (35  $\mu\text{g}$ ) stained with Coomassie brilliant blue is shown in the left two lanes. T, tubulin; H1, histone H1; CH, core histones. The nine lanes to the right of the protein stain are autoradiograms of total sperm protein (35  $\mu\text{g}$ ). +Jelly, whole egg jelly at 150  $\mu\text{g}$  fucose per ml. Sperm were exposed for 10 min to 100  $\mu\text{g}$  mAb/ml before fixation. Note that J18/2 did not induce H1 phosphorylation.



**Figure 3.** An autoradiogram showing that high concentrations of J18/2 do not induce H1 phosphorylation, whereas lower concentrations are inductive. In this experiment, J18/2 at 25  $\mu\text{g}/\text{ml}$  was as inductive as egg jelly, but with increasing concentration of this mAb, less phosphorylation of H1 occurred, whereas with J17/10, (and also J10/14 and J16/5) this decrease in H1 phosphorylation with increasing mAb concentration was not observed. Only the H1 (arrow) portion of the autoradiogram is presented. mAb concentrations in micrograms per milliliter are shown over each lane. J18/27 was noninductive from 25 to 400  $\mu\text{g}/\text{ml}$ . +Jelly, positive control; -Jelly, negative control.



**Figure 4.** Two-dimensional separation of the Endo Arg C digest of H1 phosphorylated by mAb J10/14 treatment of sperm. Procedures were exactly as previously described (Porter et al., 1988a). This autoradiogram of the thin-layer plate (20 × 20 cm) shows one major labeled peptide. o, origin; E, direction of first dimension electrophoresis; C, direction of second-dimensional chromatography.

shown that before treatment with egg jelly there is 0.16 mol phosphate/mol protein, and that after egg jelly treatment this number increases to 0.46 (Porter and Vacquier, 1986). Two-dimensional, thin-layer separation of Endo Arg C digests of the NH<sub>2</sub>-terminal CNBr-generated fragment of H1 yielded only one labeled peptide (Porter et al., 1988a). Mixing experiments showed that this same peptide was the only one labeled *in vivo* when the H1 kinase was activated by treating sperm with the phosphodiesterase inhibitor 3-isobutyl-1-methylxanthine, or when sperm were permeabilized with detergent in the presence of [ $\gamma$ -<sup>32</sup>P]ATP and cAMP. The sperm H1 kinase is cAMP dependent and the sequence of the phosphorylated peptide (RKGS(PO<sub>4</sub>)SNAR) is a preferential site for cAMP-dependent protein kinase (Porter et al., 1988a). To test whether this same site was phosphorylated in response to mAb treatment, the <sup>32</sup>P-labeled histone H1 was purified from mAb J10/14-treated sperm and the two-dimensional Endo Arg C maps were prepared. The same <sup>32</sup>P-labeled peptide previously reported as being labeled upon egg jelly treatment was the only major spot appearing on the autoradiogram of the peptide map (Fig. 4). Thus, in the highly condensed native sperm chromatin, the endogenous cAMP-dependent protein kinase phosphorylates H1 on the same peptide, whether activated by egg jelly, phosphodiesterase inhibitors, or mAbs.

#### **Binding of mAbs to the Sperm Surface**

To test if the failure of mAbs J4/4 and J18/27 to induce the phosphorylation of H1 was due to a lack of sperm surface binding, binding analyses were performed under the conditions used for the phosphorylation experiments. We previously reported that J4/4 saturates all binding sites at ~125  $\mu$ g/ml when the sperm concentration is  $4 \times 10^8$  cells/ml

(Trimmer et al., 1985). We observed no phosphorylation of H1 in J4/4 at 200  $\mu$ g/ml and sperm concentrations of  $1 \times 10^8$  cells per ml. We also reported that J10/14 IgG saturates cell surface binding sites at ~12  $\mu$ g/ml, and that there is no difference in number of binding sites in living cells as compared with paraformaldehyde/glutaraldehyde fixed cells (Trimmer et al., 1985; Trimmer, 1987).

In Table I, we present data on the relative binding of these six anti-210-kD mAbs to sperm under incubation conditions equivalent to those used for the *in vivo* phosphorylation of H1 by mAbs (0–200  $\mu$ g/ml). Fixed cells were used in these experiments, as we have never observed a quantitative difference between the binding of mAb to fixed as opposed to living sperm (the hybridomas were selected in an ELISA using fixed cells [Trimmer et al., 1985]). Also, at subsaturating concentrations, mAbs J18/2 and J18/27 cause the 210-kD protein to be shed from the surfaces of living cells (Trimmer and Vacquier, 1988). The surface binding data are presented as the relative amount of mAb bound at 100  $\mu$ g/ml as a percentage of the amount bound at 200  $\mu$ g/ml (Table I). Using this cell surface binding assay, it is evident that at 100  $\mu$ g/ml, mAbs J18/27 and J10/14 are present at saturating concentrations, whereas J4/4, J18/2, J16/5, and J17/10 are slightly below saturation. The relative amount of fluorescent second antibody bound at 200  $\mu$ g mAb/ml was lowest for J4/4, being about one third to one fourth of the group I mAbs. However, the other group II mAb, J18/27, bound roughly the same amount of second antibody as the group I-treated cells (Table I). The differences in inductive ability exhibited by group I and group II mAbs thus are not due simply to differences in cell surface binding.


To study in more detail the binding of these mAbs and how it relates to the striking differences in their ability to induce H1 phosphorylation, we performed competitive binding analysis using <sup>125</sup>I-J10/14. The group I mAbs J16/5 and J18/2 compete as well as J10/14 itself for <sup>125</sup>I-J10/14 binding to cells, whereas the group II mAbs (J4/4 and J18/27) are non-competitive at all concentrations tested, to 250  $\mu$ g/ml (Fig. 5). It therefore appears as if the group I mAbs bind at or near the same site on the 210-kD protein, whereas the group II mAbs bind a different region(s) of the protein.

From the data presented thus far, it is clear that mAb J18/2 is unique among the group I mAbs in that it displays a narrow concentration dependence in its ability to induce H1 phosphorylation, even though it clearly binds to the 210-kD protein at high levels at the noninductive concentrations (Table

**Table I. Relative Binding of Anti-210-kD mAbs**


Group	mAb	A	B
		%	Relative fluorescence
II	J4/4	87	140
II	J18/27	100	620
I	J18/2	80	400
I	J10/14	100	460
I	J16/5	89	630
I	J17/10	96	590

Column A, relative amounts of mAb bound to fixed cells at 100  $\mu$ g mAb/ml, expressed as a percentage of the amount bound at 200  $\mu$ g mAb/ml. Column B, arbitrary units of fluorescent second antibody bound after treatment of cells with 200  $\mu$ g mAb/ml.




**Figure 5.** Competitive binding using <sup>125</sup>I-J10/14. <sup>125</sup>I-J10/14 (20 µg) was mixed with the indicated micrograms of each unlabeled antibody and then sperm were added. Bound J10/14 was determined with a gamma counter (Trimmer et al., 1987). Each point is the average of three samples.

I and Fig. 5). An experiment was performed to determine if J10/14 or J16/5 could override J18/2 to yield the maximum mAb-inducible level of H1 phosphorylation. Cells were diluted into J18/2 at 100 µg/ml and after 5 min, either J10/14 or J16/5 was added to concentrations ranging from 0 to 100 µg/ml (5 µg/ml J10/14 is fully inductive). At 100 µg/ml, J18/2 does not induce the translocation and shedding of surface proteins (Trimmer and Vacquier, 1988). The results (Fig. 6) show that J18/2 at 100 µg/ml induced only a small amount of H1 phosphorylation that was not increased by later addition of J10/14 or J16/5 (as might be expected from Fig. 5). Treatment of sperm with either 100 µg/ml J10/14 or J16/5 alone gave high levels of H1 phosphorylation, as did egg jelly (Fig. 6). These data indicate that the blocking of the common binding site for these three group I mAbs by noninductive



**Figure 6.** Treatment of cells with J18/2 followed in 5 min by various concentrations of J10/14 or J16/5. In the first 12 lanes, sperm were diluted into 100 µg/ml J18/2. At 5 min, portions were removed and added to either J10/14 or J16/5 at the micrograms per milliliter indicated over each lane. After 10 min of incubation, the cells were processed for electrophoresis and autoradiography. Arrow, histone H1. Treatment with J10/14 or J16/5 did not increase the level of H1 phosphorylation in J18/2-treated cells.



**Figure 7.** Effect of mAbs on the egg jelly-induced phosphorylation of H1. Sperm were diluted into seawater only, or seawater containing 100 µg/ml of either J4/4 or J18/2. After 5 min an equal volume of seawater containing egg jelly was added to yield the final micrograms per milliliter fucose values of FSG shown on top of figure. After 10 min, the cells were processed for autoradiography. J4/4 does not alter the concentration dependence of egg jelly-induced H1 phosphorylation, whereas J18/2 inhibits the phosphorylation of H1 by egg jelly.

concentrations of mAb J18/2 disrupts the ability of both J10/14 and J16/5 to induce H1 phosphorylation.

An additional experiment was performed to determine if pretreatment of cells with J18/2 would alter the concentration dependence of FSG-induced H1 phosphorylation. As shown in Fig. 7, pretreatment with 100 µg/ml J18/2 completely inhibits the FSG-induced phosphorylation of H1. In contrast, the binding of the group II mAb J4/4 has no effect on the concentration dependence of the FSG-induced phosphorylation, relative to FSG alone. These data indicate that the site recognized by the group I mAb plays a role not only in the mAb-induced phosphorylation, but also in the phosphorylation of H1 induced by FSG.

#### *mAb Activation of Adenylate Cyclase*

To determine the relationship between the activation of adenylate cyclase and H1 phosphorylation, sperm were treated with these six mAbs or egg jelly, washed, lysed, and the adenylate cyclase activity was determined. Six such experiments are presented in Table II, the enzyme rates being calculated from the linear slope of six time points. The fold increase in enzyme activity induced by egg jelly alone ranged from 2.49 to 13.45, with a mean of 6.79. The group I mAbs stimulate adenylate cyclase activity to the same level as does egg jelly, whereas the group II mAbs (and wheat germ agglutinin [WGA]) failed to induce these large increases in adenylate cyclase activity. We have previously shown that the phosphorylation of H1 by the endogenous protein kinase is cAMP dependent (Porter et al., 1988a). The ability of group I mAbs, but not group II mAbs, to activate adenylate cyclase is consistent with the idea that increased levels of cAMP activate the kinase phosphorylating histone H1 (Porter and Vacquier, 1986; Porter et al., 1988a).

#### *Discussion*

Sea urchin spermatozoa are ideal for studying the mechanism of signal transduction. They possess a large membrane

**Table II. Activation of Adenylate Cyclase with Egg Jelly and mAb**

Exp. No.	Group	mAb* or Jelly†	Fold increase in rate over no jelly‡	Percent of rate‡ of jelly-treated sample
1	I	Jelly	3.67	—
		J18/2	3.67	100
2	II	Jelly	7.46	—
		J4/4	1.45	19
		J10/14	7.58	102
3	II	Jelly	2.49	—
		J4/4	0.92	37
		J10/14	2.66	107
4	II	Jelly	9.81	—
		J18/27	0.51	5
		J16/5	11.57	118
		J17/10	7.30	74
5	I	Jelly	13.45	—
		WGA†	1.75	13
		J18/2	13.36	99
		J18/2*	11.57	86
6*	II	Jelly	3.84	—
		J4/4	1.79	47
		J18/27	0.67	17
		J18/2	4.07	106
		J10/14	4.47	116
		J16/5	3.94	102
		J17/10	3.94	102

\* mAb concentrations were 100 µg/ml, except in one J18/2 sample in exp. 5 and for all samples of exp. 6, in which the concentrations were 200 µg/ml.

† Egg jelly was 100 µg fucose/ml.

‡ Basal adenylate cyclase specific activity in lysates of sperm not treated with egg jelly was  $0.67 \pm 0.38$  nmol cAMP formed/min per mg protein ( $n = 7$ ).

§ Rates were calculated from the linear slopes of six time points over 18 min.

¶ WGA was 60 µg/ml, which yields 100% inhibition of the FSG-induced acrosome reaction (Podell and Vacquier, 1984).

surface area (31–35 µm<sup>2</sup>; Cross, 1983) relative to their volume (8 µm<sup>3</sup>; Christen et al., 1982; Lee et al., 1983), they can be obtained as pure cells with ease and in vast quantity, and several methods exist to isolate the plasma membrane (reviewed in Vacquier, 1987). Egg jelly and its FSG component can be obtained from eggs with ease and its effects on sperm occur synchronously in a time span of seconds. In addition to FSG, egg jelly contains such small peptides as resact (Bentley et al., 1987) and speract (Schackmann and Chock, 1986; reviewed in Garbers et al., 1987) that bind to specific sperm surface receptors and alter cyclic nucleotide metabolism and ion channel activity. G proteins have also been reported in sperm membranes (Kopf et al., 1986; Bentley et al., 1986), and most recently GTP has been shown to control K<sup>+</sup> channels in the flagellar membrane of sea urchin sperm (Lee, 1988). The FSG component of egg jelly is a very large, complex molecule (SeGall and Lennarz, 1979, 1981; Garbers et al., 1983; DeAngelis and Glabe, 1987), and the mechanism by which it interacts with the sperm surface remains unknown. Because of the complexity of FSG, classical methods to identify the membrane proteins with which it interacts have been untenable. One approach to the analysis

of such complex systems is to use antibodies as probes to identify proteins mediating signal transduction. Using this approach, we have shown that the 210-kD protein plays a critical role in transduction of the FSG response (Trimmer et al., 1985, 1986, 1987).

In this study, we show that the domain of the 210-kD protein defined by the group I mAbs activates adenylate cyclase and induces the cAMP-dependent phosphorylation of H1. We propose that these mAbs use the same pathway as does FSG in inducing these changes when sperm contact eggs. The precise mechanism by which binding of mAb to the group I domain activates adenylate cyclase is of general interest, and most likely involves a conformational change in the 210-kD protein that activates Ca<sup>2+</sup> channels (Trimmer et al., 1986). The activation of the cyclase is strictly dependent on the presence of external Ca<sup>2+</sup> (Garbers and Kopf, 1980). However, ionophore A23187 (2–40 µM) does not induce H1 phosphorylation in sperm (Porter and Vacquier, 1986), but such negative data are inconclusive because ionophores act nonspecifically on cellular compartments and thus alter intracellular Ca<sup>2+</sup> concentrations unphysiologically.

The three group I mAbs tested compete equally well for binding to cells (Fig. 5), and also show the same concentration dependencies for inhibition of the FSG-induced acrosome reaction (Fig. 4 in Trimmer and Vacquier, 1986). That these mAbs activate adenylate cyclase but do not induce the acrosome reaction indicates that the cAMP-dependent protein kinase activity phosphorylating H1 is not directly involved in the acrosome reaction. The group II mAbs (and WGA), although they bind to cells as avidly as group I mAbs, neither activate adenylate cyclase nor induce H1 phosphorylation; they provide the proof that only certain domains of the 210-kD protein mediate signal transduction across the cell membrane. Additional understanding of the inductive and noninductive domains of the 210-kD protein will come from molecular cloning.

A separate topic of discussion is the possible biological significance of the phosphorylation of H1 by the endogenous kinase on only one seryl residue in a single peptide (Porter et al., 1988a). Sea urchin sperm-specific histone H1 is a unique type of H1 possessing an extended NH<sub>2</sub>-terminal tail with an unusual amino acid sequence (reviewed in Poccia, 1987). Within 10 min after fertilization, sperm H1 becomes hyperphosphorylated (~10 phosphates/mol, Green and Poccia, 1985; Porter and Vacquier, 1988; Porter et al., 1988b), and lost from the sperm pronuclear chromatin, having been replaced by cleavage stage H1 present in the egg cytoplasm (Green and Poccia, 1985; Poccia, 1987). In vitro labeling of H1 in detergent permeabilized sperm shows that the endogenous kinase is completely cAMP dependent. Reactions with purified sperm H1 and rabbit muscle cAMP-dependent protein kinase show a maximum stoichiometry of 1 mol phosphate incorporated per mol H1 (Porter et al., 1988a). We can conclude from these data that there is only one site for phosphorylation by cAMP-dependent protein kinase in native sea urchin sperm H1. It is possible that this single phosphorylation event of H1, occurring by action of the FSG-induced kinase, may cause a change in chromatin structure that could expose other sites on H1 to be phosphorylated by cAMP-independent kinases of the egg (Porter et al., 1988b), which would result ultimately in loss of sperm H1 from the male pronucleus (Green and Poccia, 1985). Regardless of the rea-

son for its existence, this is the only system in which the phosphorylation of H1 *in vivo* occurs on a single defined site. Whether it has structural consequences to the chromatin remains to be determined.

Regarding the sperm histone kinase, it could be located outside the nucleus and then move through the nuclear envelope after activation, or it could be resident in the nucleus being packaged with the highly condensed chromatin. One surprising fact about its action is that it is so specific for H1. One would assume that of the many score of different proteins in the nucleus, the preferred sequence (Kemp et al., 1977) for phosphorylation of R/K-X-S (in which X is any amino acid) would be present in many other proteins that would label if the kinase were freely diffusing in the nucleus. The H1 kinase may be resident in the nucleus, tightly associated with H1, and thus inaccessible to other substrates. Regardless of the above speculation, this system is of value in studying how signals at the cell surface are transduced across the cell membrane and through the nuclear envelope to be ultimately expressed by the action of a protein kinase on a protein involved in chromatin structure. As recently discussed (Rauscher et al., 1988), little is yet known about this fundamental process that is central to our understanding of cellular activation.

We thank Dr. Ian S. Trowbridge for help in production of mAbs and for valuable discussion, Dr. Gregory S. Kopf for advice on the adenylate cyclase assays, and Dr. David Sterns for carefully reading the manuscript.

This work was supported by National Institutes of Health grant HD-12986 to V. D. Vacquier.

Received for publication 15 June 1988, and in revised form 25 August 1988.

## References

- Bentley, J. K., D. L. Garbers, S. E. Domino, T. D. Noland, and C. Van Dop. 1986. Spermatozoa contain a guanine nucleotide-binding protein ADP-ribosylated by pertussis toxin. *Biochem. Biophys. Res. Commun.* 138:728-734.
- Bentley, J. K., A. S. Khatra, and D. L. Garbers. 1987. Receptor-mediated phosphorylation of spermatozoan proteins. *J. Biol. Chem.* 262:15708-15713.
- Christen, R., R. W. Schackmann, and B. M. Shapiro. 1982. Elevation of intracellular pH activates sperm respiration and motility of sperm of the sea urchin *Strongylocentrotus purpuratus*. *J. Biol. Chem.* 257:14881-14890.
- Cross, N. L. 1983. Isolation and electrophoretic characterization of the plasma membrane of sea urchin sperm. *J. Cell Sci.* 59:13-25.
- De Angelis, P. L., and C. G. Glabe. 1987. Polysaccharide structural features that are critical for the binding of sulfated fucans to bindin, the adhesive protein from sea urchin sperm. *J. Biol. Chem.* 262:13946-13952.
- Domino, S. E., and D. L. Garbers. 1988. The fucose sulfate glycoconjugate that induces the acrosome reaction in spermatozoa stimulates inositol 1,4,5-trisphosphate accumulation. *J. Biol. Chem.* 263:690-695.
- Garbers, D. L., and G. S. Kopf. 1980. The regulation of spermatozoa by calcium and cyclic nucleotides. *Adv. Cyclic Nucleotide Res.* 13:251-306.
- Garbers, D. L., G. S. Kopf, D. J. Tubb, and G. Olson. 1983. Elevation of sperm adenosine 3',5'-monophosphate concentrations by a fucose sulfate rich complex associated with eggs. *Biol. Reprod.* 29:1211-1220.
- Garbers, D. L., H. Shimomura, C. S. Ramarao, L. J. Dangott, and J. K. Bentley. 1987. The mechanism of action of egg peptides on spermatozoa. In *New Horizons in Sperm Cell Research*. H. Mohri, editor. Japan Scientific Societies Press, Tokyo, Japan. 159-168.
- Garvey, J. S., N. E. Cremer, and D. H. Sussdorf. 1977. *Methods in Immunology*. 3rd ed. W. A. Benjamin, Inc., London. 218-219.
- Green, G. R., and D. L. Poccia. 1985. Phosphorylation of sea urchin sperm H1 and H2B histones precedes chromatin decondensation and H1 exchange during pronuclear formation. *Dev. Biol.* 108:235-245.

- Kemp, B. E., D. J. Graves, E. Benjamin, and E. G. Krebs. 1977. Role of multiple basic residues in determining the substrate specificity of cyclic AMP-dependent protein kinase. *J. Biol. Chem.* 252:4888-4894.
- Kopf, G. S., M. J. Woolkalis, and G. L. Gerton. 1986. Evidence for a guanine nucleotide-binding regulatory protein in invertebrate and mammalian sperm. *J. Biol. Chem.* 261:7327-7331.
- Lee, H. C. 1988. Internal GTP stimulates the speract receptor mediated voltage changes in sea urchin spermatozoa membrane vesicles. *Dev. Biol.* 126:91-97.
- Lee, H. C., C. Johnson, and D. Epel. 1983. Changes in internal pH associated with initiation of motility and acrosome reaction of sea urchin sperm. *Dev. Biol.* 95:31-45.
- Laemmli, U. K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature (Lond.)*. 227:680-685.
- Lopo, A. C. 1983. Sperm-egg interactions in invertebrates. In *Mechanism and Control of Animal Fertilization*. J. F. Hartman, editor. Academic Press, Inc., New York. 269-324.
- Poccia, D. L. 1987. Regulation of chromatin condensation and decondensation in sea urchin pronuclei. In *Molecular Regulation of Nuclear Events in Mitosis and Meiosis*. R. A. Schlegel, M. S. Halleck and P. N. Rao, editors. Academic Press, Inc., New York. 149-177.
- Podell, S. B., and V. D. Vacquier. 1984. Wheat germ agglutinin blocks the acrosome reaction in *Strongylocentrotus purpuratus* sperm by binding a 210,000-mol-wt membrane protein. *J. Cell Biol.* 99:1598-1605.
- Porter, D. C., and V. D. Vacquier. 1986. Phosphorylation of sperm histone H1 is induced by the egg jelly layer in the sea urchin *Strongylocentrotus purpuratus*. *Dev. Biol.* 116:203-212.
- Porter, D. C., and V. D. Vacquier. 1988. Extraction of phosphorylated sperm specific histone H1 from sea urchin eggs: analysis of phosphopeptide maps. *Biochem. Biophys. Res. Commun.* 151:1200-1204.
- Porter, D. C., G. W. Moy, and V. D. Vacquier. 1988a. cAMP dependent protein kinase of sea urchin sperm phosphorylates histone H1 on a single site. *J. Biol. Chem.* 263:2750-2755.
- Porter, D. C., G. W. Moy, and V. D. Vacquier. 1988b. Amino terminal sequence of sea urchin sperm histone H1 and its phosphorylation by egg cytosol. *Comp. Biochem. Physiol.* In press.
- Rauscher, F. J. III, D. R. Cohen, T. Curren, T. J. Bos, P. K. Vogt, D. Bohmann, R. Tjian, and B. R. Franza, Jr. 1988. Fos-associated protein p39 is the product of the jun proto-oncogene. *Science (Wash. DC)*. 240:1010-1016.
- Schackmann, R. W. 1986. Ion measurements in sea urchin sperm. *Methods Cell Biol.* 27:57-70.
- Schackmann, R. W. 1988. The acrosome reaction. In *Fertilization*. Vol. 1. H. Schatten and G. Schatten, editors. Academic Press, Inc., New York. In press.
- Schackmann, R. W., and P. B. Chock. 1986. Alteration of intracellular  $[Ca^{2+}]$  in sea urchin sperm by the egg peptide speract. *J. Biol. Chem.* 261:8719-8728.
- SeGall, G. K., and W. J. Lennarz. 1979. Chemical characterization of the component of the jelly coat from sea urchin eggs responsible for induction of the acrosome reaction. *Dev. Biol.* 71:33-48.
- SeGall, G. W., and W. J. Lennarz. 1981. Jelly coat induction of the acrosome reaction in echinoid sperm. *Dev. Biol.* 86:87-93.
- Schultz, G., and K. H. Jakobs. 1983. Adenylate cyclase. In *Methods of Enzymatic Analysis*. Vol. 4. U. Bergmeyer, editor. Springer-Verlag, Berlin, FRG. 369-378.
- Tilney, L. G. 1985. The acrosomal reaction. In *Biology of Fertilization*. C. B. Metz and A. Monroy, editors. Academic Press, Inc., Orlando, FL. 157-213.
- Trimmer, J. S. 1987. Sea urchin antigens mediating the acrosome reaction. Ph.D. dissertation. University of California, San Diego, San Diego, CA. 273 pp.
- Trimmer, J. S., and V. D. Vacquier. 1986. Activation of sea urchin gametes. *Annu. Rev. Cell Biol.* 2:1-26.
- Trimmer, J. S., and V. D. Vacquier. 1988. Monoclonal antibodies induce the translocation, patching and shedding of surface antigens of sea urchin spermatozoa. *Exp. Cell Res.* 175:37-51.
- Trimmer, J. S., I. S. Trowbridge, and V. D. Vacquier. 1985. Monoclonal antibody to a membrane glycoprotein inhibits the acrosome reaction and associated  $Ca^{2+}$  and  $H^+$  fluxes of sea urchin sperm. *Cell*. 40:697-703.
- Trimmer, J. S., R. W. Schackmann, and V. D. Vacquier. 1986. Monoclonal antibodies increase intracellular  $Ca^{2+}$  in sea urchin spermatozoa. *Proc. Natl. Acad. Sci. USA*. 83:9055-9059.
- Trimmer, J. S., Y. Ebina, R. W. Schackmann, C. G. Meinhof, and V. D. Vacquier. 1987. Characterization of a monoclonal antibody that induces the acrosome reaction of sea urchin sperm. *J. Cell Biol.* 105:1121-1128.
- Vacquier, V. D. 1987. Plasma membranes isolated from sea urchin spermatozoa. In *New Horizons in Sperm Cell Research*. H. Mohri, editor. Japan Science Societies Press, Tokyo, Japan. 217-233.