

SI Table 1.Comparison of properties of T-type Ca^{2+} currents

Reference	Tissue	$[\text{Ca}^{2+}]_{\text{ext}}$, mM	IV_{peak} , mV	$V_{0.5\text{act}}$, mV	$V_{0.5\text{inact}}$, mV	$\tau_{\text{deact}\,1,2}$, ms	$\tau_{\text{decay}\,1,2}$, ms	τ_{recovery} , ms	Nickel-sensitive
This study	Chicken hair cells	5	-15	-38	-37	0.35 24	35 330	40 1,200	Yes
1	Olfactory	3	-40	-45	-66	-	-	-	Yes
2	Cone bipolar	10	-30	-36	-59	-	30	-	Yes
3	Myocytes	2	-30	-42	-68	1.3	13	-	Yes
4	Skeletal muscle	10	-15	-24	-41	-	16	224 5,160	Yes
5	Cochlear nucleus	2	-40	-59	-89	-	-	-	Yes

1. Kawai, F. & Miyachi, E. (2001) *J Neurosci* 21, RC144.
2. Pan, Z. H. (2000) *J Neurophysiol* 83, 513-27.
3. Leuranguer, V., Monteil, A., Bourinet, E., Dayanithi, G. & Nargeot, J. (2000) *Am J Physiol Heart Circ Physiol* 279, H2540-8.
4. Berthier, C., Monteil, A., Lory, P. & Strube, C. (2002) *J Physiol* 539, 681-91.
5. Doughty, J. M., Barnes-Davies, M., Rusznak, Z., Harasztsosi, C. & Forsythe, I. D. (1998) *J Physiol* 512 (Pt 2), 365-76.