

Transcriptional regulation of the human *c-myc* gene

J.C. Lang¹, B. Whitelaw², S. Talbot¹ & N.M. Wilkie¹

¹Beatson Institute for Cancer Research, Garscube Estate, Switchback Road, Bearsden, Glasgow G61 1BD and ²I.A.P.G.R., King's Buildings, West Mains Road, Edinburgh EH9 3JQ, UK.

Summary The involvement of *c-myc* in the genesis of animal neoplasia is now well documented for several systems. In order to define the precise role played by the *myc* gene in tumorigenesis, a better understanding of the normal regulation of *myc* expression is necessary. We have begun a study of the *cis*-acting regulatory sequences within the 5' flanking domain of the human *c-myc* gene. Regions important for *myc* promoter function have been identified by linkage to the coding sequences of the bacterial chloramphenicol acetyltransferase (*cat*) gene. Promoter deletion studies and *in vivo* competition assays for *c-myc/cat* recombinant plasmids have allowed the identification of a proximal 'core' promoter region capable of directing high levels of CAT activity. Further upstream a negative regulatory element (NRE2) has been identified which is capable of repressing *cat* gene expression and which functions by interaction with a trans-acting factor(s). Preliminary data suggests detection of NRE2 is dependent on both the type and amount of carrier DNA used in transient CAT assays. Initial experiments further indicate the involvement of at least two other distal regulatory domains, a negative regulatory domain (NRE1) and a putative enhancer-type region (E). *In vitro* footprint analysis has allowed the identification of DNA binding proteins which interact with NRE2 and the 'core' promoter. NRE2 contains binding sites for transcription factors Sp1 and CTF. The 'core' promoter domain appears to be highly complex and possesses several Sp1 binding sites.

The *myc* gene was first described as the oncogene of certain highly transforming avian retroviruses (reviewed by Hayman, 1983). Since then a substantial body of evidence has accumulated suggesting that the *myc* gene is intimately associated with the control of cellular proliferation and differentiation. Thus experimental and spontaneous tumours in rodents and cats have been observed which contain defective retroviruses expressing transduced *v-myc* genes while a variety of avian, rodent and feline tumours express *c-myc* genes activated by proviral DNA insertion (Hayward *et al.*, 1981; Corcoran *et al.*, 1984; Neil *et al.*, 1984). The *c-myc* locus is also interrupted by translocation in human Burkitt's lymphoma (BL) and murine plasmacytoma (MPC), (reviewed by Marcu, 1987) and by amplification both in some promyelocytic leukaemia cell lines and in some cases of human breast cancer (reviewed by Alitalo, 1985).

Further evidence for effects of *myc* on growth are provided by experiments which show that when the gene, linked to strong transcriptional enhancers, is transfected into cells in culture perturbations in differentiation patterns are observed (Coppola & Cole, 1986). Similarly, the *c-myc* gene is capable of inducing multiple neoplasms in transgenic mice when fused to immunoglobulin enhancers (Adams *et al.*, 1985) or a mouse mammary tumour virus LTR (Leder *et al.*, 1986).

Unlike the *ras* gene family, mutations within the coding sequences may not be an important feature in converting *myc* from a proto-oncogene to an oncogene. Thus feline *v-myc* coding sequences are virtually unaltered compared to the feline *c-myc* (Stewart *et al.*, 1986). The bulk of the present evidence suggests that aberrant expression of *c-myc* is the principal mechanism for oncogenic conversion.

The current view of the structure of the *c-myc* gene is shown in Figure 1. Expression may be regulated under 'normal' circumstances at at least three levels. Firstly, there is regulation at the level of initiation of transcription, affected by *cis*-acting regulatory domains 5' to the two main cap sites (P1 and P2) and perhaps by other domains within the body of the gene (reviewed by Marcu, 1987). During terminal differentiation of B cells, red cells and monocytes *c-myc* expression is switched off, and in the case of HL60 promyelocytic cells this partly occurs as a result of reduced initiation of transcription (Siebenlist *et al.*, 1988). Secondly, in quiescent murine fibroblasts, non-proliferating HL60 cells,




Figure 1 Structure of the human *c-myc* gene, exons 1-3 are represented by an open box and the coding regions of proposed *c-myc* proteins MYC 1 and MYC 2 (Hann *et al.*, 1988) by a solid box. Previously described *in vivo* hypersensitive sites I-III₂ (Siebenlist *et al.*, 1984) are indicated. The position of relevant restriction enzyme sites are shown relative to *c-myc* RNA cap site P1.

and in human mononuclear cells (MNC) the absence of full length *c-myc* transcripts is associated with an apparent block in elongation of already initiated RNA, located at a point close to the junction of exon 1 and the first intron (Bentley & Groudine, 1986b; Eick & Bornkamm, 1986; Nepveu *et al.*, 1987). This block is overcome on mitogen stimulation of the cell cycle (Eick *et al.*, 1987; Nepveu *et al.*, 1987). Thirdly, *c-myc* RNA is known to have a short half life which is apparently lengthened in some murine plasmacytomas and during differentiation of F9 teratocarcinoma stem cells (Piechaczyk *et al.*, 1985; Dean *et al.*, 1986).

It is not yet fully understood how the changes in the *c-myc* locus observed in tumour cells affect regulation. The translocations observed in BL and MPC often, but not always, involve breakpoints within the transcriptional control region of *c-myc*. This might disrupt normal regulatory sequences (Yang *et al.*, 1985). Alternatively, transcriptional regulatory sequences within the immunoglobulin locus translocated to the *c-myc* gene may exert a dominant effect on *c-myc* as may retroviral LTRs found adjacent to *c-myc* in lymphomas of chickens, mice and cats (Hayward *et al.*, 1981; Payne *et al.*, 1984; Corcoran *et al.*, 1984; Neil *et al.*, 1984; O'Donnell *et al.*, 1985). In many cases of BL, however, aberrant *myc* expression may be due to mutation of sequences located at the 3' end of exon 1 thought to be involved in an elongation block to transcription (Cesarman *et al.*, 1987). Alternatively, mutations located at this site might affect the expression of the proposed MYC 1 protein (Hann *et al.*, 1988). The mechanisms of alteration in cases of amplification of *c-myc*,

and in those cases where no rearrangement in the *myc* locus is apparent, remain to be investigated.

It seems obvious that greater understanding of the normal regulation of *myc* expression will be necessary for analysis of the alterations which occur during oncogenic conversion. In this paper we summarise our own studies of *cis*-acting regulatory sequences within the 5' flanking domain of the human *c-myc* gene. Such sequences were identified by linkage to the bacterial chloramphenicol acetyltransferase gene (*cat*), and in some cases further characterised by analysis of DNA binding proteins using *in vitro* DNAase I footprint analysis.

Identification of transcriptionally active domains within the 5' flanking sequence of *c-myc*

We linked the 5' flanking sequence and most of the first exon sequence of *c-myc* (nucleotides -2319 to +511, Figure 1) to the bacterial *cat* gene. This interrupts the *myc* sequences at which an elongation block to transcription has been suggested and also removes all known coding sequences for MYC1 and 2 (see Figure 1). The *cat* gene is then fused to untranslated *myc* leader sequences and placed under transcriptional control of the *myc* promoter. A series of 5' terminal deletions and internal deletions were also constructed. Following transfection into murine fibroblasts (LTK⁻) and human epithelial cells (HeLa), relative levels of CAT activity were measured using transient assays. In other experiments, various domains of the 5' flanking sequence were transferred to *cat* under control of heterologous promoters and CAT activity similarly measured. This approach permitted identification of *cis*-acting regulatory domains affecting the expression of *cat*.

The results have been reported in detail elsewhere (Whitelaw *et al.*, 1988) but are summarised in Figure 2. Evidence was obtained for a positive regulatory element, perhaps an enhancer, located between -2319 and -1980 relative to P1. From Figure 2 it can be seen that this domain does not encompass *in vivo* DNAase hypersensitive site I. Closer to the cap site, a *cis*-acting negative regulatory element (NRE1) encompassing hypersensitive site II₁ was identified, which mapped between -1527 to -1246. Between -1052 and -607 another strong negative element, NRE2 encompasses hypersensitive site II₂. Much closer to the cap site a positive acting region was found which is in the domain containing hypersensitive sites III₁ and III₂, and we presume this corresponds to the 'core' promoter.

Cis-acting negative regulatory sequences have also been reported for the human and murine *c-myc* genes by others (Chung *et al.*, 1986; Remmers *et al.*, 1986; Hay *et al.*, 1986; Lipp *et al.*, 1987). Comparison of the results reported from various laboratories suggests that the situation is complex. Remmers *et al.* and Lipp *et al.* find negative regulatory domains located between -1188 and -428 and -1052 and -511 respectively. These results are consistent with our own which locate NRE2 to within a 445 bp sequence (-1052 to -607). In contrast, Chung *et al.* and Hay *et al.* describe




Figure 2 A model for *cis*-acting transcriptional control domains within the human *c-myc* regulatory region. Transcriptional regulatory domains defined by *in vivo* CAT assays and *in vitro* DNase I footprint studies are indicated. Solid circle represents Sp1 binding sites, open circle Sp1-like binding site and solid oval CTF binding sites. The TATAA boxes (T) are indicated.

negative elements between -607 to -407 and -353 to 293 respectively, outside the map location of NRE2. Chung *et al.* (1986) also describe two additional negative regions located in the first exon and the first intron, domains which we have not yet examined. It is suggested that the latter element may be involved in the elongation block proposed by Bentley and Groudine (1986b).

It is clear that while various laboratories have detected negative regulatory elements, their reported locations (and the strength of their effects) appear to vary. The reasons for this are not yet known but one factor may be the different cell types and the different conditions used in the transient assays which most laboratories utilise.

Preliminary experiments from our laboratory indicate that one such variable may be the type and amount of carrier DNA used. We measured transient CAT expression from equimolar amounts of *cat/myc* recombinants containing the entire *myc* 5' regulatory region or with a terminal deletion to position -290, which removes hypersensitive sites I, II₁, and II₂ and also E, NRE1 and NRE2. Mouse embryo, salmon sperm or pUC18 plasmid DNA was used as carrier in varying amounts. The results are shown in Figure 3 and reveal two important effects. Firstly, no significant difference in CAT activity between the full length and deleted plasmids is obtained at carrier DNA concentrations of less than 10 μ g per plate. Secondly, the deleted plasmid directs substantially more CAT activity than the full length construction at mouse embryo and salmon sperm carrier DNA concentrations of between 10 and 60 μ g per plate. In contrast, using pUC18 DNA as carrier, no clear difference in CAT levels between the full length and deleted plasmids was observed at all carrier DNA concentrations measured. However, these effects should not be confused with the general decrease in CAT levels observed as the amount of carrier DNA increases. This additional factor is probably due to reduced efficiency of DNA transfection at higher carrier DNA concentrations and can equally be observed with recombinants containing the *cat* gene linked to other non-*myc* promoter sequences (data not shown).

We tentatively propose the following explanation for these effects. The 5' flanking sequence of *myc* contains at least four detectable *cis*-acting regulatory elements, an upstream positive element (E), two negative elements (NRE1 and NRE2) and a cap-proximal 'core' promoter (see Figure 2), which will all interact with trans-acting factors in the cell. Eukaryote DNA may contain specific sequences which act as targets for some of these trans-acting factors, and can titrate these factors if supplied at sufficiently high concentrations. There is evidence that this can occur in several systems and also in the *myc* regulatory sequence (see following section). At low carrier DNA concentrations the amount of target sequence may be too low to titrate trans-acting factors, and the effect of E balances the effects of NREs in LTK⁻ cells. As the concentration of carrier increases, specific sequences in the DNA compete with the upstream E sequences for trans-acting factors which modulate its activity, thus unveiling the negative effects of the NRE sequences which can then be detected by the terminal deletion. pUC18 plasmid DNA is unlikely to have specific target sites for eukaryote transacting factors (except coincidentally) and might be expected to have no significant effect on CAT levels.

Further characterisation of NRE2

The magnitude of the negative effect of plasmids containing NRE1 and NRE2 vary to some extent from experiment to experiment (Whitelaw *et al.*, 1988). We considered the possibility that the *cis*-acting NREs interact with cellular trans-acting factors present in limiting amounts. Increasing concentrations of DNA containing *myc* promoter sequences might then titrate out such factors leading to derepression of unbound DNA molecules which remain.




Figure 3 *Cat* activity of human *c-myc/cat* recombinant plasmids in mouse $LATK^-$ cells. Recombinants encompassing the entire *c-myc* promoter domain (pB12) or the truncated promoter (position -290 ; p240) linked to *cat* are shown. Graphs show levels of acetylated ^{14}C chloramphenicol/ 5×10^5 cells for $5 \mu g$ of recombinant plasmid pB12 and the molar equivalent for p240 after transfection into separate cultures using increasing amounts of carrier DNA. Each point represents the average of 2 values for each experiment. Protocols for DNA transfections and CAT assays have been described previously (Whitelaw *et al.*, 1988).

In order to test this we performed plasmid competition assays in which the 5' sequences containing only hypersensitive sites I, II₁ and II₂ (-2319 to -350 ; Figure 2) are used as competitor in CAT assays where *cat* is driven by the entire *myc* promoter region (plasmid pB12, Figure 3). As a control the level of CAT activity obtained with the truncated promoter (position -350) in which *cat* is driven by only the core promoter containing hypersensitive sites III₁ and III₂ (see Figures 1 & 2) was also measured. The results presented in Figure 4 show that the competitor DNA can titrate out factors responsible for repression of expression from pB12




Figure 4 Competition assay for *c-myc/cat* recombinants. *c-myc* sequences from -2319 to -350 alone were used in a competition assay against pB12 transfected into the same cells at molar plasmid ratios of 1, 2, 5 and 10 competitor to parental plasmid pB12. Results represent the CAT activity of $1 \mu g$ of a promoterless *cat* construct (dotted line) and molar equivalents of pB12 and pB14 (truncated deletion to position -350), expressed as a percentage of pB14. Each point represents the average of 2 flasks for two separate experiments A and B.

and that levels of expression equal to or greater than that obtained with *cat* driven by the truncated promoter can be obtained.

Transcription factor binding sites in the regulatory domains

It is likely that the trans-acting factors which can be titrated in the competition experiments are DNA-binding proteins which affect the activity of the *cis*-acting regulatory DNA sequences. These might be proteins which recognise and bind to specific sequences, or proteins which interact through protein-protein interactions. In order to begin analysis of the regulatory domain, we examined the interaction of specific DNA-binding proteins with NRE2 and the 'core' promoter using partially and highly purified preparations of nuclear proteins in *in vivo* DNase I protection assays (Jones *et al.*, 1985). The results have been presented in detail elsewhere (Whitelaw *et al.*, 1988) but the footprints obtained using partially purified nuclear extracts binding to NRE2 are shown in Figure 2. Four distinct 'footprints' are obtained. Using DNA-affinity purified preparations of transcription factors Sp1 and CTF (CAAT Transcription Factor) (Kadonaga & Tjian, 1985), it was possible to show that of these one binding site was due to Sp1 and two adjacent binding sites to CTF. Examination of the DNA sequence confirms typical consensus signals for these DNA-binding proteins. The fourth footprint encompasses the reported location of hypersensitive site II₂ (Siebenlist *et al.*, 1984) and although it contains an Sp1-like sequence it bound neither purified Sp1 nor CTF. This binding site may represent a target for a so far unknown DNA-binding protein.

A summary of this study and a similar analysis of the 'core' promoter domain is included in Figure 2. This shows the location and orientation of the DNA binding sites relative to the other known features of the regulatory region. We have not yet analysed NRE1 or E, or the NRE domains suggested by the reports of Chung *et al.* (1986) or Hay *et al.* (1987).

Discussion

Cumulative experimental evidence to date suggests that transcriptional control of the *c-myc* oncogene is particularly complex. Despite efforts from a number of research groups, the domains responsible for transcriptional regulation have yet to be fully characterised. The emerging picture is one of intricate interactions between multiple positive and negative control elements. *C-myc* is expressed in many different cell types. It appears to encode a protein(s) intimately involved in the regulation of cell growth and differentiation and as such may itself require to be regulated very precisely. It is plausible that varying interactions occur between control domains and trans-acting factors depending on the cell type and stage of differentiation. That negative regulatory elements identified by transient assays have *in vivo* significance is suggested by the observations that such an element detected in murine *myc* (which may correspond to our NRE2) correlates with the position of chromosomal break-points and proviral insertion sites in B- and T-cell neoplasia (Remmers *et al.*, 1986). Similarly mutations in the 3' region of exon 1 in BL cells correlates with aberrant patterns of *myc* transcription and may affect elongation block, or perhaps the expression of MYC1 (Cesarman *et al.*, 1987; Hann *et al.*, 1988). Further efforts to identify the *in vivo* regulatory function of these and the other elements described to date are obviously crucial. It is also important to determine whether alterations in the DNA sequence of the elements themselves, or alteration in the trans-acting factors which regulate their activity, are involved in the alteration of *c-myc* expression associated with various types of cancer apart from BL.

From the summary shown in Figure 2, we suggest a role for specific DNA-binding proteins Sp1 and CTF in

modulating the activity of NRE2. Binding of these general transcription factors have been associated so far with up-regulation of promoter elements. If they are also involved in negative regulation, the specificity of their effect on transcription remains to be explained. Specificity could be determined by the structure of the local DNA sequence itself, by the orientation or spacing of binding or the nature of other specific DNA binding proteins which interact with the domain. Alternatively, specificity could be determined by protein-protein interactions with other non-DNA binding proteins, or a combination of all of these. Regulation of *cis*-acting DNA sequences in transcription by the interaction of accessory proteins with specific protein-DNA complexes has been reported for other biological systems including the interaction of herpes simplex virus with host cellular proteins (Preston *et al.*, 1988). Another possible mode of action of a negative element might be to act as a promoter and lead to 'promoter occlusion effects' on downstream promoters as previously suggested for bacteriophage lambda gene expression (Adhya & Gottesman, 1982). In this regard, the presence of an additional cap site (PO) has been reported adjacent to NRE2 (Bentley & Groudine, 1986a).

It is obvious that the mechanisms by which expression of the *myc* gene is regulated is both complex and little understood at the present time. We are currently investigating the role of Sp1, CTF and the unknown protein which binds at hypersensitive site II₂, and the possible role of accessory proteins, using a combination of site directed mutagenesis, DNase I protection studies, and analysis of protein-DNA complexes.

We thank the Cancer Research Campaign of Great Britain for support.

References

- ADAMS, J.M., HARRIS, A.W., PINKERT, C.A. & 5 others (1985). The *c-myc* oncogene driven by immunoglobulin enhancers induces lymphoid malignancy in transgenic mice. *Nature*, **318**, 533.
- ALITALO, K. (1985). Amplification of cellular oncogenes in cancer cells. *TIBS*, **10**, 194.
- ADHYA, S. & GOTTESMAN, M. (1982). Promoter occlusion: Transcription through a promoter may inhibit its activity. *Cell*, **29**, 939.
- BENTLEY, D.L. & GROUDINE, M. (1986a). Novel promoter upstream of the human *c-myc* gene and regulation of *c-myc* expression in B-cell lymphomas. *Mol. Cell. Biol.*, **6**, 3481.
- BENTLEY, D.L. & GROUDINE, M. (1986b). A block to elongation is largely responsible for decreased transcription of *c-myc* in differentiated HL60 cells. *Nature*, **321**, 702.
- CESARMAN, E., DALLA-FAVERA, R., BENTLEY, G. & GROUDINE, M. (1987). Mutations in the first exon are associated with altered transcription of *c-myc* in Burkitt lymphoma. *Science*, **238**, 1272.
- CHUNG, J., SINN, E., REED, R.R. & LEDER, P. (1986). Transacting elements modulate expression of the human *c-myc* gene in Burkitt lymphoma cells. *Proc. Natl Acad. Sci. USA*, **83**, 7918.
- COPPOLA, J.A. & COLE, M.D. (1986). Constitutive *c-myc* oncogene expression blocks mouse erythroleukaemia cell differentiation but not commitment. *Nature*, **320**, 760.
- CORCORAN, L.M., ADAMS, J.M., DUNN, A.R. & CORY, S. (1984). Murine T lymphomas in which the cellular *myc* oncogene has been activated by retroviral insertion. *Cell*, **37**, 113.
- DEAN, M., LEVINE, R.A. & CAMPISI, J. (1986). *C-myc* regulation during retinoic acid-induced differentiation of F9 cells is post-transcriptional and associated with growth arrest. *Mol. Cell. Biol.*, **6**, 518.
- EICK, D. & BORNKAMM, G.W. (1986). Transcriptional arrest within the first exon is a fast control mechanism in *c-myc* gene expression. *Nucleic Acids Res.*, **14**, 8331.
- EICK, D., BERGER, R., POLACK, A. & BORNKAMM, G.W. (1987). Transcription of *c-myc* in human mononuclear cells is regulated by an elongation block. *Oncogene*, **2**, 61.
- HANN, S.R., KING, M.W., BENTLEY, D.L., ANDERSON, C.W. & EISENMAN, R.N. (1988). A non-AUG translational initiation in *c-myc* exon 1 generates an N-terminally distinct protein whose synthesis is disrupted in Burkitt's lymphoma. *Cell*, **52**, 185.
- HAY, N., BISHOP, J.M. & LEVENS, D. (1987). Regulatory elements that modulate expression of human *c-myc*. *Genes and Development*, **1**, 659.
- HAYMAN, M.J. (1983). Avian acute leukaemia viruses. *Curr. Top. Microbiol. Immunol.*, **98**, 109.
- HAYWARD, W.S., NEEL, B.G. & ASTRIN, S.N. (1981). Activation of a cellular *onc* gene by promoter insertion in ALV-induced lymphoid leukaemia. *Nature*, **290**, 475.
- JONES, K.A., YAMAMOTO, K.R. & TJIAN, R. (1985). Two distinct transcription factors bind to the HSV thymidine kinase promoter *in vitro*. *Cell*, **42**, 559.
- KADONAGA, J.T. & TJIAN, R. (1986). Affinity purification of sequence-specific DNA binding proteins. *Proc. Natl Acad. Sci. USA*, **83**, 5889.
- LEDER, A., PATTENGAL, P.K., KNO, A., STEWART, T.A. & LEDER, P. (1986). Consequence of widespread deregulation of the *c-myc* gene in transgenic mice: Multiple neoplasms and normal development. *Cell*, **45**, 485.
- LIPP, M., SCHILLING, R., WIEST, S., LAUX, G. & BORNKAMM, G.W. (1987). Target sequences for *cis*-acting regulation within the dual promoter of the human *c-myc* gene. *Mol. Cell. Biol.*, **7**, 1393.
- MARCU, K.B. (1987). Regulation of expression of the *c-myc* proto-oncogene. *BioEssays*, **6**, 28.
- NEIL, J.C., HUGHES, D., McFARLANE, R & 4 others (1984). Transduction and rearrangement of the *myc* gene by feline leukaemia virus in naturally occurring T-cell leukaemias. *Nature*, **308**, 814.
- NEPVEU, A., LEVINE, R.A., CAMPISI, J., GREENBERG, M.E., ZIFF, E.B. & MARCU, K.B. (1987). Alternative modes of *c-myc* regulation in growth factor-stimulated and differentiating cells. *Oncogene*, **1**, 243.
- O'DONNELL, P.V., FLEISSENER, E., LONIAL, H., KOEHNE, C.F. & REICIN, A. (1985). Early clonality and high frequency proviral integration into the *c-myc* locus in AKR leukaemias. *J. Virol.*, **55**, 500.
- PAYNE, G.S., BISHOP, J.M. & VARMUS, H.E. (1982). Multiple arrangements of viral DNA and an activated host oncogene in bursal lymphomas. *Nature*, **295**, 209.

- PIECHACZYK, M., YANG, J.-Q., BLANCHARD, J.-M., JEANTEUR, P. & MARCU, K.B. (1985). Post-transcriptional mechanisms are responsible for accumulation of truncated *c-myc* RNAs in murine plasma cell tumors. *Cell*, **42**, 589.
- PRESTON, C.M., FRAME, M.C. & CAMPBELL, M.E.M. (1988). A complex formed between cell components and HSV structural polypeptide binds to a viral immediate early gene regulatory DNA sequence. *Cell*, **52**, 425.
- REMMERS, E.F., YANG, J.-Q. & MARCU, K.B. (1986). A negative transcriptional control element located upstream of the murine *c-myc* gene. *EMBO J.*, **5**, 899.
- SIEBENLIST, U., HENNINGHAUSEN, L., BATTERY, J. & LEDER, P. (1984). Chromatin structure and protein binding in the putative regulatory region of the *c-myc* gene in Burkitt lymphoma. *Cell*, **37**, 381.
- SIEBENLIST, U., BRESSLER, P. & KELLY, K. (1988). Two distinct mechanisms of transcriptional control operate on *c-myc* during differentiation of HL60 cells. *Mol. Cell. Biol.*, **8**, 867.
- STEWART, M.A., FORREST, D., McFARLANE, R., ONIONS, D., WILKIE, N. & NEIL, J.C. (1986). Conservation of the *c-myc* coding sequence in transduced feline *v-myc* genes. *Virology*, **154**, 121.
- TAVASSOLI, M. & SHALL, S. (1988). Transcription of the *c-myc* oncogene is altered in spontaneously immortalized rodent fibroblasts. *Oncogene*, **2**, 337.
- WHITELAW, B., WILKIE, N.M., JONES, K.A., KADONAGA, J.T., TJIAN, R. & LANG, J.C. (1988). Transcriptionally active domains in the 5' flanking sequence of human *c-myc*. *UCLA Symposia on Molecular and Cellular Biology*, **58**, 337.
- YANG, J.-Q., BAUER, S.R., MUSHINSKI, J.F. & MARCU, K.B. (1985). Chromosome translocations clustered 5' of the murine *c-myc* gene qualitatively affect promoter usage: Implications for the site of normal *c-myc* regulation. *EMBO J.*, **4**, 1441.