			Frequency and percentage of the observations with value higher than median within each group (Binary PSI: bad performance=1)					
	Total		All hospitals		Rural PPS		САН	
PSIs	observations	Median	Frequency	Percentage	Frequency	Percentage	Frequency	Percentage
PSI-2	630	0.00	62	9.84%	49	10.47%	13	8.02%
PSI-3	624	7.47	312	50.00%	243	51.48%	69	45.39%
PSI-5	712	0.00	25	3.51%	20	3.98%	5	2.39%
PSI-6	712	0.00	92	12.92%	82	16.30%**	· 10	4.78%**
PSI-7	712	0.00	136	19.10%	119	23.66%**	· 17	8.13%**
PSI-15	712	0.00	323	45.37%	272	54.08%**	51	24.40%**
Composite score of 4PSIs (weighted by numerator)	712	0.42	356	50.00%	293	58.25%**	63	30.14%**
Composite score of 6PSIs (weighted by numerator)	594	4.78	297	50.00%	239	52.64%**	58	41.43%**
Composite score of 4PSIs (weighted by denominator)	712	0.44	356	50.00%	288	57.26%**	68	32.54%**
Composite score of 6PSIs (weighted by denominator)	594	1.12	297	50.00%	242	53.30%**	55	39.29%**

Table S2. Median for Continuous PSIs, Frequency and Percentage for Binary PSIs

\*\* Percentages of bad performance between CAH and rural PPS hospitals are significantly different (Chi-square test)