

Table 9. LCMV infection does not induce severe hemorrhagic anemia in mice lacking P-selectin or GPIb α or in mice treated with Warfarin

Treatment	Mouse strain	<i>n</i>	Days p.i.	Platelets (10 ³ / μ l)	Hct (percent)	PT (INR)
NaCl	C57BL/6J	3	6	1305 \pm 72	46.7 \pm 3.2	nd
LCMV	C57BL/6J	6	6	291 \pm 52	34.4 \pm 2.9	nd
NaCl	GPIb α ^{-/-}	3	6	622 \pm 41	47.1 \pm 3.9	nd
LCMV	GPIb α ^{-/-}	6	6	298 \pm 24	36.7 \pm 3.5	nd
NaCl	P-selectin ^{-/-}	3	6	902 \pm 43	44.7 \pm 2.5	nd
LCMV	P-selectin ^{-/-}	6	6	245 \pm 30	34.3 \pm 3.7	nd
NaCl	B10.D2	6	6	1247 \pm 34	48.4 \pm 2.3	0.8
α -PLT + LCMV	B10.D2	6	6	8 \pm 2	10.2 \pm 2.9	0.8
Diluent + LCMV	B10.D2	6	6	204 \pm 63	38.3 \pm 3.8	0.8
Warfarin + LCMV	B10.D2	6	6	222 \pm 58	38.2 \pm 4.1	5.8

All measurements were performed at the indicated time post injection (p.i.) of NaCl or LCMV Arm (10⁶ PFU/mouse). Platelets and Hct (mean \pm SD) were assessed individually. Prothrombin time (PT) was assessed in blood sample pools obtained from each group; nd: not done; INR: international normalized ratio.