

**Supplemental Table S1.** Primers used for quantification of mRNAs from common bean nodules by qRT-PCR

Gene	Identifier	Forward	Reverse
<i>Ubiquitin</i>	U77940	GCTTCGTGGTGGAAATGCAGAT	TCGCACCTTGGCAGACTACAA
<i>GMP</i>	TC466	AGCTGCACAATAATGCGAGGA	TCCACTCGAGACCATTGACCA
<i>GME</i>	TC410	CTTTGGAATTGAGTGCCGCA	TCTTCCGGCAAAAATGCAGC
<i>GalPP</i>	CV538392	CTGGAATCCGTGGAAAAGGTG	TGTTCCAATCTCTGTGGCGAG
<i>GalDH</i>	TC2872	CGATTGATGTGGTGCTTTCGT	GGAGAAGCATTGATGATGGCA
<i>GalLDH</i>	--- <sup>a</sup>	GCAAAGCCAAGCATGAAAAGAC	ATTGGGCAGCGATGGACAT
<i>γECS</i>	AF128454	AGCTGTGTCCCACCGAGTG	AAGAGTGAGCGGAGGAGGGT
<i>hGSHS</i>	AF258320	AGTTCAGGTGTGGGTTTGG	GGCCCGGGTAATAAGCAAAG
<i>DR1</i>	CB539361	CAATGTCTTCCGTTCCCCCT	AGCTTTAACGGCGATTTCGA
<i>DR2</i>	TC2364	GCAACAAACCCGAATGGTTT	ACGGGCACCTTTCCTTCAG
<i>AO</i>	Pv_ERRI_26	CAGGAGTTTGGGCCTTCCAT	TTCCCATGTGCAAATGTGGT

<sup>a</sup> Primers (5' →3') were designed based on a partial sequence obtained from a bean nodule cDNA as indicated in the Materials and Methods.