

Table S1. **Protein composition and biochemical properties of polyQ- versus silica-NP-induced NIs**

	Expanded PolyQ-induced NIs	Silica-NP-induced ubiquitin NIs	Silica-NP-induced topo I NIs
PolyQ	+ ^{a,b}	+	+ ^c
TBP	+ ^{a,d,e}	+	+ ^c
CBP	+ ^{f,g}	+	+ ^c
Huntingtin	+ ^{d,e,h,i,j,k}	+	+ ^c
Ataxin-1	+ ^{g,l}	ND	ND
Ataxin-3	+ ^{m,n}	ND	+
Ubiquitin	+ ^{a,e,g,h,i,j,l,n}	+	+ ^c
20S proteasomes	+ ^{g,h,i,j,m,o}	+	+ ^c
UbcH6	ND	+	+ ^c
MDM2	+ ^{e,p}	+	+ ^c
Topo I	ND	+	+ ^c
PML	+ ^{b,p,q}	+	+ ^c
p80 coilin	+ ^q	+	+ ^c
Fibrillarin	ND	+	+ ^c
PCNA	ND	—	— ^c
Filter retardation assay	+ ^r	+	+
Congo red sensitive	+ ^s	+	+
Trehalose sensitive	+ ^t	+	+ ^c
Proteasome inhibitor sensitive	+ ^u	+	+
UPS, PML coat	+ ^q	+	+

MDM2, murine double minute 2; PCNA, proliferating cell nuclear antigen; TBP, TATA binding protein.

^aNakamura, K., S.Y. Jeong, T. Uchihara, M. Anno, K. Nagashima, T. Nagashima, S. Ikeda, S. Tsuji, and I. Kanazawa. 2001. *Hum. Mol. Genet.* 10:1441–1448.

^bJaner, A., E. Martin, M.P. Muriel, M. Latouche, H. Fujigasaki, M. Ruberg, A. Brice, Y. Trotter, and A. Sittler. 2006. *J. Cell Biol.* 174:65–76.

^cChen, M., and A. von Mikecz. 2005. *Exp. Cell Res.* 305:51–62.

^dHuang, C.C., P.W. Faber, F. Persichetti, V. Mittal, J.P. Vonsattel, M.E. MacDonald, and J.F. Gusella. 1998. *Somat. Cell Mol. Genet.* 24:217–233.

^eSuhr, S.T., M.C. Senut, J.P. Whitelegge, K.F. Faull, D.B. Cuizon, and F.H. Gage. 2001. *J. Cell Biol.* 153:283–294.

^fKazantsev, A., E. Preisinger, A. Dranovsky, D. Goldgaber, and D. Housman. 1999. *Proc. Natl. Acad. Sci. USA.* 96:11404–11409.

^gStenoien, D.L., M. Mielke, and M.A. Mancini. 2002. *Nat. Cell Biol.* 4:806–810.

^hJana, N.R., E.A. Zemskov, Gh Wang, and N. Nukina. 2001. *Hum. Mol. Genet.* 10:1049–1059.

ⁱWaelter, S., A. Boeddrich, R. Lurz, E. Scerzinger, G. Lueder, H. Lehrach, and E.E. Wanker. 2001. *Mol. Biol. Cell.* 12:1393–1407.

^jHazeki, N., T. Tsukamoto, I. Yazawa, M. Koyama, S. Hattori, I. Someki, T. Iwatsubo, K. Nakamura, J. Goto, and I. Kanazawa. 2002. *Biochem. Biophys. Res. Com.* 294:429–440.

^kKim, S., E.A. Nollen, K. Kitagawa, V.P. Bindokas, and R.I. Morimoto. 2002. *Nature Cell Biol.* 4:826–831.

^lCummings, C.J., M.A. Mancini, B. Antalffy, D.B. DeFranco, H.T. Orr, and H.Y. Zoghbi. 1998. *Nat. Genet.* 19:148–154.

^mVerhoef, L.G., K. Lindsten, M.G. Masucci, and N.P. Dantuma. 2002. *Hum. Mol. Genet.* 11:2689–2700.

ⁿBichelmeier, U., T. Schmidt, J. Hübener, J. Boy, K. Häbig, S. Poths, M. Bonin, M. Knipper, W.J. Schmidt, J. Wilbertz, et al. 2007. *J. Neurosci.* 27:7418–7428.

^oNollen, E.A., S.M. Garcia, G. van Haften, S. Kim, A. Chavez, R.I. Morimoto, and R.H. Plasterk. 2004. *Proc. Natl. Acad. Sci. USA.* 101:6403–6408.

^pGoehler, H., M. Lalowski, U. Stelzl, S. Waelter, M. Stroedicke, U. Worm, A. Droege, K.S. Lindenberg, M. Knoblich, C. Haenig, et al. 2004. *Mol. Cell* 15:853–865.

^qYamada, M., T. Sato, T. Shimohata, S. Hayashi, S. Igarashi, S. Tsuji, and H. Takahashi. 2001. *Am. J. Pathol.* 159:1785–1795.

^rSittler, A., S. Wälter, N. Wedemeyer, R. Hasenbank, E. Scherzinger, H. Eickhoff, G.P. Bates, H. Lehrach, and E.E. Wanker. 1998. *Mol. Cell* 2:427–436.

^sCooper, J.H. 1969. *J. Clin. Path.* 22:410–413.

^tTanaka, M., Y. Machida, S. Niu, T. Ikeda, N.R. Jana, H. Doi, M. Kurosawa, M. Nekooki, and N. Nukina. 2004. *Nat. Med.* 10:148–154.

^uDing, Q., J.J. Lewis, K.M. Strum, E. Dimayuga, A.J. Bruce-Keller, J.C. Dunn, and J. N. Keller. 2002. *J. Biol. Chem.* 277:13935–13942.