

Supplementary material

DOES HUMAN NEUROGLOBIN ACT ONLY AS A SCAVENGER? REACTIVITY AND ENDOGENOUS MODIFICATION BY NITRITE AND HYDROGEN PEROXIDE*

Stefania Nicolis^{*}, Enrico Monzani^{*}, Chiara Ciaccio[†], Paolo Ascenzi[‡], Luc Moens[§], and Luigi Casella^{*}

^{*}Dipartimento di Chimica Generale, Università di Pavia, Via Taramelli 12, 27100 Pavia, Italy,

[†]Dipartimento di Medicina Sperimentale e Scienze Biochimiche, Università di Roma-Tor Vergata,

Via Montpellier 1, 00133 Roma, Italy, [‡]Dipartimento di Biologia, Università “Roma Tre”, Viale

Guglielmo Marconi 446, 00146, Roma, Italy, [§]Department of Biomedical Sciences, University of

Antwerp, Universiteit 1, B-2610 Antwerp, Belgium

Table S1. Mass Spectrometry Data of the Modified Peptides. Molecular weights and *m/z* values of the modified peptide fragments identified in NGB derivatives by MS analysis. ^aThe mass increments with respect to the molecular weights of the unmodified peptides are indicated in parentheses.

Modified residues	Peptides	MW (amu) ^a	<i>m/z</i> Bicharged ions
Y44-NO ₂	31-47	2097.4 (+ 45)	1049.7
Y88-NO ₂	68-94	2953.3 (+ 45)	1477.6
Y115-NO ₂	103-119	1949.2 (+ 45)	975.6
Y137-NO ₂	131-146	1783.0 (+ 45)	892.5
C46-SO ₂ H	31-47	2084.4 (+ 32)	1043.2
C46-SO ₂ H/Y44-NO ₂	31-47	2129.4 (+ 77)	1065.7
C46-SO ₃ H	31-47	2100.4 (+ 48)	1051.2
C46-SO ₃ H/Y44-NO ₂	31-47	2145.4 (+ 93)	1073.7
C55-SO ₂ H	48-66	2239.4 (+ 32)	1120.7
C55-SO ₃ H	48-66	2255.4 (+ 48)	1128.7
C120-SO ₂ H	120-130	1165.3 (+ 32)	583.7
C120-SO ₃ H	120-130	1181.3 (+ 48)	591.7