Supporting Information

Sun et al. 10.1073/pnas.0710867105




Fig. S1. Steady-state FRET and the effect of Imp α 1 dye labeling on cargo interaction time. (A) FRET between Alexa Fluor 568–NLS–2xGFP(4C) (100 nM) and Alexa Fluor 647–Imp α 1 (1 μ M). Blue, NLS–2xGFP(4C) alone; red, NLS–2xGFP(4C) + Imp α 1; green, Imp α 1 alone. EX = 568 nm. (B) Donor emission quenching due to FRET ($K_D = 46 \pm 3$ nM). For comparison, the K_D is \approx 40 nM in the absence of dyes (1). The data were fit to: normalized fluorescence intensity = $I_0 + (1 - I_0)(K_D/(K_D + [Imp \alpha 1]))$. [Alexa Fluor 568–NLS–2xGFP(4C)] = 100 nM. (C and D) NPC interaction time histograms for Alexa Fluor 647–NLS–2xGFP(4C) in the presence of (C) Alexa Fluor 568-Imp α 1 (7.9 \pm 0.3 ms; n = 459) and (D) Imp α 1 (7.1 \pm 0.1 ms; n = 502). Cargo interaction frequencies (2) for both C and D were 2.4 \pm 0.3 events S^{-1} , μ m⁻¹. [NLS–2xGFP(4C)] = 0.1 nM; [Imp α 1] = 250 nM; [Ran] = 2 μ M; [GTP] = 1 mM; [Imp β 1] = 0.5 μ M; [NTF2] = 1 μ M.

- 1. Catimel B, et al. (2001) Biophysical characterization of interactions involving importin-alpha during nuclear import. J Biol Chem 276:34189–34198.
- 2. Yang W, Musser SM (2006) Nuclear import time and transport efficiency depend on importin β concentration. J Cell Biol 174:951–961.




Fig. S2. Single molecule FRET efficiency of coverslip-adsorbed Alexa Fluor 568–Imp α 1/Alexa Fluor 647–NLS–2xGFP(4C) complexes. For convenience, the FRET efficiency, E, was defined as $E = I_A/(I_A + I_D)$, where I_D and I_A denote the fluorescence emission intensities observed in the donor and acceptor channels, respectively [Ha T (2001) Single-molecule fluorescence resonance energy transfer. Methods 25: 78–86]. These values were corrected for background noise and the donor emission in the acceptor channel (cross-talk). No detectable acceptor emission was observed in the donor channel. As estimated from a Gaussian fit, the FRET efficiency was 50 \pm 9% (n = 100).





Fig. S3. Decomposition of Imp α 1/cargo complexes (A \rightarrow B \rightarrow C). (A) Donor dyes on Imp α 1. The definition of the three states (A–C) and the global kinetic fit when the donor dyes were on Imp α 1 were described in the main text. The graph is reproduced from Fig. 1D. Global fitting yielded $\tau_1 = 6.7 \pm 0.3$ ms (A \rightarrow B) and $\tau_2 = 2.7 \pm 0.8$ ms (B \rightarrow C) (n = 132). [Alexa Fluor 568–Imp α 1] = 0.1 nM; [Alexa Fluor 647–NLS–2xGFP(4C)] = 250 nM. (B) Donor dyes on the cargo. The identical experiment as in A, except that the donor and acceptor dye positions were switched, yielding $\tau_1 = 6.5 \pm 0.3$ ms and $\tau_2 = 1.3 \pm 0.5$ ms (n = 122). Comparing with A, note that the Imp α 1/cargo complex dissociation times (τ_1) were identical (within error), as expected. However, the free cargo dissociated from the NPC faster than cargo-free Imp α 1 (compare τ_2 values). Also note that $\tau_1 + \tau_2 = 7.8$ ms, consistent with the cargo interaction time reported in Fig. S1C. [Alexa Fluor 568–NLS–2xGFP(4C)] = 0.1 nM; [Alexa Fluor 647–Imp α 1] = 250 nM (A and B) The position of the molecule with the acceptor dyes was unknown after Imp α 1/cargo complex dissociation. Shown is one possible spatial arrangement for the B and C states. The B state could not be identified for \approx 14% of the dissociation events in A and \approx 21% in B. These data were not included in the analysis, but indicate that the τ_2 values are upper limits. Red, A; blue, B; green, C. [CAS] = 1.3 μ M; [Ran] = 2 μ M; [GTP] = 1 mM; [Imp μ 1] = 0.5 μ M; [NTF2] = 1 μ M.



Movie \$1. Shows the transport event of Fig. 1.A. Pixels are 240 nm², each frame was acquired in 2 ms, and the playback speed is 500× slower than real-time. Note: This QuickTime video was made from 16-bit CCD camera data converted to 8-bit. The QuickTime software on late-model Apple computers may destroy the pixelation for video display. If difficulties are encountered in attempts to observe the raw pixelated data, try using the RealPlayer software (Apple) or QuickTime on a PC.

Movie S1 (MOV)

Table S1. Effect of CAS concentration on dissociation and transport efficiencies

CAS, nM	Result	Total, n	Dissociation efficiency, %	Final de Cytoplasm	stination Nucleoplasm	Transport
				•	•	
-25% Gly 0.1 nM Ale	cerol exa Fluor 568-Imp α 1 and 250 n	M Alexa Fluor	· 647-NLS-2xGF	P(4C)		
0	Did not dissociate at NPC	144	- 0	65 (45 ± 4%)	79 (55 ± 4%)	55 ± 4
0	Did dissociate at NPC	0	· · · · · · · · · · · · · · · · · · ·	0	0	33 ± 4
0.01	Did not dissociate at NPC Did dissociate at NPC	82	- 0	43 (52 ± 6%) 0	39 (48 ± 6%) 0	48 ± 6
0.02 0.08 0.2 0.8	Did not dissociate at NPC	97		44 (45 ± 5%)	53 (55 ± 5%)	
	Did dissociate at NPC	1	- 1 ± 1	0	1	55 ± 5
	Did not dissociate at NPC	103	- 1 ± 1	56 (54 ± 5%)	47 (46 ± 5%)	46 ± 5
	Did dissociate at NPC Did not dissociate at NPC	1 87		0 45 (52 ± 5%)	1 42 (48 ± 5%)	
	Did dissociate at NPC	0	- 0	0	0	48 ± 5
	Did not dissociate at NPC	76	- 1 ± 1	39 (51 ± 5%)	37 (49 ± 5%)	49 ± 6
	Did dissociate at NPC	1	1 = 1	0	1	47 ± 0
	Did not dissociate at NPC Did dissociate at NPC	92 3	3 ± 2	49 (53 ± 5%) 3	43 (47 ± 5%)	46 ± 5
	Did not dissociate at NPC	82		47 (57 ± 5%)	35 (43 ± 5%)	
3.2	Did dissociate at NPC	7	8 ± 3	2	5	44 ± 5
3.0	Did not dissociate at NPC	65	- 16 ± 4	41 (63 ± 6%)	24 (37 ± 6%)	55 ± 6
	Did dissociate at NPC Did not dissociate at NPC	12 57		1 31 (54 ± 6%)	11 26 (46 ± 6%)	
16	Did dissociate at NPC Did dissociate at NPC	22	- 28 ± 5	5 (54 ± 6%)	26 (46 ± 6%) 17	54 ± 6
32	Did not dissociate at NPC	53	38 ± 5	38 (72 ± 6%)	15 (28 ± 6%)	48 ± 5
/ <u>~</u>	Did dissociate at NPC	32	30 ± 3	6	26	70 ± 3
65	Did not dissociate at NPC Did dissociate at NPC	65 67	51 ± 4	54 (83 ± 5%) 9 (13 ± 4%)	11 (17 ± 5%) 58 (87 ± 4%)	52 ± 4
	Did not dissociate at NPC	58		48 (83 ± 5%)	10 (17 ± 5%)	
100	Did dissociate at NPC	52	47 ± 5	11 (21 ± 6%)	41 (79 ± 6%)	46 ± 5
130	Did not dissociate at NPC	73	53 ± 4	65 (89 ± 4%)	8 (11 ± 4%)	47 ± 4
130	Did dissociate at NPC	81	33 1 4	16 (20 ± 5%)	65 (80 ± 4%)	4/ = 4
160	Did not dissociate at NPC Did dissociate at NPC	43 33	- 43 ± 6	30 (70 ± 7%) 4 (12 ± 6%)	13 (30 ± 7%) 29 (88 ± 6%)	51 ± 6
	Did not dissociate at NPC	72	40 4	65 (90 ± 4%)	7 (10 ± 4%)	
320	Did dissociate at NPC	66	- 48 ± 4	8 (12 ± 4%)	58 (88 ± 4%)	47 ± 4
1300	Did not dissociate at NPC	230	- 55 ± 2	189 (82 ± 2%)	41 (18 ± 2%)	53 ± 2
	Did dissociate at NPC	280		53 (19 ± 2%)	227 (81 ± 2%)	
+25% G	lycerol					
0.1 nM	Alexa Fluor 568-NLS-2xGFP(4C) and 250 nM	Alexa Fluor 647	_		
0	Did not dissociate at NPC	128	o	59 (46 ± 4%)	69 (54 ± 4%)	54 ± 4
	Did dissociate at NPC Did not dissociate at NPC	0 177		0 101 (57 ± 5%)	0 76 (43 ± 5%)	
5	Did dissociate at NPC	16	8 ± 2	5 (31 ± 11%)	11 (69 ± 11%)	45 ± 4
20	Did not dissociate at NPC	189	35 ± 3	134 (71 ± 3%)	55 (29 ± 3%)	46 ± 4
20	Did dissociate at NPC	102	33 ± 3	24 (24 ± 4%)	78 (76 ± 4%)	70 ± 4
80	Did not dissociate at NPC Did dissociate at NPC	112 117	- 51 ± 3	91 (81 ± 4%) 25 (21 ± 4%)	21 (19 ± 4%) 92 (79 ± 4%)	49 ± 3
	Did dissociate at NPC Did not dissociate at NPC	123		96 (78 ± 4%)	27 (22 ± 4%)	
150	Did dissociate at NPC	178	— 59 ± 3	52 (29 ± 3%)	126 (71 ± 3%)	- 51 ± 3
1300	Did not dissociate at NPC	215	61 ± 2	178 (83 ± 3%)	37 (17 ± 3%)	52 ± 2
1300	Did dissociate at NPC	342	01 ± 2	87 (25 ± 2%)	255 (75 ± 2%)	32 ± 2
Glycerol						
.1 nM Al	exa Fluor 568-Imp α1 and 250 r		or 647-NLS-2xG			
0.1	Did not dissociate at NPC	53	- 4 ± 3	$31 (58 \pm 7\%)$	22 (42 ± 5%)	- 44 ± 7
	Did dissociate at NPC Did not dissociate at NPC	62		0 35 (54 ± 6%)	2 27 (46 ± 6%)	
1	Did dissociate at NPC	3	5 ± 3	$\frac{33(34 \pm 6\%)}{0}$	3	- 46 ± 6
10 50	Did not dissociate at NPC	71	12 ± 4	38 (54 ± 5%)	33 (46 ± 5%)	5 2 ± 4
	Did dissociate at NPC	11	- 13 ± 4	1	10	- 52 ± 6
	Did not dissociate at NPC	57	- 44 ± 5	40 (70 ± 6%)	17 (30 ± 6%)	- 49 ± 5
	Did dissociate at NPC	44		12 (27 ± 7%)	32 (73 ± 7%)	
100	Did not dissociate at NPC Did dissociate at NPC	56 63	- 53 ± 5	$51 (91 \pm 4\%)$ 9 (14 ± 4%)	5 (8 ± 4%) 54 (86 ± 4%)	-50 ± 5
500	Did not dissociate at NPC	43	- 56 ± 5	$32 (74 \pm 7\%)$	$11 (26 \pm 7\%)$	

[Ran] = 2 μ M; [GTP] = 1 mM; [Imp β 1] = 0.5 μ M; [NTF2] = 1 μ M.

Table S2. Dissociation of Imp lpha1/cargo complexes in the absence of RanGTP and CAS

Destination	n	Interaction time, ms	Transport efficiency*	Number dissociated	Dissociation efficiency*
Nucleoplasm Cytoplasm	87 107	27 ± 4 28 ± 3	45 ± 4%	0 0	0%

[Alexa Fluor 568–NLS–2xGFP(4C)] = 0.1 nM; [Alexa Fluor 647–Imp α 1] = 250 nM; [Imp β] = 0.5 μ M; [NTF2] = 1 μ M.

^{*}Transport and dissociation efficiencies were calculated from the data in the nucleoplasm and cytoplasm rows.