



SUPPLEMENTARY ONLINE DATA

Regulation of hypoxia-inducible factor-1 α by NF- κ B

Patrick VAN UDEN¹, Niall S. KENNETH¹ and Sonia ROCHA²

College of Life Sciences, Wellcome Trust Centre for Gene Regulation and Expression, MSI/WTB/JBC Complex, Dow Street, University of Dundee, Dundee DD1 5EH, Scotland, U.K.




Figure S1 HEK-293 cells were transfected with the indicated siRNA oligonucleotides and HIF-1 α and HIF-2 α mRNA was analysed by qRT-PCR

The histogram depicts relative levels of specific mRNA transcripts normalized to 18S mRNA levels. The mean + S.D. were calculated from a minimum of three independent experiments.




Figure S2 HEK-293 cells were transfected with the indicated siRNA oligonucleotides and PHD2 mRNA was analysed by qRT-PCR

The histogram depicts relative levels of specific mRNA transcripts normalized to 18S mRNA levels. Results are representative of a minimum of three independent experiments.

Received 29 February 2008/3 April 2008; accepted 4 April 2008
 Published as BJ Immediate Publication 4 April 2008, doi:10.1042/BJ20080476

¹ These authors contributed equally to this work.
² To whom correspondence should be addressed (email s.rocha@dundee.ac.uk).