

## Treponemal infection in the Australian Northern Territory aborigines

M. F. GARNER,<sup>1</sup> J. L. BACKHOUSE,<sup>2</sup> P. M. MOODIE,<sup>3</sup> & G. J. TIBBS<sup>4</sup>

*For many years, a moderate to high prevalence of positive reactions to standard tests for syphilis (STS) has been found in sera from the aboriginal inhabitants of the Northern Territory of Australia. In the survey reported here, people living in 18 areas of the Northern Territory were sampled to determine if the positive STS results were due to treponemal infection and, if so, to what extent this was endemic syphilis, yaws, or venereal syphilis. The results of the Treponema pallidum immobilization test were taken to indicate the presence or absence of treponemal disease in the subject. Clinically, no cases of active treponemal infection were seen, though findings on old infections were difficult to interpret in the older age groups. The prevalence of treponemal infection varied from 3.4% to 58.0% in the areas tested. No serological evidence of treponemal infection in children under 15 years of age was found in 6 areas, while in 12 areas it varied from 2.2% to 32.6%. Sera from 60 people who had no clinical signs of treponemal infection gave high-titre CWR and VDRL test results. It was concluded that yaws, endemic syphilis, and probably venereal syphilis are to be found in the aboriginal population.*

Clinical yaws has been recorded for many years among the aboriginal population in the humid north of the Northern Territory of Australia (Fig. 1), particularly up to about 1945. A condition in the arid south of the Northern Territory, sometimes referred to as *irkinja* (Hackett, 1936), has also been recorded over a long period, and was generally thought to be either climatically modified yaws or a form of endemic syphilis. Those working in laboratories where sera from aborigines are tested have long realized that a moderate to high prevalence of positive standard serological tests for syphilis (STS) is found among these people. This has caused speculation as to whether these reactions indicate the spread of venereal syphilis in the aboriginal population, perhaps as a result of the gradual disappearance of yaws and endemic syphilis.

The survey reported here was undertaken to determine if the positive results were all due to treponemal infection, and if so, the proportions that were caused by yaws, endemic syphilis, and venereal syphilis. The field work was carried out under the direction of the School of Public Health and Tropical Medicine, University of Sydney, and the sera were sent to the Institute of Clinical Pathology and Medical Research for testing.

The Northern Territory was divided into two strata, corresponding to the administrative divisions, the dividing line corresponding closely to the 19th parallel of latitude. The northern stratum, which has an annual rainfall greater than 38 cm, includes areas of monsoonal coast, savanna woodland, savanna, and grassy steppe; the southern stratum has an annual rainfall less than 38 cm and includes dry grassland, semidesert, and desert. The temperature in the whole Northern Territory varies from 36.7°C on summer days to 3.3°C on cold winter nights.

Traditional aboriginal culture was based on a nomadic existence and a hunting-collecting economy. It was very poor in the material sense, but rich in magic and symbolism, mythology, and art, and there was a very close personal identification with the natural environment. The traditional nomadic existence was carried on by small local groups of one,

<sup>1</sup> Senior Specialist Serologist, Venereal Disease Reference Laboratory, Institute of Clinical Pathology and Medical Research, Lidcombe, New South Wales, Australia.

<sup>2</sup> Microbiologist, Venereal Disease Reference Laboratory, Institute of Clinical Pathology and Medical Research, Lidcombe, New South Wales, Australia.

<sup>3</sup> Medical Officer (Research and Teaching), Department of Tropical Medicine, School of Public Health and Tropical Medicine, University of Sydney, Sydney, New South Wales, Australia.

<sup>4</sup> Lately Medical Officer, Northern Territory Medical Service, Darwin, Northern Territory, Australia.

two, or three families, who travelled independently through their territory in an endless cycle determined by season, availability of food, and ceremonial occasions. Personal contact outside the local group was infrequent and brief, and such contact as did occur, either personal or sexual, was governed by kinship and totemic rules. This traditional culture was shattered by contact with European settlement and attitudes. Rules and taboos that were accepted, and also easily enforced, in the nomadic situation were out of context in multiracial settlements, such as now exist. The aboriginal community, still mainly a rural population, became settled in relatively small groups and was generally dependent upon government or mission staff and cattle-station managements for food, shelter, employment, health and medical services, and education. The standard of living varied greatly but generally was very low materially and economically with little appreciation of hygiene.

Many families tended to move between adjacent localities but were not true nomads.

The estimated aboriginal population of the Northern Territory in late 1968 was 21 119, of which 90% were accessible. A sampling frame of all aboriginal communities of 50 or more persons was constructed from available data and the total sampling frame population was estimated to be 15 548; a 10% sample of this population was surveyed. Each stratum was divided into substrata, substratification being based partly on geography and partly on the type of aboriginal community. One or two sampling stations were selected from each substratum by probability sampling. Altogether, 11 communities were sampled in the northern stratum, of which 9 had a population greater than 100 and 2 had a population of 50-99; the corresponding data for the southern stratum are 7 areas, 5 having a population greater than 100 and 2 having a population of 50-90 (Fig. 1).




Fig. 1. Location of main towns (■) and the 1968 sampling stations (●) in the Northern Territory of Australia.

The households examined at each sampling station were drawn from a table of random numbers. A household consisted of all the people living in a dwelling with or without attached shelters, or in a group of small shelters. These people generally ate together and shared the basic necessities of daily living. Usually, this meant an extended family, with or without some attached persons. The communities were divided into three groups—namely, missions, settlements, and pastoral properties.

Each person included in this survey received as complete a clinical examination as possible. The female pudenda and the male genitalia were not always examined since it was felt that the subjects' cooperation would have been lost if this had been insisted upon. Comments in this report, such as "no clinical evidence of venereal syphilis", must be considered with this limitation in mind since most chancres or condylomata, if present, would not have been seen. No clinical cases of florid yaws, endemic syphilis, or venereal syphilis were seen, though some subjects showed signs of previous yaws infection. However, active yaws was seen during the course of this survey in people who were outside the survey sample. One case of "burned out" gangosa was found, and a few cases of marked anterior bowing of the tibia were seen even in young people. There were many cases of slight bowing of the tibia, skin eruptions, plantar hyperkeratoses, leg scars, perforated nasal septa and perforated and deformed palates. These clinical findings were difficult to interpret, especially in the older age groups, as in many instances they could not be distinguished from conditions resulting from poor nutrition, old injuries, and camp-fire burns. Perforated nasal septa referred to are the high septal perforations, which appeared to be pathological, rather than the low, soft-tissue septal perforations that had been made as a tribal custom for purposes of ornamentation. No cases of congenital, cardiovascular, or central nervous system syphilis were seen. Over the past 10 years, penicillin has been widely used, especially among the children. A clinical diagnosis of past or present treponemal disease was made in 4.1% of the total sample, and was made more often in the southern stratum (5.0%) than in the northern stratum (3.5%). Otherwise, differences in clinical signs between people in the various sub-strata or other groupings were minimal.

Sera were received from 1 570 aboriginal people aged more than 1 year. Full serological tests were carried out on only 1 542 sera and these results form the basis of this report. The following tests were

performed on each serum: cardiolipin Wassermann reaction (CWR), Venereal Disease Research Laboratory (VDRL) test, Reiter protein complement-fixation (RPCF) test, and a *Treponema pallidum* immobilization (TPI) test. Sera found to be positive to the CWR or VDRL tests were titrated. A positive TPI test result was regarded as evidence of treponemal infection and all other test results were compared with it. A previous suggestion that there was a high reactor rate to the gonococcal complement-fixation (GCF) test among the aborigines, and hence that the positive STS were caused by venereal syphilis, prompted us to perform the GCF test on each of the 1 542 sera.

## RESULTS

### *Results in total population tested*

Of the 1 542 sera tested, 765 were from males and 777 from females. The prevalence of treponemal infection in the whole group was 23.0%, 354 sera being positive to the TPI test. Males showed a prevalence of 24.7% and females 21.2%, with 189 and 165 sera, respectively, giving positive reactions to TPI tests (Table 1). The endemic treponematoses are diseases of childhood, usually contracted before the age of 15 years. For the purposes of this report, children are those aged 1–14 years and adults those aged 15 years and over. Positive TPI reactions were found with sera from 36 (5.6%) of 643 children compared with 318 (35.4%) of 899 adults (Table 1).

### *Results in each age group*

Tables 1 and 2 show the prevalence of treponemal infection in the various age groups and in males and females in each group. The percentage prevalence of infection tended to increase with age, except for those aged 5–9 years and 60 years and over. As expected, it was lower in females than in males, except those aged 5–9 years, in whom the male prevalence was 0.8% compared with 6.6% in females. In the young children aged 1–4 years the prevalence of infection in males was much higher than in females (6.9% and 3.9%, respectively).

### *Results in each area tested*

The number and proportion of TPI-reactive sera found in each area, and in males and females in each area, are shown in Table 3, while Table 4 compares these results in adults and children. The highest prevalence of treponemal infection (58.0%) was

Table 1. TPI test results in relation to age and sex of the population studied in the whole of the Northern Territory of Australia

Age (years)	No. of sera tested	TPI-reactive		Males			Females		
		No.	%	No. of sera tested	TPI-reactive		No. of sera tested	TPI-reactive	
					No.	%		No.	%
1-14	643	36	5.6	321	17	5.3	322	19	5.9
15-29	386	69	17.9	194	41	21.1	192	28	14.6
30-44	280	106	37.9	118	47	39.8	162	59	36.4
45-59	145	95	65.5	79	53	67.1	66	42	63.6
≥ 60	88	48	54.5	53	31	58.5	35	17	48.6
total adults	899	318	35.4	444	172	38.7	455	146	32.1
total sera	1 542	354	23.0	765	189	24.7	777	165	21.2

found at Maningrida, a large multi-tribal settlement in the northern stratum. Here, the prevalence rate was also the highest for both males (62.1%) and females (52.4%). Two other areas in the northern stratum had high prevalence rates—namely, Robinson River (44.4%) and Mainoru (44.0%). Sera from 50.0% of the males and 38.9% of the females at Robinson River gave positive TPI reactions; the corresponding proportions for Mainoru were 38.5% and 50.0%. Of the 18 areas tested, 11 had over 20% of sera reactive to the TPI test; the same 11 areas had over 20% prevalence of infection in males; 9 of these areas had over 20% prevalence of infection in

females. The lowest rate of infection was found at Hermannsburg Mission in the southern stratum, where of 59 sera tested only 2, from an adult male and an adult female, were positive to the TPI test.

In adults, the highest prevalence of treponemal infection was seen at Maningrida (77.2%), then Mainoru (67.9%), Robinson River (66.7%), and Bamyili (54.1%), all in the northern stratum, and Yuendumu (49.2%) in the southern stratum. Of the 18 areas tested, only 6 had prevalence rates less than 20% in adults, 4 of these being in the southern stratum.

In the children, 14 (32.6%) of 43 sera tested at Maningrida were reactive to the TPI test, 3 (13.6%)

Table 2. Results of TPI test in relation to age and sex in the children studied

Age (years)	No. of sera tested	TPI-reactive		Males			Females		
		No.	%	No. of sera tested	TPI-reactive		No. of sera tested	TPI-reactive	
					No.	%		No.	%
1-4	204	11	5.4	102	7	6.9	102	4	3.9
5-9	239	9	3.8	118	1	0.8	121	8	6.6
10-14	200	16	8.0	101	9	8.9	99	7	7.1
total	643	36	5.6	321	17	5.3	322	19	5.9

Table 3. Number and percentage of TPI-reactive sera in each area, by sex

Area	No. of sera tested	TPI-reactive		Males			Females		
		No.	%	No. of sera tested	TPI-reactive		No. of sera tested	TPI-reactive	
					No.	%		No.	%
Maningrida	100	58	58.0	58	36	62.1	42	22	52.4
Angurugu	87	9	10.3	39	5	12.8	48	4	8.3
Milingimbi	101	28	27.7	38	11	29.0	63	17	27.0
Bathurst Island	100	26	26.0	43	12	27.9	57	14	24.6
Delissaville	90	11	12.2	48	8	16.7	42	3	7.1
Roper River	98	22	22.5	47	10	21.3	51	12	23.5
Bamyili	102	33	32.4	54	17	31.5	48	16	33.3
Bagot	100	22	22.00	57	14	24.6	43	8	18.6
Victoria River Downs	110	31	28.2	54	17	31.5	56	14	25.0
Robinson River	36	16	44.4	18	9	50.0	18	7	38.9
Mainoru	50	22	44.0	26	10	38.5	24	12	50.0
Hermannsburg	59	2	3.4	30	1	3.3	29	1	3.5
Yuendumu	95	29	30.5	42	15	35.7	53	14	26.4
Areyonga	99	5	5.1	44	4	9.1	55	1	1.8
Amoonguna	105	9	8.6	58	5	8.6	47	4	8.5
Mary Vale	99	12	12.1	52	4	7.7	47	8	17.0
Aileron	70	14	20.0	33	7	21.2	37	7	18.9
Alroy Downs	41	5	12.2	24	4	16.7	17	1	5.9

of 22 at Mainoru and 2 (13.3%) of 15 at Robinson River. No children had sera giving positive TPI reactions at Bamyili and Yuendumu, where high prevalence rates were found in the adults. In 6 of the 18 areas studied, no child had a serum that gave positive reactions to the TPI test; 5 of these areas were in the southern stratum. In each of the two remaining areas in the southern stratum, serum from only one child gave a positive TPI test result.

*Results in the northern and southern strata*

Table 5 shows the number of sera reactive in the TPI test for the total population studied and for adults and children in each stratum.

In the northern stratum, of 974 sera tested, 278 (28.5%) were found to be positive to the TPI test, the adults having a prevalence rate of 42.4%. The rates for the southern stratum were approximately half those for the northern stratum for these two groups. The most marked contrast in prevalence of infection occurred in the children. In the northern

stratum, 34 (8.5%) of 398 sera reacted positively to TPI tests compared with 2 (0.8%) of 245 sera in the southern stratum.

*Comparison of test results with the TPI test*

Table 6 compares the results of the CWR, RPCF, and VDRL tests with the TPI test. Altogether, 354 sera gave positive reactions to the TPI test. The number of sera reactive to both TPI and VDRL tests was 265; to TPI and RPCF tests, 227; and to TPI and CWR tests, 207. The VDRL test was the only one to show complete agreement with the TPI test in any individual area tested, and then in only 4 of the 18 areas.

Biological false positive (BFP) reactions occurred in 61 sera as follows: CWR in 1, VDRL in 16, CWR and VDRL in 4, and RPCF in 40 sera.

A total of 60 sera, reactive in all tests, had high CWR and/or VDRL test titres. Test titres were regarded as high when the CWR titre was 40, 80, 160, or 320 and the VDRL test was positive for serum

Table 4. Number and percentage of TPI-reactive sera in adults and children in each area

Area	Total no. of sera tested	Adults			Children		
		No. of sera tested	TPI-reactive		No. of sera tested	TPI-reactive	
			No.	%		No.	%
Maningrida	100	57	44	77.2	43	14	32.6
Angurugu	87	45	8	17.8	42	1	2.4
Milingimbi	101	57	27	47.4	44	1	2.3
Bathurst Island	100	62	22	35.5	38	4	10.5
Delissaville	90	54	8	14.8	36	3	8.3
Roper River	98	53	19	35.9	45	3	6.7
Bamyili	102	61	33	54.1	41	0	0
Bagot	100	62	21	33.9	38	1	2.6
Victoria River Downs	110	76	29	38.2	34	2	5.9
Robinson River	36	21	14	66.7	15	2	13.3
Mainoru	50	28	19	67.9	22	3	13.6
Hermannsburg	59	21	2	9.5	38	0	0
Yuendumu	95	59	29	49.2	36	0	0
Areyonga	99	57	5	8.8	42	0	0
Amoonguna	105	59	8	13.6	46	1	2.2
Mary Vale	99	63	11	17.5	36	1	2.8
Aileron	70	42	14	33.3	28	0	0
Alroy Downs	41	22	5	22.7	19	0	0

Table 5. Number and percentage of TPI-reactive sera in northern and southern strata; totals for adults and children

Northern Territory division	Total no. of sera tested	TPI-reactive		Adults			Children		
		No.	%	No. of sera tested	TPI-reactive		No. of sera tested	TPI-reactive	
					No.	%		No.	%
northern stratum	974	278	28.5	576	244	42.4	398	34	8.5
southern stratum	568	76	13.4	323	74	22.9	245	2	0.8

diluted 1 : 8, 1 : 16, or 1 : 32. There were only 4 high-titre reactions among children under 15 years of age—namely, in 2 males at Maningrida and 1 female each at Robinson River and Mainoru; all were in the 10–14-year age group and showed no significant clinical indications of infection.

#### GCF test results

A total of 21 sera were reactive in the GCF test; 10 of these also gave positive reactions to the TPI test.

#### DISCUSSION

It is not possible from the clinical findings and serological results of this survey to determine the exact separate prevalence of endemic syphilis, yaws, and venereal syphilis in the aboriginal population tested in the Northern Territory of Australia. Because of the climatic conditions, the absence of congenital syphilis, and the lack of clinical evidence of venereal syphilis, it is reasonable to assume that the

Table 6. Comparison of results of CWR, RPCF, and VDRL tests with TPI test

Area	No. of sera tested	Positive											
		CWR	RPCF	VDRL	TPI	In all tests	CWR, RPCF, TPI only	CWR, VDRL, TPI only	CWR, TPI only	VDRL, RPCF, TPI only	RPCF, TPI only	VDRL, TPI only	TPI only
Maningrida	100	32	28	38	58	27	—	4	1	1	—	6	19
Angurugu	87	1	4	3	9	1	—	—	—	—	2	1	5
Milingimbi	101	21	22	23	28	17	—	4	—	—	1	1	5
Bathurst Island	100	16	26	17	26	14	—	2	—	—	5	1	4
Delissaville	90	7	8	9	11	6	—	—	—	—	—	1	4
Roper River	98	14	15	21	22	11	—	2	—	3	—	3	3
Bamyili	102	18	25	32	33	15	—	3	—	8	1	5	1
Bagot	100	15	20	21	22	13	—	1	—	3	1	3	1
Victoria River Downs	110	22	21	23	31	19	—	3	—	1	—	—	8
Robinson River	36	12	14	14	16	11	—	1	—	—	1	1	2
Mainoru	50	11	13	12	22	10	—	1	—	1	2	—	8
Hermannsburg	59	1	4	4	2	1	—	—	—	—	—	1	—
Yuendumu	95	19	25	29	29	13	1	4	—	5	4	2	—
Areyonga	99	4	10	5	5	2	—	1	—	1	—	—	1
Amoonguna	105	3	6	9	9	3	—	—	—	2	—	2	2
Mary Vale	99	4	8	8	12	3	—	1	—	2	—	2	4
Aileron	70	8	12	13	14	8	—	—	—	3	—	1	2
Alroy Downs	41	4	6	5	5	4	—	—	—	1	—	—	—

disease in the southern stratum is endemic syphilis and that that in the northern stratum is yaws.

It has been found that history or clinical signs are no guide to past treponemal disease as indicated by the TPI test reaction. On the usual assumption that the TPI test reaction is specific, it can be stated that few of the signs elicited were peculiar to positive TPI test reactors, and those that were occurred too seldom to be useful indicators of infection, e.g., perforated nasal septum and bone deformity. Similarly, there were no significant clinical signs to account for high-titre reactivity in persons whose sera showed this phenomenon. There was a negative association of stillbirths to parous women whose sera were reactive in the TPI test, suggesting endemic treponemal disease rather than venereal syphilis. There was a positive association between the past history of leprosy and positive TPI tests in all age groups. The most likely explanation is that cases of leprosy and cases of treponemal infection came from the same micropopulations because of common environmental

factors, and that the association has been produced artificially by combining heterogeneous micro-populations.

Statistically, there was a strong association between wives and husbands whose sera gave positive reactions to the TPI test. This could have arisen from the origin of husbands and wives in the same community in areas where childhood yaws was prevalent, or from cross-infections between husband and wife. If the latter occurred, the disease could have been either yaws or venereal syphilis, particularly in marriages between very young brides and older men, i.e., yaws infection could have been transmitted from recently infected girls to susceptible older men, or from first or relapsing latent infections in older men to uninfected girls. The transmission of venereal syphilis could have followed a similar pattern.

There was no association of positive reactions to TPI tests amongst husbands and wives in the cattle-station populations. Here, the partners were closer

in age, there was less regard for tribal custom and taboos, husbands were frequently absent in cattle camps for weeks at a time, and there was a general lack of European supervision of camp behaviour. High-titre seroreactivity, suggestive of active syphilis in adults, was most common in the females living on cattle stations.

In the northern stratum, TPI reactivity in mother and child was considered for offspring aged 1-14 and 15-29 years. In all, 15 (14.4%) of 104 children of TPI-reactive mothers were themselves positive, compared with 11 (4.8%) of 230 children whose mothers' sera were nonreactive in the TPI test. However, in the 15-29-year age group, all TPI-positive offspring had positive mothers, suggesting that yaws cross-infection between mother and child may have been usual prior to 1953 before penicillin was used widely, but that it no longer occurs to the same extent. Children were also more likely to be TPI-positive in households where other children were also TPI-positive.

The difference in prevalence of treponemal infection between the northern and southern strata is marked. The endemic syphilis of the southern areas seems to be disappearing; only 2 of 245 children tested showed serological evidence of infection. By contrast, the prevalence of yaws varied greatly from place to place among the children in the northern stratum, from as high as 32.6% at Maningrida to 0 at Bamyili. This indicated active foci of infection in some areas and thus a continuing condition.

Endemic syphilis and yaws are both diseases usually contracted in childhood. Reagin, the antibody detected by the CWR and VDRL tests, fades naturally with time, even without treatment. Thus it is unlikely that adults with high reagin test titres have either endemic syphilis or yaws, especially when they have no clinical signs of either disease in a florid form, unless the endemic treponematoses have altered in their clinical signs in these people. Venereal syphilis is well known to occur with relatively fleeting signs in the primary, secondary, and early latent stages and the serum in these stages usually has a high reagin test titre. It is therefore highly probable that venereal syphilis has occurred in approximately 60 of the 354 cases of treponemal infection among persons past the age of puberty in both the northern and southern strata. Not a single high-titre reaction was found in children under 10 years of age. Another possible explanation of high-titre seroreactivity is a very marked serological relapse of old latent yaws.

The TPI test was the test of most value in this

survey. It detected 354 cases of treponemal infection and 61 sera that showed BFP reactions to the more conventional treponemal serology tests. The VDRL test results showed closer agreement with those of the TPI test than did the results of the CWR and RPCF tests, but it gave BFP reactions with 20 sera and failed to detect 89 cases of treponemal infection. As in previous yaws surveys in New Guinea (Garner & Hornabrook, 1968), the CWR and VDRL test results, both detecting the same antibody (reagin), showed marked discrepancies. The RPCF test showed a high number of false positive results, probably as the result of group antibody in the sera against nonpathogenic mouth and genital treponemes. This is a feasible explanation since the knowledge and practice of personal hygiene is limited in the people under survey.

In view of recent work on sorbents used in the fluorescent treponemal antibody absorption (FTS-ABS) test (Rathlev, 1968; Wilkinson & Ferguson, 1968), this test was not used in the present survey. Its value would have been to detect early cases of treponemal infection, but as Rathlev (1968) pointed out, it is unwise to make a diagnosis of treponemal infection on the results of a positive FTA-ABS test on a serum, with nothing else to support it serologically or, as in our survey, no clinical support.

The GCF test is generally considered to be an unreliable test, useful only in old cases of arthritis and salpingitis. The 10 sera reactive in the GCF and TPI tests may have come from people with gonorrhoea as well as treponemal infections. It does not follow that the associated positive TPI test result indicates venereal syphilis, unless there is clinical evidence to support this. Gonorrhoea, as judged by the results of the GCF test, does not appear to be a problem.

The most important question is not whether venereal syphilis exists in the Northern Territory aborigines, but is whether venereal syphilis is spreading and replacing yaws and endemic syphilis, which, on serological, clinical, and analogous evidence, are declining. The question of the extent to which venereal syphilis has replaced endemic treponemal disease remains unsolved. There is no evidence of extensive venereal syphilis, nor of extensive gonorrhoeal seroreactivity.

Syphilis, if it occurs, is probably most common in the north, is not widespread, and was only recently introduced and spread. Whatever the nature of the treponemal infection, it is usually subclinical at any point in time; it is still affecting children in the north.



As judged by seronegativity to the TPI test, 65% of adults in the sample have no acquired or cross-immunity to syphilis. In view of the increasing opportunities for venereal transmission, venereal syphilis is a real threat to the population. Its spread is probably imminent if it is not already occurring.

To eradicate endemic treponemal infection, the best method is the mass treatment of all active and latent cases and contacts in the whole area, as recommended by Guthe & Luger (1966). The people whose sera show high-titre CWR and/or VDRL test results are best treated as cases of active syphilis.

### ACKNOWLEDGEMENTS

We acknowledge the assistance of the Northern Territory Department of Health, Darwin, Dr J. Donovan, Department of Mathematical Statistics, University of Sydney, and the Computing Research Centre, CSIRO, University of Sydney. This paper is published with the permission of the Director-General of Health for Australia.

### RÉSUMÉ

#### LES INFECTIONS TRÉPONÉMIQUES CHEZ LES ABORIGÈNES DU TERRITOIRE DU NORD EN AUSTRALIE

En recherchant la présence des anticorps spécifiques dans les sérums prélevés chez 1542 aborigènes vivant dans le Territoire du Nord (Australie), on a constaté que la prévalence globale des infections tréponémiques était de 23,0%, atteignant 35,4% chez les adultes et 5,6% chez les enfants.

Dans la partie septentrionale du Territoire, où le climat se prête à la propagation du pian, le taux global d'infection tréponémique atteignait 28,5%; il était de 42,4% chez les adultes et de 8,5% chez les enfants. Dans la zone sud, apparemment plus favorable à la transmission de la syphilis endémique, les taux de prévalence étaient respectivement de 13,4% (taux global), 22,9% (adultes) et 0,8% (enfants). Soixante sérums, examinés en épreuves de

Wassermann et VDRL, ont fourni des titres d'anticorps très élevés. Aucun signe clinique net de pian, de syphilis endémique ou de syphilis vénérienne en évolution n'a été décelé dans le groupe de population étudié.

La conclusion des auteurs est que les tréponématoses responsables des modifications sérologiques sont le pian dans la partie nord du Territoire et la syphilis endémique dans le sud. La forte réactivité de 60 sérums résulte probablement d'une atteinte de syphilis vénérienne.

L'éradication des infections tréponémiques dans cette région ne pourra être obtenue que par le traitement de masse de tous les cas, actifs ou latents, et des contacts, associé à des mesures destinées à élever le niveau général d'hygiène de la population.

### REFERENCES

- Garner, M. F. & Hornabrook, R. W. (1968) *Bull. Wld Hlth Org.*, **38**, 189-195  
 Guthe, T. & Luger, A. (1966) *Derm. int.*, **5**, 179-199  
 Hackett, C. J. (1936) *Boomerang leg and yaws in Australian aborigines*, London, Royal Society of Tropical Medicine and Hygiene  
 Rathlev, T. (1968) *Brit. J. vener. Dis.*, **44**, 295-297  
 Wilkinson, A. E. & Ferguson, H. G. (1968) *Brit. J. vener. Dis.*, **44**, 291-294