

## ERRATUM

### Fatty Acid Distribution in Mesophilic and Thermophilic Strains of the Genus *Bacillus*

P. Y. SHEN, E. COLES, J. L. FOOTE, AND J. STENESH

*Department of Chemistry, Western Michigan University, Kalamazoo, Michigan 49001*

Volume 103, no. 2, p. 480, Table 2: The values shown in columns B and C under "Mesophiles" should be reversed; the "Avg chain length" values which appear under "Mesophiles" should be under "Thermophiles," and vice versa. The corrected table is as follows:

TABLE 2. General characteristics of the fatty acid distribution

Fatty acid	Mesophiles				Thermophiles			
	A	B	C	Avg	D	E	F	Avg
i-15:0 + i-17:0	65.2	64.1	53.7	61.0	64.7	75.1	58.6	66.1
a-15:0 + a-17:0	25.5	27.5	37.8	30.2	18.7	11.2	14.4	14.8
i-14:0 + i-16:0	2.8	3.5	5.5	3.9	9.3	3.9	10.3	7.8
n-14:0 + n-16:0	1.9	1.2	3.1	2.1	6.8	1.8	1.2	3.3
Avg chain length	15.5	15.3	15.7	15.5	16.1	15.9	16.1	16.0