

# Mouse Hepatitis Virus S RNA Sequence Reveals that Nonstructural Proteins ns4 and ns5a Are Not Essential for Murine Coronavirus Replication

KYOKO YOKOMORI AND MICHAEL M. C. LAI\*

*Howard Hughes Medical Institute and Department of Microbiology, University of Southern California School of Medicine, Los Angeles, California 90033-1054*

Received 14 May 1991/Accepted 4 July 1991

**Genes 4 and 5 of mouse hepatitis virus (MHV) are known to encode nonstructural proteins ns4, ns5a, and ns5b, whose function is unknown. In this study, we demonstrated that one of the MHV strains, MHV-S, did not synthesize mRNA 4 and made a smaller mRNA 5. Sequence analysis showed that the transcription initiation site for gene 4 of MHV-S was mutated from the consensus UC UAAAC to UU UAAAC, consistent with the idea that mutations in this region abolish mRNA synthesis. Furthermore, within gene 5 there were deletions totaling 307 nucleotides which deleted almost all of open reading frame 5a, but preserved open reading frame 5b of gene 5. Comparison of the growth of MHV-S with other MHV strains in DBT cells revealed no significant growth defect in MHV-S. These results suggest that ns4 and ns5a are not essential for viral replication in tissue culture cells, and thus join gene 2 and the hemagglutinin-esterase (HE) gene as nonessential viral genes in MHV.**

Mouse hepatitis virus (MHV), a member of the family *Coronaviridae*, contains a single-stranded, positive-sense RNA genome of 31 kb (8, 17). The viral particle is enveloped and consists of three or four structural proteins (6). The spike (S) protein is a 180-kDa envelope glycoprotein which is frequently cleaved into two 90-kDa species and is apparently responsible for receptor binding on target cells (3, 26, 29). The membrane (M) protein is a membrane-spanning glycoprotein of 23 kDa, and the nucleocapsid (N) protein is a phosphorylated RNA-binding protein of 55 to 60 kDa (6). While these three proteins are present in every MHV isolate, a fourth structural protein, the hemagglutinin-esterase (HE) protein, is present only in certain MHV strains (23, 27). The HE protein is an envelope glycoprotein of 65 kDa whose function is unknown. These proteins are translated from virus-specific mRNAs which show a nested-set structure (7). Most MHV strains synthesize seven or eight mRNAs with common 3' termini but unique 5' regions (7, 11). Although these mRNAs are structurally polycistronic, only their 5' unique regions are translated (10).

In addition to the four structural proteins, there are at least four genes that encode nonstructural (ns) proteins. The order of genes on the MHV genome is, from 5' to 3', ns1-ns2-HE-S-ns4-ns5a-5b-M-N (6), and the genes are encoded by mRNAs 1, 2, 2-1, 3, 4, 5 (for both 5a and 5b), 6 and 7, respectively. The gene for ns1, the biggest gene (22 kb) on the MHV genome, presumably encodes a polyprotein of 800 kDa which includes RNA polymerase, protease, and probably some additional activities (1, 8), although no distinct ns1 proteins have been detected in MHV-infected cells. ns2 protein is a 30-kDa cytoplasmic protein which, on the basis of its amino acid sequence, is postulated to have a nucleotide-binding capacity (12). ns4 protein has been detected as a 15-kDa protein in infected cells (4, 25), while the ns5 gene has two open reading frames (ORFs), 5a and 5b (2, 24). In the JHM and A59 strains of MHV, ORFs 5a and 5b have the capacity to encode a 13-kDa basic protein and a 10-kDa

neutral protein with a hydrophobic N terminus, respectively (2, 9, 24). Both ORFs 5a and 5b are encoded from a single mRNA 5; the downstream ORF, 5b, was translated preferentially in vitro, suggesting the utilization of an internal AUG in this mRNA (2, 9).

Although the function of the ns proteins is not clear, two recent studies have suggested that some of these proteins are not essential for coronavirus replication. Schwarz et al. isolated a variant of the JHM strain of MHV which, despite a defective ns2 gene, had normal growth properties, suggesting that the protein was not necessary for viral replication in tissue culture (22). Furthermore, porcine respiratory coronavirus apparently lacks protein ns3a (equivalent to ns4 of MHV in genomic location) (19).

In this study, we examined a strain of MHV, MHV-S, and found that gene 4 of this virus was not expressed and gene 5a was deleted. However, the growth of MHV-S in cell culture was not significantly different from that of the other MHV strains tested. These data suggest that ns4 and ns5a are not essential for viral replication in tissue culture and add to the growing list of nonessential viral genes in the coronavirus genome.

MHV-S, which was initially isolated from mice dying of acute hepatitis (20), was a generous gift from Paul Masters of Wadsworth Center for Laboratories and Research, New York State Department of Health. A59, a prototype MHV, was used for comparison and was derived from a source described previously (28). Virus was propagated in a murine astrocytoma cell line, DBT (5), throughout the study.

We first examined the intracellular virus-specific RNA patterns of MHV-S and A59 (Fig. 1). <sup>32</sup>P-labeled RNA was extracted from infected DBT cells, denatured by glyoxal treatment, and analyzed by agarose gel electrophoresis as described previously (15). Although seven mRNAs were detected in both MHV-S- and A59-infected cells, MHV-S synthesized neither mRNA 4 nor mRNA 5. Instead, a novel mRNA (Fig. 1, arrowhead), with a size intermediate between those of mRNAs 5 and 6, was observed. Furthermore, mRNAs 1, 2, and 3 of MHV-S migrated slightly faster than the corresponding mRNAs of A59 but their relative migra-

\* Corresponding author.




FIG. 1. Intracellular virus-specific RNA from A59- or MHV-S-infected DBT cells. DBT cells were infected with either A59 or MHV-S at a multiplicity of infection of 5 to 10. The medium was changed to phosphate-free minimum essential medium containing 1% dialyzed fetal calf serum and 2.5  $\mu$ g of actinomycin D (Sigma) per ml at 3.5 h after infection. Two hours later, 0.6 mCi of  $^{32}$ P<sub>i</sub> (ICN) was added and cells were labeled for 2 h. RNA was then extracted from cytoplasm as described previously (15). The numbers indicate each viral mRNA (6). The smaller mRNA 5 in MHV-S is indicated by the arrowhead.

tion rates remained unchanged. In contrast, mRNAs 6 and 7 of both viruses had the same migration rate. As demonstrated previously, mRNA 2-1, which encodes HE, was not detected in A59-infected cells, probably as a result of a mutation in its transcription initiation signal (12, 23, 27). The presence of several minor RNA species between mRNAs 3 and 5 was also detected. These RNAs varied with cell types and were most likely minor viral transcripts and rRNAs, since most of them were not detectable after poly(A) purification (unpublished observation). These minor RNAs have also been observed in cells infected with other strains of MHV (13, 15).

Since MHV mRNAs possess a nested-set structure (6, 7, 11), the patterns of viral mRNA detected (Fig. 1) suggested that a deletion occurred within gene 5 and possibly also in other genes upstream of gene 5 in the MHV-S genome. To identify the possible site(s) of deletion, we first examined the sizes of the intracellular viral proteins. MHV-S-infected cells were labeled with [ $^{35}$ S]methionine and immunoprecipitated with anti-MHV polyclonal antibody as described previously (27, 28). Figure 2 shows that the S and M proteins of MHV-S had the same size as those of A59. Although the N protein of A59 migrated faster than that of MHV-S, there was no difference in N gene size between these two viruses (18).




FIG. 2. Intracellular virus-specific proteins from MHV-S- and A59-infected cells. DBT cells were infected with A59 or MHV-S at a multiplicity of infection of 5 to 10 or incubated with medium only as a control (mock infection). Cells were starved for methionine for 30 min and then labeled with 50  $\mu$ Ci of [ $^{35}$ S]methionine (Translabel; 1,193 Ci/mmol; ICN) per ml for 20 min at 7 to 8 h after infection. Lysates were prepared as described previously (28), immunoprecipitated with polyclonal antibodies made against MHV (23, 28), and analyzed by electrophoresis in a 7.5 to 15% gradient polyacrylamide gel containing 0.1% sodium dodecyl sulfate. Lane M contained  $^{14}$ C-labeled molecular weight markers (BRL), whose sizes ( $10^3$ ) are indicated on the left.

These results suggested that the possible deletion was localized in the genes encoding nonstructural proteins, more likely genes 4 and 5. As predicted, the HE protein was synthesized only in MHV-S-infected cells (27).

To determine the sites of deletions within gene 4 or 5, cDNA clones corresponding to these genes were obtained after amplification by polymerase chain reaction on the total intracellular RNA of MHV-S-infected cells by using two primers: one that binds to the 5' unique region of gene 7, no. 327 (5'-GCATTTTCTTGCCAGAAC-3'), and one that binds to either the leader sequence, no. 78 (5'-AGCTTTA CGTACCCTCTCTACTCTAAAACCTTTGTAGTTT-3'), or the sequence near the 3' end of gene 3, no. 347 (5'-GGACACCAGGACAGTATTGT-3'). The sequence used to synthesize primer 347 was obtained by direct RNA sequencing of MHV-S RNA (data not shown) by using various primers complementary to the known gene 3 sequence of JHM (21). The PCR protocol employed has been described previously (14, 16, 27). By this approach, cDNA clones covering 800 nucleotides of the 5' unique region of mRNA 6, 1.1 kb of the 5' unique region plus some downstream



in DBT cells (data not shown). Furthermore, the total amounts and peaks of synthesis of viral RNA and proteins and virus yields were similar between MHV-S and most of the other MHV strains (Fig. 1 and 2). These results suggest that ns4 and ns5a are not essential for MHV replication, at least in DBT cell culture. Different MHV strains did show minor differences in their growth properties, but none of these differences could be correlated with the presence or absence of ns gene expression.

These results add to the growing list of nonessential genes in the MHV genome. So far, ns2 (22), ns4, ns5a, and a structural protein, HE (12, 27), have been demonstrated not to be essential for MHV replication. However, these proteins may contribute to the biology or pathogenicity of the virus under certain situations. Elucidation of these functions will need additional studies.

**Nucleotide sequence accession number.** The nucleotide sequence reported here has been deposited in the EMBL-GenBank data base under accession no. M64835.

We thank Tom McNaughton for editorial help.

This work was supported by U.S. Public Health Service research grants AI 19244 and NS 18146. K.Y. was supported partially by a Fieger predoctoral fellowship from Norris Cancer Center, University of Southern California. M.M.C.L. is an Investigator of Howard Hughes Medical Institute.

#### REFERENCES

- Baker, S. C., C.-K. Shieh, L. H. Soe, M.-F. Chang, D. M. Vannier, and M. M. C. Lai. 1989. Identification of a domain required for the autoproteolytic cleavage of murine coronavirus gene A polypeptide. *J. Virol.* **63**:3693-3699.
- Budzilowicz, C. J., and S. R. Weiss. 1987. *In vitro* synthesis of two polypeptides from a nonstructural gene of coronavirus mouse hepatitis virus strain A59. *Virology* **157**:509-515.
- Collins, A. R., R. L. Knobler, H. Powell, and M. J. Buchmeier. 1982. Monoclonal antibodies to murine hepatitis virus-4 (strain JHM) define the viral glycoprotein responsible for attachment and cell-cell fusion. *Virology* **119**:358-371.
- Ebner, D., T. Raabe, and S. G. Siddell. 1988. Identification of the coronavirus MHV-JHM mRNA 4 product. *J. Gen. Virol.* **69**:1041-1050.
- Hirano, N., K. Fujiwara, S. Hino, and M. Matsumoto. 1974. Replication and plaque formation of mouse hepatitis virus (MHV-2) in mouse cell line DBT culture. *Arch. Gesamte Virusforsch.* **44**:298-302.
- Lai, M. M. C. 1990. Coronavirus: organization, replication and expression of genome. *Annu. Rev. Microbiol.* **44**:303-333.
- Lai, M. M. C., P. R. Brayton, R. C. Armen, C. D. Patton, C. Pugh, and S. A. Stohlman. 1981. Mouse hepatitis virus A59: mRNA structure and genetic localization of the sequence divergence from hepatotropic strain MHV-3. *J. Virol.* **39**:823-834.
- Lee, H.-J., C.-K. Shieh, A. E. Gorbalenya, E. V. Koonin, N. La Monica, J. Tuler, A. Bagdzhadzhyan, and M. M. C. Lai. 1991. The complete sequence (22 kilobases) of murine coronavirus gene 1 encoding the putative proteases and RNA polymerase. *Virology* **180**:567-582.
- Leibowitz, J. L., S. Perlman, G. Weinstock, J. R. de Vries, C. Budzilowicz, J. M. Weissman, and S. Weiss. 1988. Detection of a murine coronavirus non-structural protein encoded in a downstream open reading frame. *Virology* **164**:156-164.
- Leibowitz, J. L., S. R. Weiss, E. Paavola, and C. W. Bond. 1982. Cell-free translation of murine coronavirus RNA. *J. Virol.* **43**:905-913.
- Leibowitz, J. L., K. C. Wilhelmsen, and C. W. Bond. 1981. The virus specific intracellular RNA species of two murine coronaviruses: MHV-A59 and MHV-JHM. *Virology* **114**:39-51.
- Luytjes, W., P. J. Bredenbeek, A. F. H. Noten, M. C. Horzinek, and W. J. M. Spaan. 1988. Sequence of mouse hepatitis virus A59 mRNAs: indications for RNA recombination between coronaviruses and influenza C virus. *Virology* **166**:415-422.
- Makino, S., and M. M. C. Lai. 1989. High-frequency leader sequence switching during coronavirus defective interfering RNA replication. *J. Virol.* **63**:5285-5292.
- Makino, S., L. H. Soe, C.-K. Shieh, and M. M. C. Lai. 1988. Discontinuous transcription generates heterogeneity at the leader fusion sites of coronavirus mRNA. *J. Virol.* **62**:3870-3873.
- Makino, S., F. Taguchi, H. Hirano, and K. Fujiwara. 1984. Analysis of genomic and intracellular viral RNAs of small plaque mutants of mouse hepatitis virus, JHM strain. *Virology* **139**:138-151.
- Makino, S., K. Yokomori, and M. M. C. Lai. 1990. Analysis of efficiently packaged defective interfering RNAs of murine coronavirus: localization of a possible RNA-packaging signal. *J. Virol.* **64**:6045-6053.
- Pachuk, C. J., P. J. Bredenbeek, P. W. Zoltick, W. J. M. Spaan, and S. R. Weiss. 1989. Molecular cloning of the gene encoding the putative polymerase of mouse hepatitis coronavirus strain A59. *Virology* **171**:141-148.
- Parker, M. M., and P. S. Masters. 1990. Sequence comparison of the N genes of five strains of the coronavirus mouse hepatitis virus suggests a three domain structure for the nucleocapsid protein. *Virology* **179**:463-468.
- Rasschaert, D., M. Duarte, and H. Laude. 1990. Porcine respiratory coronavirus differs from transmissible gastroenteritis virus by a few genomic deletions. *J. Gen. Virol.* **71**:2599-2607.
- Rowe, W., J. Hartley, and W. Capps. 1963. Mouse hepatitis virus infection as a highly contagious, prevalent enteric infection of mice. *Proc. Soc. Exp. Biol. Med.* **112**:161-165.
- Schmidt, I., M. Skinner, and S. Siddell. 1987. Nucleotide sequence of the gene encoding the surface projection glycoprotein of coronavirus MHV-JHM. *J. Gen. Virol.* **68**:47-56.
- Schwarz, B., E. Routledge, and S. G. Siddell. 1990. Murine coronavirus nonstructural protein ns2 is not essential for virus replication in transformed cells. *J. Virol.* **64**:4784-4791.
- Shieh, C.-K., H.-J. Lee, K. Yokomori, N. La Monica, S. Makino, and M. M. C. Lai. 1989. Identification of a new transcriptional initiation site and the corresponding functional gene 2b in the murine coronavirus RNA genome. *J. Virol.* **63**:3729-3736.
- Skinner, M. A., D. Ebner, and S. G. Siddell. 1985. Coronavirus MHV-JHM mRNA 5 has a sequence arrangement which potentially allows translation of a second, downstream open reading frame. *J. Gen. Virol.* **66**:581-592.
- Skinner, M. A., and S. G. Siddell. 1985. Coding sequence of coronavirus MHV-JHM mRNA 4. *J. Gen. Virol.* **66**:593-596.
- Sturman, L. S., C. S. Ricard, and K. V. Holmes. 1985. Proteolytic cleavage of the E2 glycoprotein of murine coronavirus: activation of cell fusing activity of virions by trypsin and separation of two different 90K cleavage fragments. *J. Virol.* **56**:904-911.
- Yokomori, K., L. R. Banner, and M. M. C. Lai. 1991. Heterogeneity of gene expression of the hemagglutinin-esterase (HE) protein of murine coronaviruses. *Virology* **183**:647-657.
- Yokomori, K., N. La Monica, S. Makino, C.-K. Shieh, and M. M. C. Lai. 1989. Biosynthesis, structure, and biological activities of envelope protein gp65 of murine coronavirus. *Virology* **173**:683-691.
- Yoshikura, H., and S. Tejima. 1981. Role of protease in mouse hepatitis virus-induced cell fusion. *Virology* **113**:503-511.