

# Genetic and Regulatory Aspects of Methionine Biosynthesis in *Saccharomyces cerevisiae*

H. CHEREST, F. EICHLER, AND H. DE ROBICHON-SZULMAJSTER

Laboratoire d'Enzymologie, Centre National de la Recherche Scientifique, 91-Gif-sur-Yvette, France

Received for publication 2 August 1968

Methionine biosynthesis and regulation of four enzymatic steps involved in this pathway were studied in *Saccharomyces cerevisiae*, in relation to genes concerned with resistance to ethionine (*eth*<sub>1</sub> and *eth*<sub>2</sub>). Data presented in this paper and others favor a scheme which excludes cystathionine as an obligatory intermediate. Kinetic data are presented for homocysteine synthetase [ $K_m$ (*O*-acetyl-L-homoserine) =  $7 \times 10^{-3}$  M;  $K_i$ (L-methionine) =  $1.9 \times 10^{-3}$  M]. Enzymes catalyzing steps 3, 4, 5, and 9 were repressible by methionine. Enzyme 4 (homoserine-*O*-transacetylase) and enzyme 9 (homocysteine synthetase) were simultaneously derepressed in strains carrying the mutant allele *eth*<sub>2</sub><sup>r</sup>. Studies on diploid strains confirmed the dominance of the *eth*<sub>2</sub><sup>s</sup> allele over *eth*<sub>2</sub><sup>r</sup>. Regulation of enzyme 3 (homoserine dehydrogenase) and enzyme 5 (adenosine triphosphate sulfurylase) is not modified by the allele *eth*<sub>2</sub><sup>r</sup>. The other gene *eth*<sub>1</sub> did not appear to participate in regulation of these four steps. Gene enzyme relationship was determined for three of the four steps studied (steps 3, 4, and 9). The structural genes concerned with the steps which are under the control of *eth*<sub>2</sub> (*met*<sub>3</sub>: enzyme 9 and *met*<sub>4</sub>: enzyme 4) segregate independently, and are unlinked to *eth*<sub>2</sub>. These results are compatible with the idea that the gene *eth*<sub>2</sub> is responsible for the synthesis of a pleiotropic methionine repressor and suggest the existence of at least two different methionine repressors in *S. cerevisiae*. Implications of these findings in general regulatory mechanisms have been discussed.

Methionine biosynthesis in *Saccharomyces cerevisiae* can be divided in three parts: (i) synthesis of the four-carbon skeleton; (ii) sulfur assimilation; (iii) synthesis and attachment of the methyl groups. Regulation by end-product inhibition and repression occurs at different steps of this pathway. This paper is concerned only with biochemical and regulatory aspects of parts (i) and (ii).

The four-carbon skeleton of methionine and threonine originates from aspartate. Homoserine dehydrogenase [L-homoserine:nicotinamide adenine dinucleotide (NAD) oxidoreductase; EC 1.1.1.3], the enzyme catalyzing the last common step leading to these two end products, is repressed by methionine (65% repression in the wild-type strains); both methionine and threonine participate in the feedback inhibition of this enzyme (12). Aspartate kinase (L-aspartate 4-phosphotransferase; EC 2.7.2.4), catalyzing the first step in threonine-methionine biosynthesis, is repressed by threonine; a mutation altering this repression does not affect the repressibility of homoserine dehydrogenase (27).

From homoserine, the biosynthesis of methionine starts with an acetylation reaction leading to

*O*-acetyl-homoserine which is apparently the active substrate for the entry of sulfur into the pathway. The enzyme that catalyzes homoserine *O*-acetylation (homoserine-*O*-transacetylase) is under the control of methionine for its synthesis and of *S*-adenosyl-methionine for its activity (25).

The sulfur assimilation originates with sulfate activation (22). In yeast, this reaction is subject to feedback inhibition by sulfide. In addition, adenosine 5'-phosphosulfate (APS) and 3'-phospho-adenosine-5'-phosphosulfate (PAPS), which are the intermediary compounds leading to sulfite, are also inhibitors. The synthesis of the first enzyme, adenosine triphosphate sulfurylase (ATP: sulfate adenylyltransferase; EC 2.7.7.4), is repressed by methionine and derepressed by cysteine (35). On the other hand, the enzyme sulfite reductase [hydrogen sulfide:nicotinamide adenine dinucleotide phosphate (NADP) oxidoreductase; EC 1.8.1.2] is repressed in cysteine-grown cells (35). Consequently, sulfide synthesis appears to be under double, and somehow complementary, control of the two sulfur-containing amino acids, methionine and cysteine.

Direct sulfide utilization occurred by condensation of homoserine and H<sub>2</sub>S in *Neurospora* (37)



Homocysteine formed was estimated according to Kredich and Tomkins (13).

Protein estimation was carried out according to Lowry et al. (16), with bovine serum albumin as reference.

**Genetic techniques.** Sporulation of diploids was induced according to McClary et al. (17). Ascospores were isolated according to Johnston and Mortimer (11).

**Chemicals.** *O*-acetyl-DL-homoserine was synthesized for us by M. Cherest according to Sakami and Toennies (30).

DL-C<sub>14</sub>-<sup>14</sup>C-homoserine was purchased from the C.E.A., France.

Inorganic pyrophosphatase was obtained from Worthington Biochemical Corp., Freehold, N.J.

Aspartic semialdehyde was prepared according to Black and Wright (1) and titered with purified yeast homoserine dehydrogenase (12).

## RESULTS

**Homoserine *O* transacetylase.** Earlier studies on methionine auxotrophs devoided of homoserine-*O*-transacetylase activity have shown that homoserine-*O*-acetylation is the first step in methionine biosynthesis in *S. cerevisiae* (25). These studies also led to the following conclusions: (i) that activity of this enzyme in crude extracts is subject to feedback inhibition, not by free methionine but by its activated product *S*-adenosyl-methionine; (ii) that the synthesis of homoserine-*O*-transacetylase is repressed by methionine (70% repression by DL-methionine  $2 \times 10^{-3}$  M). In addition, we pointed out the possible role of *eth2<sup>r</sup>* in this repression. Further investigation on the role of this gene is presented in this paper.

After growth in the presence of DL-methionine ( $2 \times 10^{-3}$  M), the wild-type strains 4094-B (*eth1<sup>s</sup>, eth2<sup>s</sup>*) synthesized very little acetyl-homoserine (Fig. 1, spot 2); moreover, the compound derived from it (spot 3) was absent; on the contrary, the two other strains CH82-7D (*eth1<sup>r</sup>, eth2<sup>r</sup>*) and CC30-ID (*eth1<sup>s</sup>, eth2<sup>r</sup>*) synthesized considerable amounts of these two compounds.

A quantitative estimation of homoserine-*O*-transacetylase activity is summarized in Table 1. All strains carrying the wild-type allele (*eth2<sup>s</sup>*) were repressible by methionine, whereas the strains carrying the mutated recessive allele (*eth2<sup>r</sup>*) lost this repressibility. The results show unequivocally that the alleles *eth1<sup>r</sup>* or *eth1<sup>s</sup>* do not modify this regulation. In either case, *S*-adenosyl-methionine inhibition is maintained. Furthermore, Table 1 shows the absence of homoserine-*O*-transacetylase in mutant strains D6 and CC92-8D (gene *met<sub>a</sub>*).

**Homocysteine synthetase.** Direct synthesis of homocysteine has been shown to occur in *Neurospora* from homoserine and H<sub>2</sub>S by Wiebers and

TABLE 1. Homoserine-*O*-transacetylase activity<sup>a</sup> and repression in different strains of *S. cerevisiae*<sup>b</sup>

Strains	Genotypes	Growth conditions		Avg repression
		Minimal medium	Minimal medium + DL-methionine ( $2 \times 10^{-3}$ M)	
4094-B	<i>eth1<sup>s</sup> eth2<sup>s</sup></i>	0.381	0.125	66
CH82-7A	<i>eth1<sup>r</sup> eth2<sup>s</sup></i>	0.405	0.141	
CC30-1D	<i>eth1<sup>s</sup> eth2<sup>r</sup></i>	0.359	0.333	3
CH82-7D	<i>eth1<sup>r</sup> eth2<sup>r</sup></i>	0.311	0.316	
D6	? <i>eth2<sup>s</sup></i>		0-0.050	0
CC92-8D	<i>eth1<sup>r</sup> eth2<sup>s</sup></i>		0	

<sup>a</sup> Expressed as nanomoles per minute per milligram of protein.

<sup>b</sup> Strains are grown in the presence of minimal medium complemented with adenine and uracil. Their genotype is given only regarding the ethionine resistance alleles known to be present. Complete genotypes are given in Materials and Methods

TABLE 2. Homocysteine synthetase activity<sup>a</sup> and repression in different strains of *S. cerevisiae*<sup>b</sup>

Strains	Growth conditions		Avg repression
	Minimal medium	Minimal medium DL-methionine ( $2 \times 10^{-3}$ M)	
			%
<b>Haploids</b>			
4094-B( <i>eth1<sup>s</sup> eth2<sup>s</sup></i> )	444	62	87
CH82-7A( <i>eth1<sup>r</sup> eth2<sup>s</sup></i> )	425	74	
D6(? <i>eth2<sup>s</sup></i> )	162	0	
CC30-1D( <i>eth1<sup>s</sup> eth2<sup>r</sup></i> )	416	236	31
CH82-7D( <i>eth1<sup>r</sup> eth2<sup>r</sup></i> )	525	273	
CC92-8D( <i>eth1<sup>r</sup> eth2<sup>r</sup></i> )	214	172	
<b>Heterozygous diploids</b>			
CC18-CC53 ( <i>eth2<sup>s</sup>/eth2<sup>r</sup></i> )	343	67	81
<b>Homozygous diploids</b>			
CC17( <i>eth2<sup>r</sup>/eth2<sup>r</sup></i> )	248	153	42

<sup>a</sup> Expressed as nanomoles per minute per milligram of protein.

<sup>b</sup> See legend of Table 1. The mutant strains D6 and CC92-8D are grown in the minimal medium described, supplemented with *O*-acetyl-DL-homoserine  $10^{-3}$  M. For heterozygous diploids, we have given the mean value obtained for two homologous diploids.




FIG. 1. Homoserine-*O*-transacetylase activity in different strains grown in a methionine-supplemented medium. Strains used are of the following genotypes: 4094-B:  $\alpha$ ,  $ade_2$ ,  $ura_1$ ,  $eth_1^+$ ,  $eth_2^+$ ; CH82-7D:  $\alpha$ ,  $ade_2$ ,  $ura_1$ ,  $eth_1^+$ ,  $eth_2^+$ ; CC30-1D:  $\alpha$ ,  $ade_2$ ,  $ura_1$ ,  $eth_1^+$ ,  $eth_2^+$ . Minimal medium was supplemented with DL-methionine  $2 \times 10^{-3}$  M. Spot identification: 1,  $^{14}\text{C}$ -4-DL-homoserine; 2, *O*-acetyl-DL-homoserine; 3, compound (ninhydrin-negative) derived from *O*-acetyl-homoserine; 4 and 5, radioactive impurities present in  $^{14}\text{C}$ -4-labeled-DL-homoserine. The assay and solvent system have been described (25).

Garner (37). However, in view of our previous finding of the occurrence of homoserine-*O*-transacetylation in *S. cerevisiae* we favored *O*-acetyl-homoserine as a candidate for direct homocysteine synthesis (25). More recently Wiebers and Garner (38) found that acyl derivatives of homoserine are better substrates than homoserine for homocysteine synthesis in *N. crassa*, *Escherichia coli*, and in commercial yeast (38).

We investigated the properties of this enzyme in our strains grown in synthetic or methionine-supplemented media (Table 2 and Fig. 2).

It can be first pointed out that the specific activities found for cells grown in synthetic medium were much higher than activities observed by previous authors for commercial cells (360 nmoles per min per mg of protein at 30 C). This could be

caused by the fact that enzymes obtained from commercial cells are often found at their repressed level. This was found to be the case for aspartate kinase, aspartic semialdehyde dehydrogenase, homoserine dehydrogenase, and threonine deaminase (de Robichon-Szulmajster, unpublished data).

It can be seen in Fig. 2A that the affinity of the enzyme for *O*-acetyl-L-homoserine ( $K_m = 7 \times 10^{-3}$  M) is of the same order of magnitude as the one already published (38). However, the enzyme of our wild-type strains exhibited a much higher affinity for methionine ( $K_i$  for L-methionine =  $1.9 \times 10^{-3}$  M) than that found by Wiebers and Garner (38; Fig. 2B, 2C). In addition, the inhibition caused by methionine seems, in our experiments, competitive with *O*-acetyl-homoserine.




FIG. 2. Kinetic studies of homocysteine synthetase from *S. cerevisiae*. (A) Variation of activity with *O*-acetyl-*L*-homoserine concentration in the absence (●) or in the presence (⊗) of *L*-methionine  $5 \times 10^{-3} \text{ M}$  [Lineweaver-Burk transformation (15)]. (B) Variation of activity ( $1/V$ ) with *L*-methionine concentration in the presence of *O*-acetyl-*L*-homoserine  $10^{-3} \text{ M}$ . Same enzyme preparation was used for A and B. (C) Variation of activity ( $1/V$ ) with *L*-methionine concentration in the presence of *O*-acetyl-*L*-homoserine  $4 \times 10^{-3} \text{ M}$  (●) and  $8 \times 10^{-3} \text{ M}$  (⊗).  $K_i$  values have been calculated according to Dixon and Webb (4).

Our results with homoserine-*O*-transacetylase regulation and the role of *eth*<sub>2</sub><sup>r</sup> prompted us to search for homocysteine synthetase repressibility.

Methionine was a very potent repressor of homocysteine synthetase (Table 2) in strains carrying the wild-type allele *eth*<sub>2</sub><sup>s</sup> (87% repression by DL-methionine  $2 \times 10^{-3} \text{ M}$ ). As in the case of homoserine-*O*-transacetylase, it appears that the presence of the mutated allele *eth*<sub>2</sub><sup>r</sup> strongly diminished this repression (31%), whereas the state of the *eth*<sub>1</sub> gene had no effect. In all cases, methionine inhibition has been found to be unmodified.

**Homoserine dehydrogenase.** It has already been shown that homoserine dehydrogenase synthesis is regulated by methionine (12). In view of our findings on the role of *eth*<sub>2</sub> gene in the repressibility of homoserine-*O*-transacetylase and homocysteine synthetase, we investigated the repressibility of homoserine dehydrogenase in the two extreme genotypes. Results have shown no difference in this regard (*eth*<sub>1</sub><sup>s</sup> *eth*<sub>2</sub><sup>s</sup>: 41 to 45%; *eth*<sub>1</sub><sup>r</sup> *eth*<sub>2</sub><sup>r</sup>: 35 to 52% repression by DL-methionine  $5 \times 10^{-3} \text{ M}$ ).

**ATP sulfurylase.** The first step in sulfur assimilation in yeast is repressed by methionine (35). Table 3 shows that repression of ATP-sulfurylase by DL-methionine ( $2 \times 10^{-3} \text{ M}$ ) was identical in all strains, regardless of the *eth* alleles present.

The specific activities found in these experiments were much higher than the values published by deVito and Dreyfuss (35). Again, strains used and culture conditions are quite different.

As for the other enzymes tested, inhibitory properties of ATP-sulfurylase (here by sodium sulfide) were identical in all cases.

**Dominance of the allele *eth*<sub>2</sub><sup>s</sup> on *eth*<sub>2</sub><sup>r</sup>.** Earlier genetic and physiological studies, based on methionine resistance-sensitivity of diploid strains carrying *eth*<sub>2</sub> alleles in homoallelic or heteroallelic combinations, have shown a complete dominance of *eth*<sub>2</sub><sup>s</sup> (24). Because of the knowledge gained on the regulation of homocysteine synthetase we were able to use repressibility of this enzyme as a criterion of dominance of *eth*<sub>2</sub> alleles.

The repression of homocysteine synthetase in heteroallelic strains (*eth*<sub>2</sub><sup>s</sup>/*eth*<sub>2</sub><sup>r</sup>) was of the same

TABLE 3. ATP-sulfurylase activity<sup>a</sup> and repression in different strains of *S. cerevisiae*<sup>b</sup>

Strains	Growth conditions		Avg repression
	Minimal medium	Minimal medium + DL-methionine ( $2 \times 10^{-3}$ M)	
Haploids			%
4094-B( <i>eth</i> <sub>1</sub> <sup>+</sup> <i>eth</i> <sub>2</sub> <sup>+</sup> )	148	15	93
CH82-7A( <i>eth</i> <sub>1</sub> <sup>+</sup> <i>eth</i> <sub>2</sub> <sup>+</sup> )	81	0	
D6(? <i>eth</i> <sub>2</sub> <sup>+</sup> )	130	12	
CC30-1D( <i>eth</i> <sub>1</sub> <sup>+</sup> <i>eth</i> <sub>2</sub> <sup>+</sup> )	181	12	81
CH82-7D( <i>eth</i> <sub>1</sub> <sup>+</sup> <i>eth</i> <sub>2</sub> <sup>+</sup> )	246	37	
CC92-8D( <i>eth</i> <sub>1</sub> <sup>+</sup> <i>eth</i> <sub>2</sub> <sup>+</sup> )	230	81	
Heterozygous diploids			
CC18-CC53 ( <i>eth</i> <sub>2</sub> <sup>+</sup> / <i>eth</i> <sub>2</sub> <sup>+</sup> )	131	5	96
Homozygous diploids			
CC17( <i>eth</i> <sub>2</sub> <sup>+</sup> / <i>eth</i> <sub>2</sub> <sup>+</sup> )	168	40	75, 5

<sup>a</sup> Expressed as nanomoles per minute per milligram of protein.

<sup>b</sup> See legend of Table 2.

order of magnitude as that found in haploid *eth*<sub>2</sub><sup>+</sup> strains (Table 2). In addition, in the homoallelic diploid *eth*<sub>2</sub><sup>+</sup>/*eth*<sub>2</sub><sup>+</sup>, the repression was comparable to that found in haploid *eth*<sub>2</sub><sup>+</sup> strains. These data confirm our previous conclusion on the dominance of *eth*<sub>2</sub><sup>+</sup> over *eth*<sub>2</sub><sup>-</sup>.

The results of a control experiment showed that the presence of *eth*<sub>2</sub><sup>+</sup> did not affect the repressibility of ATP-sulfurylase in haploid or diploid strains (Table 3).

Enzymatic study of mutant strains *met*<sub>8</sub> and *met*<sub>10</sub>. In a survey of methionine auxotrophs, we estimated the activity of ATP-sulfurylase and homocysteine synthetase (Table 4). It appears

that the strain carrying the *met*<sub>8</sub> allele, which has been already mapped on chromosome II (19), is deficient in homocysteine synthetase. This result is compatible with the idea that the marker *met*<sub>8</sub> represents mutation in the structural gene for homocysteine synthetase. The strain carrying the *met*<sub>10</sub> allele presents rather low values for both enzymes when compared with reference strains grown in the same conditions. The exact nature of this mutant remains then to be determined.

**Genetic analysis.** We made a genetic analysis of appropriate crosses to investigate further the relationship between the regulatory gene *eth*<sub>2</sub> and the two structural genes which are under its control (*met*<sub>8</sub> and *met*<sub>10</sub>). The two crosses analyzed thus far were subjected to tetrad analysis. They are the following: CC125:CC92-8D × EY9 (*met*<sub>8</sub>; *eth*<sub>2</sub><sup>+</sup> × *met*<sub>8</sub>; *eth*<sub>2</sub><sup>+</sup>) CC92:D6 × CH82-9C (*met*<sub>8</sub>; *eth*<sub>2</sub><sup>+</sup> × *eth*<sub>2</sub><sup>+</sup>).

The cross involving *met*<sub>8</sub> and *met*<sub>10</sub> (CC125) shows, for 20 tetrads: 6 parental ditypes, 3 non-parental ditypes, and 11 tetratypes. In spite of the slight excess of parental ditypes over the non-parental ditypes, which could be caused by the small number of tetrads analyzed in this cross, the large percentage of tetratypes (55% of the total) is in favor of genetic independence and, in any case, excludes any close linkage between the two genes.

The cross involving *met*<sub>8</sub> and *eth*<sub>2</sub><sup>+</sup> (CC92) produced 88 *met*<sup>-</sup> and 88 *met*<sup>+</sup>. Since sensitivity to ethionine is never manifested in the presence of methionine (18), the distribution of the *eth*<sub>2</sub><sup>+</sup> allele could not be tested in the first class. Among the second class, we obtained 44 ethionine-resistant spores. This segregation is typical of two independent characters.

Segregation of the alleles *met*<sub>8</sub> and *eth*<sub>2</sub><sup>+</sup> was considered thus far only in the cross described above (CC125). Since this cross involved two independent methionine auxotrophic characters,

TABLE 4. Enzymatic activities<sup>a</sup> in various methionine auxotrophs<sup>b</sup>

Strains	Genotypes	Growth conditions (minimal medium + DL-homocysteine, $10^{-3}$ M)			
		ATP-sulfurylase		Homocysteine synthetase	
		Avg	Extreme values	Avg	Extreme values
EY 9	<i>met</i> <sub>8</sub> MET <sub>10</sub>	109	82-140	4	0-9
X 1266/1C	MET <sub>8</sub> <i>met</i> <sub>10</sub>	29	6-42	100	75-171
Reference	MET <sub>8</sub> MET <sub>10</sub>	121	62-240	279	109-450

<sup>a</sup> Expressed as nanomoles per minute per milligram of protein.

<sup>b</sup> Complete genotypes are given in Materials and Methods. Since the true parental strains of these two mutants were not in our possession, we have taken as reference activities the average of determinations made on related methionine auxotrophs which are blocked at other steps in the pathway.

only 17 methionine prototrophs were recovered. These yielded 7 ethionine-sensitive and 10 ethionine-resistant spores. Again, this segregation is in favor of independence between the two characters.

### DISCUSSION

The results presented in this paper concern two different aspects of methionine biosynthesis in *S. cerevisiae*. One is the direct synthesis of homocysteine, the other is the more general problem of relationship between structural and regulatory genes in a multichromosomal organism like *S. cerevisiae*.

Homocysteine synthetase has already been described in yeast. We found this enzyme at a very high level in our wild-type strains grown in minimal medium. In addition, a completely methionineless mutant, devoid of homocysteine synthetase, was found and characterized. The activity of this enzyme is under strict control by methionine (37, 38). Furthermore, the gene *eth<sub>2</sub>*, which affects repressibility of homoserine-*O*-transacetylase, also affects repressibility of homocysteine synthetase.

Flavin and Slaughter (7) have reported the ability of *Salmonella* cystathionine  $\gamma$ -synthetase to carry out additional reactions, one which is direct synthesis of homocysteine from *O*-succinyl-homoserine and H<sub>2</sub>S.

However, recent data from Sorsoli et al. (32) indicate that, although cystathionine is actively transported and the  $\beta$ -cleavage enzyme can be demonstrated in *S. cerevisiae*, the strains studied are unable to use exogenous cystathionine for growth. Giovanelli and Mudd (9) have reported that, in crude extracts of spinach, the conversion of *O*-acetyl-homoserine to *S*-adenosyl-homocysteine is at least 10 times more efficient than its conversion to cystathionine. In addition, the same investigators (10) show that cystathionine fails to decrease the label of homocysteine synthesized from <sup>35</sup>SO<sub>4</sub> and that cystathionine is not labeled in the process. Utilization of cystathionine <sup>14</sup>C-labeled in the four-carbon chain has shown that this chain is not incorporated in methionine by *Neurospora* (36).

These results, in addition to (i) the existence of a very active homocysteine synthetase, (ii) the regulatory properties of this enzyme, and (iii) the existence of the mutant EY 9 already described, lead to the obligatory participation of homocysteine synthetase as a step in methionine biosynthesis in *S. cerevisiae* and make it unnecessary to implicate cystathionine as an obligatory intermediate in this pathway.

These considerations led us to favor the following scheme of methionine biosynthesis in *S.*




FIG. 3. Metabolic and regulatory scheme for methionine biosynthesis in *S. cerevisiae*. Step numbers correspond to the following enzymes: 1, aspartate kinase (*L*-aspartate 4-phosphotransferase; EC 2.7.2.4). 2, ASA dehydrogenase [*L*-aspartate- $\beta$ -semialdehyde: NADP oxidoreductase (phosphorylating); EC 1.2.1.11]. 3, HS-dehydrogenase (*L*-homoserine: NAD oxidoreductase; EC 1.1.1.3). 4, *L*-Homoserine-*O*-transacetylase (Nagai and Flavin, 20). 5, ATP-sulfurylase (ATP: sulfate adenylate transferase; EC 2.7.7.4). 6, APS-kinase (ATP: adenylylsulfate-3'-phosphotransferase; EC 2.7.1.25). 7, PAPS-reductase (Pasternak et al., 21). 8, Sulfite reductase (hydrogen sulphide: NADP oxidoreductase; EC 1.8.1.2). 9, Homocysteine synthetase (Wiebers and Garner, 38). Relevant structural genes are connected to reactions by broken arrows. Compounds are abbreviated as in the scheme given in the introduction. Large empty arrows indicate enzymatic steps repressed by methionine.

*cerevisiae* which excludes, thus far, cystathionine as an obligatory intermediate.

Although a few examples of clustered structural genes have been reported (5, 6, 14, 34) in *S. cerevisiae*, it is commonly found that genes controlling sequential steps in a metabolic pathway are located on different chromosomes (19). This is the case for the threonine-methionine biosynthetic pathway studied in this laboratory (23, 28).

Data obtained by de Vito and Dreyfuss (35) and from this laboratory (12, 25) have shown that the synthesis of four enzymes involved in methionine biosynthesis in *S. cerevisiae* responds to exogenous methionine (indicated by large arrow on Fig. 3). Results presented in this paper show that the regulation of two of these enzymes (homoserine-*O*-transacetylase, step 4, and homocysteine synthetase, step 9) is under the control of the gene *eth<sub>2</sub>*, whereas regulation of the two other

enzymes (homoserine dehydrogenase, step 3, and ATP-sulfurylase, step 5) is not.

Previous genetic analysis has shown that ethionine resistance conferred in haploid strains by the mutant allele  $eth_2^s$  is recessive in the heterozygous diploid,  $eth_2^s/eth_2^r$ . Present data confirm the dominance of the  $eth_2^s$  allele when characterized by assay of homocysteine synthetase repressibility on various diploid strains. In addition, it is shown that  $eth_2$  segregates independently of the gene  $met_a$  and of the gene  $met_s$ , and that  $met_a$  and  $met_s$  are unlinked as well.

These findings suggest that the  $eth_2$  gene may be involved in the synthesis of a pleiotropic methionine repressor on two unlinked structural genes  $met_a$  and  $met_s$ . The two other enzymes, whose synthesis is repressible by exogenous methionine but is nevertheless unaffected by  $eth_2$ , must then remain under the control of another (or other) methionine repressor(s).

Different types of repressor formed responding to the presence of a single amino acid can be postulated. They could differ either by the nature of the apo-repressor or by the exact nature of the corepressor: free amino acid, derivatives, and activated forms. In the case of methionine, two activated forms can be considered: the methionyl-transfer ribonucleic acid and the sulfonium derivative, *S*-adenosyl-methionine (SAM). The latter is accumulated in the presence of exogenous methionine (31), whereas the free methionine pool remains, in contrast, remarkably constant under these conditions (H. Cherest, unpublished data). It can be recalled that homoserine-*O*-transacetylase is inhibited by SAM and is insensitive to free methionine. We suggested that, by analogy, the corepressor of the gene  $met_a$  might also be SAM (25). Homocysteine synthetase, on the other hand, is very sensitive to inhibition by free methionine. The quantitative difference in the derepression effect on these two enzymes, in response to  $eth_2^r$ , could be due to a slight difference between the two operator-like structures of the two structural genes or to a slight difference in the nature of the corepressors involved.

#### ACKNOWLEDGMENTS

This investigation was supported by grants from the D.G.R.S.T. (66-00-140) and from the C.E.A., France.

#### LITERATURE CITED

- Black, S., and N. G. Wright. 1955. Aspartic  $\beta$ -semi-aldehyde dehydrogenase and aspartic  $\beta$ -semi-aldehyde. *J. Biol. Chem.* 213:39-50.
- Cherest, H., and H. de Robichon-Szulmajster. 1966. Résistance à l'éthionine chez *S. cerevisiae*. I. Etude génétique. *Genetics* 54:981-991.
- Delavier-Klutchko, C., and M. Flavin. 1965. Enzymatic synthesis and cleavage of cystathionine in fungi and bacteria. *J. Biol. Chem.* 240:2537-2549.
- Dixon, M., and E. C. Webb. 1964. *Enzymes*, 2nd ed., p. 329. Longmans, Green and Co., London.
- Douglas, H. C., and G. Pelroy. 1963. A gene controlling inducibility of the galactose pathway enzymes in *S. cerevisiae*. *Biochim. Biophys. Acta* 68:155-156.
- Fink, G. R. 1966. A cluster of genes controlling three enzymes in histidine biosynthesis. *Genetics* 53:445-449.
- Flavin, M., and C. Slaughter. 1967. Enzymatic synthesis of homocysteine or methionine directly from *O*-succinyl-homoserine. *Biochim. Biophys. Acta* 132:400-405.
- Galzy, P., and P. P. Slonimski. 1957. Evolution de la constitution enzymatique de la levure cultivée sur acide lactique ou sur glucose comme seule source de carbone. *Compt. Rend.* 245:2556-2558.
- Giovanelli, J., and S. H. Mudd. 1966. Enzymatic synthesis of cystathionine by extracts of spinach, requiring *O*-acetyl-homoserine or *O*-succinyl-homoserine. *Biochem. Biophys. Res. Commun.* 25:366-371.
- Giovanelli, J., and S. H. Mudd. 1967. Synthesis of homocysteine and cysteine by enzyme extracts of spinach. *Biochem. Biophys. Res. Commun.* 27:150-156.
- Johnston, J. R., and R. K. Mortimer. 1959. Use of snail digestive juice in isolation of yeast spore tetrads. *J. Bacteriol.* 78:292.
- Karasevitch, Y., and H. de Robichon-Szulmajster. 1963. Régulation métabolique de la biosynthèse de la méthionine et de la thréonine chez *S. cerevisiae*. II. Répression et inhibition de l'homosérine deshydrogénase. *Biochim. Biophys. Acta* 73:414-426.
- Kredich, N. M., and G. N. Tomkins. 1966. The enzyme synthesis of L-cysteine in *E. coli* and *S. typhimurium*. *J. Biol. Chem.* 241:4955-4965.
- Lacroute, F. 1968. Regulation of pyrimidine biosynthesis in *Saccharomyces cerevisiae*. *J. Bacteriol.* 95:824-832.
- Lineweaver, H., and D. Burk. 1934. The determination of enzyme dissociation constants. *J. Am. Chem. Soc.* 56:658-666.
- Lowry, O. H., N. J. Rosebrough, A. L. Farr, and R. J. Randall. 1951. Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* 193:265-275.
- McClary, D. O., W. L. Nulty, and G. R. Miller. 1959. Effect of potassium versus sodium in the sporulation of *Saccharomyces*. *J. Bacteriol.* 78:362-368.
- Maw, G. A. 1961. Ability of S-methyl-cysteine to annul the inhibition of yeast growth by L-ethionine and by S-ethyl-L-cysteine. *J. Gen. Microbiol.* 25:441-449.
- Mortimer, R. K. and D. C. Hawthorne, 1966. Genetic mapping in *S. cerevisiae*. *Genetics* 53:165-173.
- Nagai, S., and M. Flavin. 1967. Acetyl-homoserine: an intermediate in the fungi biosynthesis of methionine. *J. Biol. Chem.* 242:3884-3895.
- Pasternak, C. A., R. J. Ellis, M. C. Jones, R. K. Mortimer, and C. E. Crichton. 1965. The control of sulphate reduction in bacteria. *Biochem. J.* 96:270.
- Robbins, P. W., and F. Lipmann. 1958. Separation of the two enzymatic phases in active sulfate synthesis. *J. Biol. Chem.* 233:681-685.
- de Robichon-Szulmajster, H. 1967. Régulation du fonctionnement de deux chaînes de biosynthèse chez *S. cerevisiae*: thréonine-méthionine et isoleucine-valine. *Bull. Soc. Chim. Biol.* 49:1431-1462.
- de Robichon-Szulmajster, H., and H. Cherest, 1966. Résistance à l'éthionine chez *S. cerevisiae*. II. Etude physiologique. *Genetics* 54:993-1006.
- de Robichon-Szulmajster, H., and H. Cherest. 1967. Regulation of homoserine-*O*-transacetylase, first step in methionine biosynthesis in *S. cerevisiae*. *Biochem. Biophys. Res. Commun.* 28:256-262.
- de Robichon-Szulmajster, H., and P. P. Magee. 1968. The regulation of isoleucine-valine biosynthesis in *S. cerevisiae*. I. Threonine deaminase. *European J. Biochem.* 3:492-501.
- de Robichon-Szulmajster, H., Y. Surdin, Y. Karasevitch,



- and D. Corrivaux. 1965. Régulations métaboliques chez *S. cerevisiae*: biosynthèse de la méthionine et de la thréonine. Colloq. Intern. Centre Natl. Rech. Sci. (Paris). no. 124:255-269.
28. de Robichon-Szulmajster, H., Y. Surdin, and R. K. Mortimer. 1966. Genetic and biochemical studies on genes controlling the synthesis of threonine and methionine in *Saccharomyces*. *Genetics* 53:609-619.
29. Rowbury, R. J., and D. D. Woods. 1964. O-succinyl-homoserine as an intermediate in the synthesis of cystathionine by *E. coli*. *J. Gen. Microbiol.* 36:341-358.
30. Sakami, W., and G. Toennies. 1942. The investigation of amino acid reactions by methods of non aqueous titrimetry. II. Differential acylation of hydroxy groups, and a method for the preparation of the acetyl derivatives of hydroxy amino acids. *J. Biol. Chem.* 144:203-217.
31. Schlenk, F. 1955. Biochemical and cytological studies with sulfonium compounds, p. 48-65. In S. K. Shapiro and F. Schlenk (ed.), *Transmethylation and methionine biosynthesis*. Chicago Univ. Press, Chicago.
32. Sorsoli, W. A., M. Buettner, and L. W. Parks. 1968. Cystathionine metabolism in methionine auxotrophic and wild-type strains of *Saccharomyces cerevisiae*. *J. Bacteriol.* 95:1024-1029.
33. Surdin, Y., W. Sly, J. Sire, A. M. Bordes, and H. de Robichon-Szulmajster. 1965. Système d'accumulation des acides aminés chez *S. cerevisiae*: propriétés et contrôle génétique. *Biochim. Biophys. Acta* 107:546-566.
34. Tingle, M., A. Herman, and H. O. Halvorson. 1968. Characterization and mapping of histidine genes in *S. lactis*. *Genetics* 58:361-371.
35. de Vito, P. C., and J. Dreyfuss. 1964. Metabolic regulation of adenosine triphosphate sulfurylase in yeast. *J. Bacteriol.* 88:1341-1348.
36. Wiebers, J. L., and H. R. Garner. 1960. Metabolic relationship between cystathionine and methionine in *Neurospora*. *J. Bacteriol.* 80:51-60.
37. Wiebers, J. L., and H. R. Garner. 1967. Homocysteine and cysteine synthetases of *Neurospora crassa*. Purification, properties and feedback control of activity. *J. Biol. Chem.* 242:12-23.
38. Wiebers, J. R., and H. R. Garner. 1967. Acyl derivatives of homoserine as substrates for homocysteine synthesis in *N. crassa*, yeast and *E. coli*. *J. Biol. Chem.* 242:5644-5649