

Differential Sensing of Protein Influences by  
NO and CO Vibrations in Heme Adducts


**- Supporting Information -**

*Mohammed Ibrahim, Changliang Xu and Thomas G. Spiro\**

Department of Chemistry, Princeton University, Princeton, New Jersey 08544

**Table S1.** Key structural parameters and  $S^2$  values calculated by the different DFT functionals in this study, with selected structural information from experiments and other computations for comparison

	Fe-NO(Å)	N-O (Å)	Fe-N-O (deg)	Fe-L (Å)	Fe-Np(Å)	$S^2$	Ref
<b>Five-coordinate</b>							
<b>Experimental</b>							
Fe(TPP)(NO)	1.717(7)	1.122(12)	149.2(6)		2.001(3)		1
Fe(TpivPP)(NO)	1.716(15)	1.197(9)	143.8		1.981(26)		2
<Fe(Por)(NO)>	1.728(5)	1.168(1)	143.4(9)		2.008(3)		3
<b>Functional, basis (calc.)</b>							
B3LYP, VTZ	1.709	1.173	140.4		2.0725/2.0037	0.84	t.w.
B3PW91, VTZ	1.780	1.173	142.0		2.018/2.0023	1.42	t.w.
B3P86, VTZ	1.761	1.172	141.2		2.0133/1.9959	1.31	t.w.
BLYP, VTZ	1.716	1.196	142.5		2.0436/2.0154	0.77	t.w.
BPW91, VTZ	1.699	1.189	143.6		2.0296/2.001	0.77	t.w.
BP86, VTZ	1.699	1.191	143.2		2.0279/1.9995	0.77	t.w.
PW91/STO-TZP	1.704	1.182	144.8		2.025/2.002		4
BP86, TZVP	1.700	1.180	146.0				5
<b>Six-coordinate</b>							
<b>Experimental</b>							
(N-Melm)Fe(TPP)(NO)	1.750(2)	1.182(3)	137.7(2)	2.173(2)	2.008(13)		3
(Py)Fe(TpivPP)(NO)	1.742(5)	1.194(9)	133.4(5)	2.260(5)	2.009(2)		3
<(L)Fe(Por)(NO)>	1.752(4)	1.174(6)	138.5(11)		2.007		3
<b>Functional, basis (calc.)</b>							
B3LYP, VTZ	1.782	1.174	139.6	2.123	2.0299/2.02	0.86	t.w.
B3PW91, VTZ	1.763	1.171	140.0	2.099	2.0196/2.0075	0.86	t.w.
B3P86, VTZ	1.757	1.171	139.6	2.080	2.0149/2.0027	0.85	t.w.
BLYP, VTZ	1.749	1.199	139.3	2.236	2.0456/2.0263	0.77	t.w.
BPW91, VTZ	1.735	1.192	139.8	2.186	2.0315/2.0105	0.77	t.w.
BP86, VTZ	1.735	1.194	139.4	2.161	2.0304/2.0101	0.76	t.w.
PW91/STO-TZP	1.741	1.186	140.1	2.173	2.027/2.008		4
BP86, TZVP	1.730	1.190	140.0	2.180			5



**Figure S1.** Three highest HOMOs of Fe(II)P(CO) and (ImH)Fe(II)P(CO), which correspond to the selected HOMOs of Fe(II)P(NO) and (ImH)Fe(II)P(NO) in Figure 9 in the main article.

**Complete Reference 15:**

Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Montgomery, Jr., J. A.; Vreven, T.; Kudin, K. N.; Burant, J. C.; Millam, J. M.; Iyengar, S. S.; Tomasi, J.; Barone, V.; Mennucci, B.; Cossi, M.; Scalmani, G.; Rega, N.; Petersson, G. A.; Nakatsuji, H.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Klene, M.; Li, X.; Knox, J. E.; Hratchian, H. P.; Cross, J. B.; Bakken, V.; Adamo, C.; Jaramillo, J.; Gomperts, R.; Stratmann, R. E.; Yazyev, O.; Austin, A. J.; Cammi, R.; Pomelli, C.; Ochterski, J. W.; Ayala, P. Y.; Morokuma, K.; Voth, G. A.; Salvador, P.; Dannenberg, J. J.; Zakrzewski, V. G.; Dapprich, S.; Daniels, A. D.; Strain, M. C.; Farkas, O.; Malick, D. K.; Rabuck, A. D.; Raghavachari, K.; Foresman, J. B.; Ortiz, J. V.; Cui, Q.; Baboul, A. G.; Clifford, S.; Cioslowski, J.; Stefanov, B. B.; Liu, G.; Liashenko, A.; Piskorz, P.; Komaromi, I.; Martin, R. L.; Fox, D. J.; Keith, T.; Al-Laham, M. A.; Peng, C. Y.; Nanayakkara, A.; Challacombe, M.; Gill, P. M. W.; Johnson, B.; Chen, W.; Wong, M. W.; Gonzalez, C.; and Pople, J. A. *Gaussian 03*; Gaussian, Inc.: Wallingford, CT, 2004.

**References**

1. Scheidt, W. R.; Frisse, M. E. *J. Am. Chem. Soc.* **1975**, *97*, 17-21.
2. Nasri, H.; Haller, K. J.; Wang, Y.; Hanh, H. B.; Scheidt, W. R. *Inorg. Chem.* **1992**, *31*, 3459 – 3467.
3. Wyllie, G. R. A.; Schulz, C. E.; Scheidt, W. R. *Inorg. Chem.* **2003**, *42*, 5722-5734.
4. Tangen, E.; Svadberg, A.; Ghosh, A. *Inorg. Chem.* **2005**, *44*, 7802-7805.
5. Praneeth, V. K. K.; Neese, F.; Lehnert, N. *Inorg. Chem.* **2005**, *44*, 2570-2572.