

NeuroImage

www.elsevier.com/locate/ynimg NeuroImage 42 (2008) 1045-1046

Erratum

Erratum to "A novel tensor distribution model for the diffusion-weighted MR signal" [NeuroImage 37 (2007) 164–176]

Bing Jian,^a Baba C. Vemuri,^{a,*} Evren Özarslan,^b Paul R. Carney,^c and Thomas H. Mareci^d

^aDepartment of Computer and Information Science and Engineering, University of Florida, P.O. Box 116120, Gainesville, FL 32611, USA ^bSection on Tissue Biophysics and Biomimetics, LIMB, NICHD, National Institutes of Health, Bethesda, MD 20892, USA ^cDepartment of Pediatrics, University of Florida, Gainesville, FL 32610, USA ^dDepartment of Biochemistry and Molecular Biology, University of Florida, P.O. Box 100245, Gainesville, FL 32610, USA

Available online 1 April 2008

Because of errors introduced by the typesetter, the following corrections should be made:

• Equation (4) currently displayed as

$$S(\mathbf{q}) = S_0 \int_{\mathcal{P}_n} \exp\left[-b\mathbf{g}^T \mathsf{D}\mathbf{g}\right]$$
$$dF = S_0 \int_{\mathcal{P}_n} f(\mathsf{D}) \exp\left[-b\mathbf{g}^T \mathsf{D}\mathbf{g}\right] d\mathsf{D}$$

is incorrect. The correct form is

$$\begin{split} S(\mathbf{q}) &= S_0 \int_{\mathcal{P}_n} \exp\left[-b\mathbf{g}^T \mathbf{D}\mathbf{g}\right] \mathrm{d}F \\ &= S_0 \int_{\mathcal{P}_n} f(\mathbf{D}) \exp\left[-b\mathbf{g}^T \mathbf{D}\mathbf{g}\right] \mathrm{d}\mathbf{D}. \end{split}$$

E-mail address: vemuri@cise.ufl.edu (B.C. Vemuri). Available online on ScienceDirect (www.sciencedirect.com).

DOI of original article: 10.1016/j.neuroimage.2007.03.074 * Corresponding author.

^{1053-8119/\$ -} see front matter ${\ensuremath{\mathbb C}}$ 2008 Elsevier Inc. All rights reserved. doi:10.1016/j.neuroimage.2008.03.033

• Equation (5) currently displayed as

$$S(\mathbf{q}) = S_0 \int_{\mathcal{P}_n} \exp(-b\mathbf{g}^T \mathsf{D}\mathbf{g})$$
$$\mathbf{d}F = S_0 \int_{\mathcal{P}_n} \exp(-trace(\mathsf{B}\mathsf{D}))$$

is incorrect. The correct form is

$$\begin{split} S(\mathbf{q}) &= S_0 \int_{\mathcal{P}_n} \exp\bigl(-b \mathbf{g}^T \mathsf{D} \mathbf{g}\bigr) \mathrm{d} F \\ &= S_0 \int_{\mathcal{P}_n} \exp(-trace(\mathsf{B}\mathsf{D})) \mathrm{d} F. \end{split}$$

- In Eq. (8), $(\Theta + \sum -1) \in \mathcal{P}_n$ is incorrect and should read $(\Theta + \sum^{-1}) \in \mathcal{P}_n$.
- Equation (10) currently displayed as,

$$S(\mathbf{q})/S_0(1+(b\mathbf{g}^T\hat{\mathbf{D}}\mathbf{g})/p)^{-p}$$

is incorrect. The correct form is

$$S(\mathbf{q}) = S_0 \left(1 + \left(b \mathbf{g}^T \hat{\mathbf{D}} \mathbf{g} \right) / p \right)^{-p}$$

- On page 168, in the last line of text before Eq. (16), F(D) should be changed to dF.
- In Fig. 2, the range of the x axis, "std. dev. of noise," should be from 0 to 0.1 as stated in the text, not from 0 to 1 as currently displayed in Fig. 2.
- Eq. (18) currently displayed as

$$(2\pi i)^{-n(n+1)/2} \int_{Re\mathbb{Z}=X_0} \mathscr{D}f(\mathbb{Z}) \exp[\operatorname{trace}(\mathbb{Y}\mathbb{Z})]$$
$$d\mathbb{Z} = \begin{cases} f(\mathbb{Y}), & \text{for } \mathbb{Y} \in \mathcal{P}_n\\ 0, & \text{otherwise} \end{cases}$$

is incorrect. The correct form is

$$(2\pi i)^{-n(n+1)/2} \int_{Re\mathbb{Z}=\mathsf{X}_0} \mathscr{L}f(\mathbb{Z}) \exp[\operatorname{trace}(\mathsf{Y}\mathbb{Z})] d\mathbb{Z}$$
$$= \begin{cases} f(\mathsf{Y}), & \text{for } \mathsf{Y} \in \mathcal{P}_n \\ 0, & \text{otherwise.} \end{cases}$$