



a

IC70 titer

	A1	A2	A3	A4	A5	A6	A/D1	A/D2	D1	D2	D3	C1	C2	C3	B5	B15	Mean	Median	Breadth score	Potency score
QA013 264 dpi	91	52				87	57										37	25	4	21.0
QC370 302 dpi	84																29	25	1	16.0
QA465 227 dpi	64	83		66	65	90	62	63		*auto	56			63	62		51	62	9	31.1
QD435 299 dpi																	25	25	0	15.3
QB008 303 dpi	55																27	25	0	15.7
QC406 379 dpi	156	55												*auto			35	25	2	17.1
QB099 283 dpi	80																28	25	0	16.0
QD370 278 dpi																	25	25	0	15.3
QA413 275 dpi	56																27	25	0	15.7
QC890 306 dpi		60															27	25	1	16.7
QA509 368 dpi	68					89	62				52						36	25	3	20.9
QC036 385 dpi	174	67															37	25	2	18.8
QB045 695 dpi																	25	25	0	15.3
QA281 656 dpi	57					56											29	25	1	16.9
QA692 707 dpi	424	297	118			56	63	88						147			89	25	7	45.0
QB247 667 dpi	50																27	25	0	15.6
QB726 468 dpi	100	83															33	25	2	18.5
QC413 416.5 dpi	190	57			76												40	25	3	20.6
QA874 463 dpi	65	82					60										33	25	2	19.5
QB670 321 dpi	119	66	82		58	58	57	58		57	61	69		64	57		57	58	12	33.0
QD022 439 dpi																	25	25	0	15.3
QA779 481 dpi	170	203															45	25	2	24.2
QA028 451 dpi	143	58															34	25	2	18.1
QA538 410 dpi																	25	25	0	15.3



Supplemental Figure 1. NAb responses by IC70 at ~1 year post-infection. (a) IC70 values for plasma/virus combinations are presented as described in the legend to Figure 2a. (b) Comparison of breadth scores calculated using IC70 values between superinfection cases and matched controls prior to superinfection, as for Figure 2b. (c) Comparison of potency scores calculated using IC70 values between superinfection cases and matched controls prior to superinfection, as for Figure 2c.