

Supplementary Table 2. Clinical and biochemical characteristics of three groups

	Control	LVD		SVD	
	n=158	n=221	P	n=134	P
Age (years)	65.08±10.44	66.61±10.47	0.670	66.96±7.62	0.086
Male/Female	103/55	160/61	0.143	76/58	0.149
BMI(kg/m ²)	23.30±2.20	24.40±2.40	0.845	24.50±3.20	0.127
Hypertension (%)	41.14	67.87	<0.0001	61.19	0.001
DM2 (%)	17.72	31.67	0.003	25.37	0.117
Smoking (%)	7.59	14.93	0.036	14.17	0.86
Alcohol intake(%)	2.63	4.66	0.500	3.05	1.000
HDL-C(mmol/L)	2.73±1.13	2.17±1.14	<0.0001	2.68±1.02	0.026
LDL-C(mmol/L)	1.42±0.64	1.98±1.10	<0.0001	1.46±0.80	0.007
TC(mmol/L)	4.50±0.76	4.72±0.95	0.002	4.74±0.92	0.010
TG(mmol/L)	1.48±0.62	1.75±1.10	<0.0001	1.65±0.98	0.043
FIB(g/L)	3.30±0.76	4.04±1.24	<0.0001	3.75±1.04	0.005

LVD: stroke caused by large-artery atherosclerosis; SVD: stroke caused by small-vessel occlusion; BMI: body mass index; DM2: diabetes mellitus type 2; LDL-C: low density lipoprotein cholesterol; HDL-C: high density lipoprotein cholesterol; TC: total cholesterol; TG: triglycerides; FIB: fibrinogen.