

**Variant IRAK-1 haplotype is associated with increased NF- κ B
activation and worse outcomes in sepsis**

**John Arcaroli, Eliezer Silva, James P Maloney, Qianbin He, Daiva
Svetkauskaite, James R Murphy, Edward Abraham**

Online Data Supplement

PRIMARY DATA FOR FIGURE 1

Patient Number	Haplotype	LPS 0	LPS 30	LPS 60	NF- κ B AUC
pt020	Variant	1	2.52	2.81	148.24
pt021	Variant	1	3.26	1.66	173.20
pt024	Variant	1	4.06	3.06	191.43
pt025	Variant	1	1.32	1.00	65.79
pt028	Variant	1	1.82	1.00	91.08
pt035	Variant	1	2.00	1.75	128.27
pt056	Variant	1	2.12	1.83	105.78
pt018	Wild type	1	1.78	1.11	85.83
pt019	Wild type	1	1.61	2.43	95.79
pt022	Wild type	1	1.00	1.36	66.43
pt026	Wild type	1	1.48	1.99	93.98
pt027	Wild type	1	1.80	1.59	104.04
pt038	Wild type	1	1.06	1.00	67.72
pt039	Wild type	1	1.24	1.44	84.60
pt040	Wild type	1	1.60	1.27	67.00
pt041	Wild type	1	1.25	1.11	73.13
pt042	Wild type	1	1.28	1.36	77.97
pt043	Wild type	1	1.64	1.41	88.66
pt044	Wild type	1	1.16	1.43	72.76
pt045	Wild type	1	1.28	1.12	72.17
pt046	Wild type	1	1.36	1.07	72.25
pt047	Wild type	1	1.90	2.04	99.03
pt048	Wild type	1	1.28	1.27	76.31
pt049	Wild type	1	1.94	1.25	95.01
pt050	Wild type	1	1.00	1.20	66.93
pt051	Wild type	1	1.78	1.12	96.27
pt052	Wild type	1	3.15	3.07	161.28
pt053	Wild type	1	1.16	1.08	78.42
pt055	Wild type	1	1.09	1.00	62.32
pt057	Wild type	1	1.57	1.42	87.65