

Supplementary Table 4: Table summarizing the prevalence (percentage of infected hosts) and the mean intensity (number of parasite individuals of a given species per infected host) of all recorded parasites in the 53 dissected fish.

	Prevalence (%)	Mean Intensity
Protozoa:		
<i>Trichodina sp.</i>	88.46	18.62
<i>Apiosoma sp.</i>	9.62	5.50
<i>Ichthyophthirius multifiliis</i>	11.54	2.33
Monogenea:		
<i>Gyrodactylus sp.</i>	28.85	17.80
Digenea:		
<i>Diplostomum sp. (total)</i>	100.00	15.98
<i>D. pseudospathaceum (lab infection)</i>	100.00	5.06
<i>Cyathocotyle prussica</i>	98.08	4.35
<i>Echinochasmus sp.</i>	98.08	4.83
<i>Apatemon cobitidis</i>	71.15	4.70
<i>Tylodelphis clavata</i>	98.08	6.93
<i>Phyllodistomum folium</i>	25.00	3.75
Cestoda:		
<i>Valipora campylancristrota</i>	84.62	11.93
<i>Proteocephalus filicolis</i>	86.54	6.87
Nematoda:		
<i>Anguillicola crassus</i>	69.23	1.64
<i>Contraecaecum sp.</i>	13.46	1.00
<i>Camallanus lacustris</i>	86.54	4.33
<i>Raphidascaaris acus</i>	71.15	2.97
Acanthocephala:		
<i>Acanthocephalus clavula</i>	34.62	2.00
<i>Acanthocephalus lucii</i>	78.85	1.00
Crustacea:		
<i>Argulus foliaceus</i>	15.38	1.91
<i>Ergasilus sp.</i>	9.62	1.00
Mollusca:		
Glochidia	23.08	1.00