

Suppl. Table 2. FWHM changes in the widening of the plasmalemma-bound GFP fluorescence signal in BRI1-GFP expressing hypocotyl and root cells before (0 min) and 30 min after application of 10 nM BL (30 min). The statistical analysis revealed an expansion of the cell wall by $34 \pm 22 \%$ ($n = 31$, $p = 0,0003$). The measurements were performed on 31 independent cells derived from 5 independent seedlings.

FWHM [μm]	FWHM [μm]	expansion [%]
0 min	30 min	
0,95	1,26	33
1,09	1,60	47
1,05	1,20	14
1,35	1,58	17
1,25	1,35	8
0,93	1,01	9
0,85	1,51	78
0,84	1,18	40
0,87	0,98	13
1,43	1,98	38
1,63	2,84	74
1,40	1,60	14
1,26	1,50	19
0,80	1,20	50
1,44	1,59	10
1,37	2,32	69
1,01	1,25	24
0,79	1,05	33
1,13	1,88	66
0,84	1,07	27
0,79	1,24	57
0,63	0,85	35
0,83	1,31	58
1,01	1,68	66
0,94	1,46	55
0,92	1,19	29
2,07	2,18	5
1,40	1,70	21
1,85	2,15	16
1,36	1,48	9
1,56	1,93	24
mean		34
standard deviation		22