

TABLE S1. Plasmids and bacterial strains.

Strain/plasmid	Relevant genotype or features	Source
Strains	Genotype	
BL21(DE3)	<i>E. coli</i> B F ⁻ <i>ompT hsdS_B(r_B⁻ m_B⁻) gal dcm</i> (DE3)	(4)
EN2	BL21(DE3) Δ <i>dnaK::kan</i>	This study
PopC4617	MC4100 Δ <i>dnaK::kan</i>	P. Delepelaire
Plasmids	Features	
pTrc99	<i>trc</i> promotor containing expression vector	Amersham Bioscience
pTrc-DnaK	<i>E. coli dnaK</i> cloned into the pTrc99 vector	This study
pIVEX-CpxP	<i>E. coli cpxP</i> cloned into the pIVEX2.4d vector	M. Miot and J-M. Betton
pET-PA28 α	Rat PA28 α cDNA cloned into the pET16b vector	(3)
pET-ClpP1	<i>Mycobacterium tuberculosis Rv2461c</i> cloned into the pET22b vector	M. Ortiz and N. Benaroudj
pET-ClpP2	<i>M. tuberculosis Rv2460c</i> cloned into the pET22b vector	M. Ortiz and N. Benaroudj
pNB41	<i>Methanococcus janaschii s4</i> cloned into the pET14b vector	(2)
pET-GFP	GFP cDNA cloned into the pET16b vector	(5)
pRSETa/Nemo	Murine Nemo cDNA cloned into pRSETa vector	(1)

1. **Agou, F., F. Ye, S. Goffinont, G. Courtois, S. Yamaoka, A. Israel, and M. Veron.** 2002. NEMO trimerizes through its coiled-coil C-terminal domain. *J Biol Chem* **277**:17464-75.
2. **Benaroudj, N., and A. L. Goldberg.** 2000. PAN, the proteasome-activating nucleotidase from archaeobacteria, is a protein-unfolding molecular chaperone. *Nat Cell Biol* **2**:833-9.
3. **Song, X., J. D. Mott, J. von Kampen, B. Pramanik, K. Tanaka, C. A. Slaughter, and G. N. DeMartino.** 1996. A model for the quaternary structure of the proteasome activator PA28. *J Biol Chem* **271**:26410-7.
4. **Studier, F. W., A. H. Rosenberg, J. J. Dunn, and J. W. Dubendorff.** 1990. Use of T7 RNA polymerase to direct expression of cloned genes. *Methods Enzymol* **185**:60-89.
5. **Weber-Ban, E. U., B. G. Reid, A. D. Miranker, and A. L. Horwich.** 1999. Global unfolding of a substrate protein by the Hsp100 chaperone ClpA. *Nature* **401**:90-3.