

## Supplementary Methods

### Strains and reporter transgenes

N2 Bristol wild-type <sup>1</sup> and CB4856 Hawaiian wild-type isolates <sup>2</sup> were used. The transgenic animals used were *vtIs1 [dat-1::gfp]* <sup>3</sup> in the manual screen and *vtIs1 plus vsIs33 [dop-3::rfp]* <sup>4</sup> for the worm sorter screen.

### Screening process

In all screens, animals were mutagenized with EMS according to standard protocols <sup>1</sup>. Worms were kept at 25°C at all times. In the manual screen 5 parental (P<sub>0</sub>) mutagenized animals were placed in each of 10 founder P<sub>0</sub> plates. Three days later, 400 F<sub>1</sub> progeny of the mutagenized P<sub>0</sub> animals were singled. Their ensuing progeny (F<sub>2</sub> and F<sub>3</sub> generation) were screened under a stereomicroscope equipped with a fluorescent light source.

In the worm sorter screen, 5 large (150 mm) founder P<sub>0</sub> plates were each seeded with 5 to 10 mutagenized P<sub>0</sub> worms. After 6 days the F<sub>2</sub>/F<sub>3</sub> generation was directly screened by rinsing the 5 P<sub>0</sub> plates and passing half of the worm population through the COPAS biosorter.

The sorting process for 5 P<sub>0</sub> plates lasted on average 5 hours. The work involved mostly waiting time during sample sorting and short washing steps in between plates. The reported total time of 25 days for performing the worm sorter screen includes all stages of the procedure, i.e., mutagenesis, worm handling, actual sorting, manually inspecting the sorted animals and lastly mutant handling (confirm the heritability of the phenotype, filing and freezing). The worm sorter screened all larval and adult stages.

In total we screened in the worm sorter screen 110 plates that translate to over 2,000,000 worms. Mutants were recovered in 21 of the plates. The mutants presented here are all independent isolates, i.e. mutants coming from different plates or mutants coming from the same plate showing different phenotype. When multiple mutants of the same phenotype were retrieved from the same P<sub>0</sub> plate they were considered as multiple isolates of the first isolated mutant and

are not reported in this paper. Note that this may lead to a slight underestimation of the number of mutants retrieved from the screen. The number of genomes screened with the worm sorter was estimated based on the initial number of  $P_0$  worms used (5 to 10) and the average number of  $F_1$  progeny our strain produced after mutagenesis (about 50). Therefore we estimated that we screened 500 to 1000 haploid genomes per  $P_0$  plate depending on whether the initial number of  $P_0$  worms was 5 or 10 respectively.

To calculate the false-positive and false-negative rates of recovering non-*gfp* animals from a *vt/s1* population (presented in **Supplementary Table1**) we mixed 50 N2 individuals with 10,000 *vt/s1* worms. For comparison we made the same experiment using a transgene that carries *vs/s33* in the background. The results are summarized in **Supplementary Table 1**.

In our experimental setup we would get on average 60 false positives from each plate of 25,000  $F_2$ s sorted. It would take us about 5 minutes to go through this number of worms under the dissecting scope and this would add a total of 30 minutes in the daily sorting routine; this number is included in the overall time effort calculation shown in **Table 2**.

### **Worm sorter settings**

The sorting process was performed according to the manufacturer's instructions and (Union Biometrica). The worms were gated to exclude eggs, small L1s and debris using as gating parameters Time of Flight vs. Extinction. Then they were sorted based on fluorescence parameters, in particular Red Peak Height vs. Green Peak Height. We set the parameters for the fluorescence signal amplification as follows:

Full scale: ToF 2048, Ext 2048, RedPH 65536, GreenPH 65536

Gains (Signal): Ext 3, Green 4, Red 4 Trigger: Ext

Thresholds: Singal 30, TOF Minimum: 10

PMT Control: Green 750, Red 750

For the test runs in which we determined the efficiency of sorting non-*gfp* animals when *rfp* transgene (*vs/s33*) was present or not present in the

background (**Supplementary Table 1**), we used as sorting parameters Tof vs. GreenPH.

### **Mapping and allele identification**

We first tested mutants for linkage to the X chromosome by crossing them with wild-type males and examining a potential mutant phenotype in hemizygous, male F<sub>1</sub> cross-progeny. If a phenotype was observed in the F<sub>1</sub> male progeny, it was tested whether this is due to dominance rather than to X-linkage.

For placing the mutants into complementation groups, we first grouped them into phenotypic categories. Then we performed complementation tests among mutants of the same phenotypic category. Subsequently we mapped at least one -and in most cases more than one- mutant per complementation group, using a rapid SNP mapping procedure<sup>5</sup>. Last, we complemented mutants of different phenotypic categories that mapped in the same region.


The two retrieved mutants that showed no *gfp* expression (*ot344* and *ot373*) are likely array mutations based on the following analysis: mutant animals were crossed with N2 males and the cross progeny showed no recovery of the *gfp* expression indicating either a mutation on the array or a dominant mutation. Then mutants were crossed with wild-type *vtIs1* males. The cross progeny were not "off" for *gfp* as we would expect in the case of a dominant mutation but showed normal fluorescence levels.

### **References**


1. Brenner, S. The genetics of *Caenorhabditis elegans*. *Genetics* **77**, 71-94 (1974).
2. Hodgkin, J. & Doniach, T. Natural variation and copulatory plug formation in *Caenorhabditis elegans*. *Genetics* **146**, 149-64 (1997).
3. Nass, R. et al. A genetic screen in *Caenorhabditis elegans* for dopamine neuron insensitivity to 6-hydroxydopamine identifies dopamine transporter mutants impacting transporter biosynthesis and trafficking. *J Neurochem* **94**, 774-85 (2005).

4. Chase, D.L., Pepper, J.S. & Koelle, M.R. Mechanism of extrasynaptic dopamine signaling in *Caenorhabditis elegans*. *Nat Neurosci* **7**, 1096-103 (2004).
5. Davis, M.W. et al. Rapid single nucleotide polymorphism mapping in *C. elegans*. *BMC Genomics* **6**, 118 (2005).

**Supplementary Fig.1: Phenotypes of isolated dopy mutants.**  
 Micrographs of representative animals and quantification of aberrant *dat-1::gfp* expression patterns are shown.



**Supplementary Figure 2 Molecular identity of mutant alleles of known genes retrieved from screens.** Names of alleles isolated from our screens are in blue, previously described reference alleles are in black. Lines indicate deletions. For quantification see Supplementary Tables 1 to 3.



**Supplementary Table 1: False positive/negative rates**

<b>Efficiency of sorting non-<i>gfp</i> (non-<i>vtIs1</i>) animals</b>						
<b>Input</b>		<b>Recovery</b>			<b>Rates</b>	
# of <i>vtIs1</i> animals	# of N2 animals	# worms sorted	N2 recovered	False positive worms	% false negative (N2 <i>not</i> recovered)	% false positive
10,000	50	63	38	25	24% (12/50)	40% (25/63)
<b>Efficiency of sorting non-<i>gfp</i> (non-<i>vtIs1</i>) animals with <i>rfp</i> transgene (<i>vsIs33</i>) in the background</b>						
<b>Input</b>		<b>Recovery</b>			<b>Rates</b>	
# <i>vtIs1;vsIs33</i> animals	# <i>vsIs33</i> animals	# worms sorted	<i>vsIs33</i> recovered	False positive worms	% false negative ( <i>vsIs33 not</i> recovered)	% false positive
10,000	50	71	50	21	0% (0/50)	30% (21/71)

Supplementary Table 2: Quantification of defects in *lin-32* mutants.

	% animals with <i>dat-1::gfp</i> expression:					% animals showing any phenotype	n
	<i>missing in CEPVs</i>	<i>missing in CEPDs</i>	<i>missing in ADEs</i>	<i>missing in PDEs</i>	<i>in extra cells near ADEs</i>		
wild type	0	0	0	0	0	0	100
<i>ot259</i>	0	61	0	77	7	95	55
<i>ot297</i>	0	94	17	100	74	100	42
<i>ot341</i>	0	84	1	99	93	100	38
<i>ot343</i>	0	71	4	100	14	100	38
<i>ot338</i>	0	100	7	100	60	100	30
<i>ot366</i>	0	100	20	100	76	100	42
<i>tm2044</i>	1	81	31	100	33	100	80
<i>tm1446</i>	0	100	8	100	95	100	42
<i>u282</i>	49	100	40	100	2	100	34

*lin-32(u282)* was used as reference allele. The *tm* alleles are putative null alleles (deleting parts of the bHLH domain; [www.wormbase.org](http://www.wormbase.org)) and were provided by the *C. elegans* knockout consortium in Japan.



**Supplementary Table 3: Quantification of defects in *vab-3* mutant.**

	% animals with <i>dat-1::gfp</i> expression:						n
	<i>in extra cells</i>	<i>in 1 extra cells</i>	<i>in 2 extra cells</i>	<i>in 3 extra cells</i>	<i>in 4 extra cells</i>	<i>missing in 1 CEPD</i>	
<b>wild type</b>	0	0	0	0	0	0	0
<b><i>ot346</i></b>	30	22	8	0	0	21	52
<b><i>u648</i></b>	88	40	21	17	4	0	48

Supplementary Table 4: Quantification of defects in *ham-1* mutants.

	% animals with <i>dat-1::gfp</i> expression:							<i>n</i>
	<i>missing in 1 or 2 CEPDs</i>	<i>in extra cells near CEPDs</i>	<i>missing in 1 or 2 ADEs</i>	<i>in extra cells near ADEs</i>	<i>missing in 1 or 2 PDEs</i>	<i>in extra cells near PDEs</i>	<i>any phenotype</i>	
wild type	0	0	0	0	0	0	0	100
<b><i>ot253</i></b>	23	13	0	95	0	15	98	40
<b><i>ot257</i></b>	40	14	0	80	0	0	94	50
<b><i>ot342</i></b>	47	17	0	83	0	23	90	30
<b><i>ot361</i></b>	17	0	3	90	0	30	93	30
<b><i>ot367</i></b>	40	0	3	90	0	20	100	30
<b><i>ot339</i></b>	47	20	3	90	0	13	97	30
<b><i>ot371</i></b>	40	10	3	73	27	6	100	30
<b><i>ot364</i></b>	45	13	0	80	0	26	94	31

**Supplementary Table 5: Calculating the degree of saturation of the worm sorter screen.**

Model	Maximum likelihood	$\Delta$ AICc	Weight	% saturation	95% credible interval [lower - upper]
Poisson	$1.46 \times 10^{-5}$	0	.817	93.3%	77.9%-98.6%
Gamma	$1.84 \times 10^{-5}$	4.53	.085	84.1%	51.4%-97.7%
2 Class	$2.11 \times 10^{-5}$	4.27	.097	80.7%	42.1%-99.7%
3 Class	$2.29 \times 10^{-5}$	14.10	.000	76.8%	72.5%-99.9%
4 Class	$2.22 \times 10^{-5}$	14.10	.000	73.8%	85.1%-99.9%
2 Class Variable	$2.21 \times 10^{-5}$	14.17	.000	79.8%	65.1%-99.6%

We calculated the degree of saturation based on the allele frequencies for the worm sorter isolated mutants *dopy-3* (3 alleles), *dopy-3* (1 allele), *dopy-7* (2 alleles), *lin-32* (4 alleles), *ham-1* (6 alleles) and *vab-3* (1 allele). Summary of Maximum Likelihood, AICc (Akaike Information Criterion) values and weights are shown. Poisson, multiple rates, and gamma distribution-based predictions for this mutagenesis were obtained as described (Pollock and Larkin, 2004). AICc, a measure of the quality of fit to an estimated statistical model, accounts for smaller data sets and therefore was used in this study to ascertain best fit to the given models (Akaike, 1978).  $\Delta$ AIC represents the difference between the AIC for each model and the lowest AIC. The weight of evidence in favor of each model being the best fit is given as 'weight.' The "95% credible interval" indicates the upper and lower bounds of the saturation estimates and were determined by removing the highest and lower 2.5% of saturation values computed by SatMut. The Poisson model assumes an equal rate of mutability. The 2-, 3- and 4-class models are multiple rate models that assume two to four mutation rate classes which occur at equal frequencies, i.e., an equal number of genes that are highly mutable as are lowly mutable. The 2-class variable frequency models are multiple rate models which assume two to four mutation rate classes that occur at two frequencies throughout the genome, i.e., a few genes highly mutable and many lowly mutable. The Poisson model (shaded in grey) best fits this data.

**Supplementary Table 6: Mutants retrieved from genetic screens**

Gene name	Molecular identity	Manual screen	Sorter screen	Allele names	Affected cells	Dom/Rec	Visible pleiotropies	Location
<i>dopy-2</i>	unknown	0 alleles	3 alleles	<i>ot340, ot345, ot406</i>	CEPs CEPDs, ADEs	R	-	III:-7 to -1
<i>dopy-3</i>	unknown	0 alleles	1 allele	<i>ot337</i>	CEPs, ADEs	D	-	I:+14 to +26
<i>dopy-4</i>	unknown	1 allele *	0 alleles	<i>ot260</i>	PDEs	R	-	I: left of -6
<i>dopy-5</i>	unknown	4 alleles **	0 alleles	<i>ot283, ot284, ot296, ot298</i>	CEPDs	R	sterile	III:-7 to -1
<i>dopy-6</i>	unknown	1 allele *	0 alleles	<i>ot263</i>	PDEs	R	-	X:-17 to -8
<i>dopy-7</i>	unknown	0 alleles	2 alleles	<i>ot399, ot347</i>	CEPs, PDEs	R	sick	X: left of -4
<i>lin-32</i>	bHLH	2 alleles	4 alleles	<i>ot259, ot297, ot341, ot343, ot338, ot366</i>	CEPDs, ADEs, PDEs	R	-	X:-16.01
<i>ham-1</i>	no homologies	2 alleles	6 alleles	<i>ot253, ot257, ot342, ot361, ot367, ot339, ot371, ot364</i>	CEPDs, ADEs, PDEs	R	-	IV:+6.02
<i>vab-3</i>	paired + homeodomain	0 alleles	1 allele	<i>ot346</i>	unknown	R	notched head	X:+2.22
<b>Total allele number</b>		10	17					
<b>Genomes screened</b>		11,000	80,000					
<b>Allele frequency per genomes screen</b>		1/1,100	1/4,700					
<b>Time investment</b>		100 days §	25 days §					
<b>Allele frequency per time</b>		1 allele / 10 days	1 allele / 1.5 days					

This is a more extensive version of Table 2, shown in the main body of the paper. Dom/Rec indicates “dominant/recessive”. § These days are differentially spent. 100 days dissecting scope work mean full time work at the microscope while 25 days of worm sorting involves mainly machine running and casual observation of functioning of sorter.