

**Supplemental Data****Mitochondrial Haplogroup U5b3:****A Distant Echo of the Epipaleolithic in Italy****and the Legacy of the Early Sardinians**

Maria Pala, Alessandro Achilli, Anna Olivieri, Baharak Hooshiar Kashani, Ugo A. Perego, Daria Sanna, Ene Metspalu, Kristiina Tambets, Erika Tamm, Matteo Accetturo, Valeria Carossa, Hovirag Lancioni, Fausto Panara, Bettina Zimermann, Gabriela Huber, Nadia Al-Zahery, Francesca Brisighelli, Scott R. Woodward, Paolo Francalacci, Walther Parson, Antonio Salas, Doron M. Behar, Richard Villems, Ornella Semino, Hans-Jürgen Bandelt, and Antonio Torroni

**Table S1. List of U5b3 Complete Sequences Included in Figure 1**

<b>Sequence ID<sup>a</sup></b>	<b>Haplogroup</b>	<b>Geographic Origin</b>	<b>GenBank ID</b>	<b>References</b>
1	U5b3	Sicily	GQ129143	[1]
2 <sup>b</sup>	U5b3	Southern France	GQ129144	[1]
3	U5b3	Bosnia	GQ129145	[1]
4	U5b3	Croatia	GQ129146	[1]
5	U5b3	Spain	GQ129147	[1]
6	U5b3	Piedmont - Northern Italy	GQ129148	[1]
7	U5b3a1a	Sardinia	GQ129149	[1]
8	U5b3a1a	Sardinia	GQ129150	[1]
9 <sup>c</sup>	U5b3a1a	Sardinia	GQ129151	[1]
10	U5b3a1a	Sardinia	DQ523655	[2]
11	U5b3a1a	Sardinia	DQ523658	[2]
12	U5b3a1a	Sardinia	DQ523664	[2]
13	U5b3a1a	Sardinia	GQ129152	[1]
14	U5b3a1a	Sardinia	GQ129153	[1]
15	U5b3a1a	Sardinia	DQ523628	[2]
16	U5b3a1a	Sardinia	DQ523644	[2]
17	U5b3a1a	Sardinia	DQ523669	[2]
18	U5b3a1a	Sardinia	GQ129154	[1]
19	U5b3a1a	Sardinia	GQ129155	[1]
20	U5b3a1a	Sardinia	EU597508	[3]
21	U5b3a1a	Sardinia	GQ129156	[1]
22	U5b3a1a	Undefined - Italy	AF346988	[4]
23	U5b3a1a	Sardinia	DQ523624	[2]
24 <sup>b</sup>	U5b3a1b	Southern France	GQ129157	[1]
25	U5b3a1b	Delaware - USA	GQ129158	[1]
26	U5b3a2	Marche - Central Italy	GQ129159	[1]

27	U5b3a2	Morocco	GQ129160	[1]
28	U5b3a2	Marche - Central Italy	GQ129161	[1]
29	U5b3a2	Marche - Central Italy	GQ129162	[1]
30	U5b3a2	England - UK	EU926618	[1]
31	U5b3a2	Pennsylvania - USA	GQ129163	[1]
32	U5b3a2	Estonia	GQ129164	[1]
33	U5b3b	Crete - Greece	GQ129165	[1]
34	U5b3b	Spain	GQ129166	[1]
35	U5b3b	Bohemia - Czech Rep.	GQ129167	[1]
36	U5b3b	Marche - Central Italy	GQ129168	[1]
37	U5b3b	England - UK	EU130681	[1]
38	U5b3b	Palatinate - Germany	GQ129169	[1]
39	U5b3c	Sardinia	GQ129170	[1]
40	U5b3c	Cantabria - Spain	GQ129171	[1]
41	U5b3c	Sicily	GQ129172	[1]
42	U5b3d	Southern Spain	GQ129173	[1]
43	U5b3d	Iraq	GQ129174	[1]
44	U5b3e	Scotland - UK	GQ129175	[1]
45 <sup>d</sup>	U5b3e	Southern Germany	GQ129176	[1]
46	U5b3e	Poland	GQ129177	[1]
47	U5b3e	Bulgaria	GQ129178	[1]
48	U5b3f	USA	GQ129179	[1]
49	U5b3f	Latiun - Central Italy	GQ129180	[1]
50	U5b3f	Latiun - Central Italy	GQ129181	[1]
51	U5b3f	Catalonia - Spain	GQ129182	[1]
52	U5b3f	Latiun - Central Italy	GQ129183	[1]

<sup>a</sup> ID numbers correspond to the numbers in Figure 1.<sup>b</sup> The control-region sequence of this mtDNA was previously published.<sup>5</sup><sup>c</sup> The control-region sequence of this mtDNA was previously published.<sup>6</sup><sup>d</sup> The control-region sequence of this mtDNA was previously published.<sup>7</sup>

**Table S2. Population Distribution of Haplogroup U5b3**

Region, Population/Country	Sample size	U5b3 <sup>a</sup>		References
		Number of subjects	%	
<i>Western Europe:</i>				
Austria	374	0	-	[8, 9]
Balearic Islands	256	4	1.56	[10, 11]
Basques	237	0	-	[5, 6, 12, 13]
Denmark	247	0	-	[1, 6]
England	335	2	0.60	[1, 6, 14]
Finland	405	0	-	[1, 6, 15, 16]
France	234	1	0.43	[1, 5, 6, 17]
France Centre-South	240	2	0.83	[5, 18]
France Centre-West	124	0	-	[5]
France North & Belgium	267	1	0.37	[5, 6, 19]
France-Corse	99	0	-	[11, 20]
France-Normandie-Bretagne	383	0	-	[5, 18]
Germany Centre	115	0	-	[1, 21, 22]
Germany North	426	0	-	[21, 23-25]
Germany South	356	1	0.28	[7, 21, 26]
Iceland	457	2	0.44	[6, 27]
Ireland	300	1	0.33	[28]
Norway	554	1	0.18	[6, 14]
Portugal Centre	160	0	-	[29, 30]
Portugal North	184	2	1.09	[29, 30]
Portugal South	196	0	-	[29, 30]
Saami	255	0	-	[1, 15, 31, 32]
Scotland	1199	3	0.25	[14]
Spain	554	3	0.54	[1, 33-36]
Spain North-East	562	2	0.36	[1, 13, 37]
Spain North-West	439	1	0.23	[1, 13, 30, 38, 39]
Spain South	641	5	0.78	[1, 10, 11, 13, 34, 35]
Sweden	512	0	-	[1, 6]
Switzerland	434	1	0.23	[1, 40, 41]
Wales	92	0	-	[21]
<i>Western Europe-Italy:</i>				
Apulia Calabria	228	0	-	[1]
Campania	314	0	-	[1]
Latium	736	6	0.82	[1, 42, 43]
Lombardy	215	0	-	[1, 43]
Marche	1125	8	0.71	[1, 43]
Piedmont	1006	5	0.50	[1, 43]
Sardinia	1224	47	3.84	[1, 6, 11, 44]

Sicily	699	3	0.43	[1, 6, 45]
South Tyrol	540	0	-	[46, 47]
Tuscany	483	0	-	[1, 11, 43, 48, 49]
Umbria	202	0	-	[1]
<i>Eastern Europe:</i>				
Albania	242	0	-	[1, 6]
Bashkirs	332	0	-	[1]
Bosnia	512	2	0.39	[1, 50]
Bulgaria	268	1	0.37	[1, 6]
Byelorussia	351	0	-	[1]
Croatia	1405	4	0.28	[1, 42]
Czech Republic	83	0	-	[6]
Estonia	646	1	0.15	[6, 51, 52]
Greece	735	0	-	[1, 6, 53, 54]
Greece-Crete	395	1	0.25	[1, 53]
Greece-Lemnos	60	0	-	[1]
Hungary	647	2	0.31	[1, 55, 56]
Karelians	176	0	-	[1]
Macedonia	162	0	-	[1]
Montenegro	138	1	0.72	[1]
Poland	572	1	0.17	[1, 6, 57]
Rumania	453	0	-	[1, 6]
Russia	1879	0	-	[1, 6, 57-59]
Serbia	126	0	-	[1]
Slovakia	885	2	0.23	[1, 60, 61]
Slovenia	104	0	-	[50]
Ukraine	1142	0	-	[1, 59]
<i>Caucasus:</i>				
Armenia	191	1	0.52	[6]
Azerbaijan	48	0	-	[6]
Caucasus North	356	0	-	[6, 62, 63]
Georgia	113	0	-	[62, 64, 65]
<i>Middle East:</i>				
Iran	1311	0	-	[1, 63, 66]
Iraq	261	1	0.38	[1, 6]
Jordan	146	0	-	[6]
Kurds	82	0	-	[6, 64]
Pakistan	109	0	-	[62, 67]
Palestinians	227	0	-	[6, 68]
Syria	118	0	-	[6, 53]
Turkey	340	0	-	[6, 62, 69]
<i>Africa:</i>				
Algeria	125	0	-	[13, 35]
Egypt	525	0	-	[1, 70, 71]

Libya	129	0	-	[1]
Mauritania-Morocco	999	2	0.20	[1, 11, 35, 71, 72]
Senegal	240	0	-	[72, 73]
Tunisia	724	0	-	[1, 35, 74]
<b>TOTAL</b>	<b>34766</b>	<b>120</b>		

<sup>a</sup> For many of the published datasets only control-region data were available (from np 16030 to np 16365). In these cases, U5 mtDNAs were affiliated within U5b3 when lacking 16189 or 16256 and harbouring 16304 (with the exception of U5b3a1a clade in which 16304 is reverted).

### Supplemental References

1. This Study
2. Fraumene, C., Belle, E.M.S., Castrì, L., Sanna, S., Mancosu, G., Cocco, M., Marras, F., Barbujani, G., Pirastu, M., Angius, A. (2006). High resolution analysis and phylogenetic network construction using complete mtDNA sequences in Sardinian genetic isolates. *Mol. Biol. Evol.* 23, 2101-2111.
3. Hartmann, A., Thieme, M., Nanduri, L.K., Stempfl, T., Moehle, C., Kivisild, T., Oefner, P.J. (2009). Validation of microarray-based resequencing of 93 worldwide mitochondrial genomes. *Hum. Mutat.* 30, 115-122.
4. Ingman, M., Kaessmann, H., Pääbo, S., Gyllensten, U. (2000). Mitochondrial genome variation and the origin of modern humans. *Nature* 408, 708-713.
5. Richard, C., Pennarun, E., Kivisild, T., Tambets, K., Tolk, H.V., Metspalu, E., Reidla, M., Chevalier, S., Giraudet, S., Barać Lauc, L., et al. (2007). An mtDNA perspective of French genetic variation. *Ann. Hum. Biol.* 34, 68-79.
6. Richards, M., Macaulay, V., Hickey, E., Vega, E., Sykes, B., Guida, V., Rengo, C., Sellitto, D., Cruciani, F., Kivisild, T., et al. (2000). Tracing European founder lineages in the Near Eastern mtDNA pool. *Am. J. Hum. Genet.* 67, 1251-1276.
7. Brandstätter, A., Klein, R., Duftner, N., Wiegand, P., Parson, W. (2006). Application of a quasi-median network analysis for the visualization of character conflicts to a population sample of mitochondrial DNA control region sequences from southern Germany (Ulm). *Int. J. Legal. Med.* 120, 310-314.
8. Parson, W., Parsons, T.J., Scheithauer, R., Holland, M.M. (1998). Population data for 101 Austrian Caucasian mitochondrial DNA d-loop sequences: application of mtDNA sequence analysis to a forensic case. *Int. J. Legal. Med.* 111, 124-132.
9. Brandstätter, A., Niederstätter, H., Pavlic, M., Grubwieser, P., Parson, W. (2007). Generating population data for the EMPOP database - an overview of the mtDNA

- sequencing and data evaluation processes considering 273 Austrian control region sequences as example. *Forensic. Sci. Int.* 166, 164-175.
10. Picornell, A., Gómez-Barbeito, L., Tomàs, C., Castro, J.A., Ramon, M.M. (2005). Mitochondrial DNA HVRI variation in Balearic populations. *Am. J. Phys. Anthropol.* 128, 119-130.
11. Falchi, A., Giovannoni, L., Calo, C.M., Piras, I.S., Moral, P., Paoli, G., Vona, G., Varesi, L. (2006). Genetic history of some western Mediterranean human isolates through mtDNA HVR1 polymorphisms. *J. Hum. Genet.* 51, 9-14.
12. Bertranpetti, J., Sala, J., Calafell, F., Underhill, P.A., Moral, P., Comas, D. (1995). Human mitochondrial DNA variation and the origin of Basques. *Ann. Hum. Genet.* 59, 63-81.
13. Cörte-Real, H.B., Macaulay, V.A., Richards, M.B., Hariti, G., Issad, M.S., Cambon-Thomsen, A., Papiha, S., Bertranpetti, J., Sykes, B.C. (1996). Genetic diversity in the Iberian Peninsula determined from mitochondrial sequence analysis. *Ann. Hum. Genet.* 60, 331-350.
14. Helgason, A., Hickey, E., Goodacre, S., Bosnes, V., Stefánsson, K., Ward, R., Sykes, B. (2001). MtDNA and the islands of the North Atlantic: estimating the proportions of Norse and Gaelic ancestry. *Am. J. Hum. Genet.* 68, 723-737.
15. Lahermo, P., Sajantila, A., Sistonen, P., Lukka, M., Aula, P., Peltonen, L., Savontaus, M.L. (1996). The genetic relationship between the Finns and the Finnish Saami (Lapps): analysis of nuclear DNA and mtDNA. *Am. J. Hum. Genet.* 58, 1309-1322.
16. Hedman, M., Brandstätter, A., Pimenoff, V., Sistonen, P., Palo, J.U., Parson, W., Sajantila, A. (2007). Finnish mitochondrial DNA HVS-I and HVS-II population data. *Forensic Sci. Int.* 172, 171-178.
17. Rousselet, F., Mangin, P. (1998). Mitochondrial DNA polymorphisms: a study of 50 French Caucasian individuals and application to forensic casework. *Int. J. Legal. Med.* 111, 292-298.

18. Dubut, V., Chollet, L., Murail, P., Cartault, F., Béraud-Colomb, E., Serre, M., Mogentale-Profizi, N. (2004). MtDNA polymorphisms in five French groups: importance of regional sampling. *Eur. J. Hum. Genet.* 12, 293-300.
19. Decorte, R., Jehaes, E., Xiao, F.X. and Cassiman, J.J. (1996). Genetic analysis of single hair shafts by automated sequence analysis of the mitochondrial d-loop region. In *Advances in Forensic Haemogenetics* 6, A. Carracedo, B. Brinkmann, and W. Bärr, ed. (Berlin, Germany: Springer-Verlag) pp. 17-19.
20. Varesi, L., Memmi, M., Cristofari, M.C., Mameli, G.E., Calò, C.M., Vona, G. (2000). Mitochondrial control-region sequence variation in the Corsican population, France. *Am. J. Hum. Biol.* 12, 339-351.
21. Richards, M., Cörte-Real, H., Forster, P., Macaulay, V., Wilkinson-Herbots, H., Demaine, A., Papiha, S., Hedges, R., Bandelt, H.-J., Sykes, B. (1996). Paleolithic and neolithic lineages in the European mitochondrial gene pool. *Am. J. Hum. Genet.* 59, 185-203.
22. Hofmann, S., Jaksch, M., Bezold, R., Mertens, S., Aholt, S., Paprotta, A., Gerbitz, K.D. (1997). Population genetics and disease susceptibility: characterization of central European haplogroups by mtDNA gene mutations, correlation with D loop variants and association with disease. *Hum. Mol. Genet.* 6, 1835-1846.
23. Baasner, A., Schäfer, C., Junge, A., Madea, B. (1998). Polymorphic sites in human mitochondrial DNA control region sequences: population data and maternal inheritance. *Forensic Sci. Int.* 98, 169-178.
24. Pfeiffer, H., Brinkmann, B., Hühne, J., Rolf, B., Morris, A.A., Steighner, R., Holland, M.M., Forster, P. (1999). Expanding the forensic German mitochondrial DNA control region database: genetic diversity as a function of sample size and microgeography. *Int. J. Legal. Med.* 112, 291-298.
25. Tetzlaff, S., Brandstätter, A., Wegener, R., Parson, W., Weirich, V. (2007). Mitochondrial DNA population data of HVS-I and HVS-II sequences from a northeast German sample.

Forensic Sci. Int. 172, 218-224.

26. Lutz, S., Weisser, H.J., Heizmann, J., Pollak, S. (1998). Location and frequency of polymorphic positions in the mtDNA control region of individuals from Germany. Int. J. Legal. Med. 111, 67-77.
27. Helgason, A., Sigurðardóttir, S., Gulcher, J.R., Ward, R., Stefánsson, K. (2000). MtDNA and the origin of the Icelanders: deciphering signals of recent population history. Am. J. Hum. Genet. 66, 999-1016.
28. McEvoy, B., Richards, M., Forster, P., Bradley, D.G. (2004). The Longue Durée of genetic ancestry: multiple genetic marker systems and Celtic origins on the Atlantic facade of Europe. Am. J. Hum. Genet. 75, 693-702.
29. Pereira, L., Prata, M.J., Amorim, A. (2000). Diversity of mtDNA lineages in Portugal: not a genetic edge of European variation. Ann. Hum. Genet. 64, 491-506.
30. González, A.M., Brehm, A., Pérez, J.A., Maca-Meyer, N., Flores, C., Cabrera, V.M. (2003). Mitochondrial DNA affinities at the Atlantic fringe of Europe. Am. J. Phys. Anthropol. 120, 391-404.
31. Sajantila, A., Lahermo, P., Anttinen, T., Lukka, M., Sistonen, P., Savontaus, M.L., Aula, P., Beckman, L., Tranebjærg, L., Gedde-Dahl, T., et al. (1995). Genes and languages in Europe: an analysis of mitochondrial lineages. Genome Res. 5, 42-52.
32. Delghandi, M., Utsi, E., Krauss, S. (1998). Saami mitochondrial DNA reveals deep maternal lineage clusters. Hum. Hered. 48, 108-114.
33. Pinto, F., González, A.M., Hernández, M., Larruga, J.M., Cabrera, V.M. (1996). Genetic relationship between the Canary Islanders and their African and Spanish ancestors inferred from mitochondrial DNA sequences. Ann. Hum. Genet. 60, 321-330.
34. Larruga, J.M., Díez, F., Pinto, F.M., Flores, C., González, A.M. (2001). Mitochondrial DNA characterisation of European isolates: the Maragatos from Spain. Eur. J. Hum. Genet. 9, 708-716.

35. Plaza, S., Calafell, F., Helal, A., Bouzerna, N., Lefranc, G., Bertranpetti, J., Comas, D. (2003). Joining the pillars of Hercules: mtDNA sequences show multidirectional gene flow in the western Mediterranean. *Ann. Hum. Genet.* 67, 312-328.
36. Database Comisaría General Policía Científica de Madrid (<http://www.policia.es/cgpc/index.htm>).
37. Crespillo, M., Luque, J.A., Paredes, M., Fernández, R., Ramírez, E., Valverde, J.L. (2000). Mitochondrial DNA sequences for 118 individuals from northeastern Spain. *Int. J. Legal. Med.* 114, 130-132.
38. Salas, A., Comas, D., Lareu, M.V., Bertranpetti, J., Carracedo, A. (1998). MtDNA analysis of the Galician population: a genetic edge of European variation. *Eur. J. Hum. Genet.* 6, 365-375.
39. Maca-Meyer, N., Sánchez-Velasco, P., Flores, C., Larruga, J.M., González, A.M., Oterino, A., Leyva-Cobián, F. (2003). Y chromosome and mitochondrial DNA characterization of Pasiegos, a human isolate from Cantabria (Spain). *Ann. Hum. Genet.* 67, 329-339.
40. Pult, I., Sajantila, A., Simanainen, J., Georgiev, O., Schaffner, W., Pääbo, S. (1994). Mitochondrial DNA sequences from Switzerland reveal striking homogeneity of European populations. *Biol. Chem. Hoppe. Seyler.* 375, 837-840.
41. Dimo-Simonin, N., Grange, F., Taroni, F., Brandt-Casadevall, C., Mangin, P. (2000). Forensic evaluation of mtDNA in a population from south west Switzerland. *Int. J. Legal. Med.* 113, 89-97.
42. Babalini, C., Martínez-Labarga, C., Tolk, H.V., Kivisild, T., Giampaolo, R., Tarsi, T., Contini, I., Barać L., Janićijević, B., Martinović Klarić, I., et al. (2005). The population history of the Croatian linguistic minority of Molise (southern Italy): a maternal view. *Eur. J. Hum. Genet.* 13, 902-912.
43. Turchi, C., Buscemi, L., Previderè, C., Grignani, P., Brandstätter, A., Achilli, A., Parson, W., Tagliabruni, A., Ge.F.I. Group. (2008). Italian mitochondrial DNA database: results of

- a collaborative exercise and proficiency testing. *Int. J. Legal. Med.* *122*, 199-204.
44. Di Rienzo, A., Wilson, A.C. (1991). Branching pattern in the evolutionary tree for human mitochondrial DNA. *Proc. Natl. Acad. Sci. USA* *88*, 1597-1601.
45. Vona, G., Ghiani, M.E., Calò, C.M., Vacca, L., Memmi, M., Varesi, L. (2001). Mitochondrial DNA sequence analysis in Sicily. *Am. J. Hum. Biol.* *13*, 576-589.
46. Pichler, I., Mueller, J.C., Stefanov, S.A., De Grandi, A., Volpato, C.B., Pinggera, G.K., Mayr, A., Ogriseg, M., Ploner, F., Meitinger, T., et al. (2006). Genetic structure in contemporary south Tyrolean isolated populations revealed by analysis of Y-chromosome, mtDNA, and Alu polymorphisms. *Hum. Biol.* *78*, 441-464.
47. Thomas, M.G., Barnes, I., Weale, M.E., Jones, A.L., Forster, P., Bradman, N., Pramstaller, P.P. (2008). New genetic evidence supports isolation and drift in the Ladin communities of the South Tyrolean Alps but not an ancient origin in the Middle East. *Eur. J. Hum. Genet.* *16*, 124-134.
48. Francalacci, P., Bertranpetti, J., Calafell, F., Underhill, P.A. (1996). Sequence diversity of the control region of mitochondrial DNA in Tuscany and its implications for the peopling of Europe. *Am. J. Phys. Anthropol.* *100*, 443-460.
49. Achilli, A., Olivieri, A., Pala, M., Metspalu, E., Fornarino, S., Battaglia, V., Accetturo, M., Kutuev, I., Khusnutdinova, E., Pennarun, E., et al. (2007). Mitochondrial DNA variation of modern Tuscan supports the near eastern origin of Etruscans. *Am. J. Hum. Genet.* *80*, 759-768.
50. Malyarchuk, B.A., Grzybowski, T., Derenko, M.V., Czarny, J., Drobnić, K., Miścicka-Sliwka, D. (2003). Mitochondrial DNA variability in Bosnians and Slovenians. *Ann. Hum. Genet.* *67*, 412-425.
51. Tambets, K., Rootsi, S., Kivisild, T., Help, H., Serk, P., Loogväli, E.L., Tolk, H.V., Reidla, M., Metspalu, E., Pliss, L., et al. (2004). The western and eastern roots of the Saami—the story of genetic "outliers" told by mitochondrial DNA and Y chromosomes. *Am. J. Hum. Genet.* *75*, 98-112.

- Genet. 74, 661-682.
52. Pliss, L., Tambets, K., Loogväli, E.L., Pronina, N., Lazdins, M., Krumina, A., Baumanis, V., Villem, R. (2006). Mitochondrial DNA portrait of Latvians: towards the understanding of the genetic structure of Baltic-speaking populations. Ann. Hum. Genet. 70, 439-458.
53. Vernesi, C., Di Benedetto, G., Caramelli, D., Secchieri, E., Simoni, L., Katti, E., Malaspina, P., Novelletto, A., Marin, V.T., Barbujani, G. (2001). Genetic characterization of the body attributed to the evangelist Luke. Proc. Natl. Acad. Sci. USA 98, 13460-13463.
54. Irwin, J., Saunier, J., Strouss, K., Paintner, C., Diegoli, T., Sturk, K., Kovatsi, L., Brandstätter, A., Cariolou, M.A., Parson, W., Parsons, T.J. (2008). Mitochondrial control region sequences from northern Greece and Greek Cypriots. Int. J. Legal. Med. 122, 87-89.
55. Brandstätter, A., Egyed, B., Zimmermann, B., Duftner, N., Padar, Z., Parson, W. (2007). Migration rates and genetic structure of two Hungarian ethnic groups in Transylvania, Romania. Ann. Hum. Genet. 71, 791-803.
56. Irwin, J., Egyed, B., Saunier, J., Szamosi, G., O'Callaghan, J., Padar, Z., Parsons, T.J. (2007). Hungarian mtDNA population databases from Budapest and the Baranya county Roma. Int. J. Legal. Med. 121, 377-383.
57. Malyarchuk, B.A., Grzybowski, T., Derenko, M.V., Czarny, J., Woźniak, M., Miścicka-Sliwka, D. (2002). Mitochondrial DNA variability in Poles and Russians. Ann. Hum. Genet. 66, 261-283.
58. Orekhov, V., Poltoraus, A., Zhivotovsky, L.A., Spitsyn, V., Ivanov, P., Yankovsky, N. (1999). Mitochondrial DNA sequence diversity in Russians. FEBS Lett. 445, 197-201.
59. Malyarchuk, B.A., Derenko, M.V. (2001). Mitochondrial DNA variability in Russians and Ukrainians: implication to the origin of the Eastern Slavs. Ann. Hum. Genet. 65, 63-78.
60. Metspalu, E., Kivisild, T., Kaldma, K., Reidla, M., Villem, R. (1998). Genbank direct submission.
61. Lehocký, I., Baldovic, M., Kádasi, L., Metspalu, E. (2008). A database of mitochondrial

- DNA hypervariable regions I and II sequences of individuals from Slovakia. *Forensic Sci. Int. Genet.* 2, e53-59.
62. Quintana-Murci, L., Chaix, R., Wells, R.S., Behar, D.M., Sayar, H., Scozzari, R., Rengo, C., Al-Zahery, N., Semino, O., Santachiara-Benerecetti, A.S., et al. (2004). Where west meets east, the complex mtDNA landscape of the southwest and Central Asian corridor. *Am. J. Hum. Genet.* 74, 827-845.
63. Derenko, M., Malyarchuk, B., Grzybowski, T., Denisova, G., Dambueva, I., Perkova, M., Dorzhu, C., Luzina, F., Lee, H.K., Vanecek, T., et al. (2007). Phylogeographic analysis of mitochondrial DNA in northern Asian populations. *Am. J. Hum. Genet.* 81, 1025-1041.
64. Comas, D., Calafell, F., Bendukidze, N., Fañanás, L., Bertranpetti, J. (2000). Georgian and kurd mtDNA sequence analysis shows a lack of correlation between languages and female genetic lineages. *Am. J. Phys. Anthropol.* 112, 5-16.
65. Alfonso-Sánchez, M.A., Martínez-Bouzas, C., Castro, A., Peña, J.A., Fernández-Fernández, I., Herrera, R.J., de Pancorbo, M.M. (2006). Sequence polymorphisms of the mtDNA control region in a human isolate: the Georgians from Swanetia. *J. Hum. Genet.* 51, 429-439.
66. Metspalu, M., Kivisild, T., Metspalu, E., Parik, J., Hudjashov, G., Kaldma, K., Serk, P., Karmin, M., Behar, D.M., Gilbert, M.T., et al. (2004). Most of the extant mtDNA boundaries in south and southwest Asia were likely shaped during the initial settlement of Eurasia by anatomically modern humans. *BMC Genet.* 5, 26.
67. Kivisild, T., Bamshad, M.J., Kaldma, K., Metspalu, M., Metspalu, E., Reidla, M., Laos, S., Parik, J., Watkins, W.S., Dixon, M.E., et al. (1999). Deep common ancestry of Indian and western-Eurasian mitochondrial DNA lineages. *Curr. Biol.* 9, 1331-1334.
68. Behar, D.M., Metspalu, E., Kivisild, T., Rosset, S., Tzur, S., Hadid, Y., Yudkovsky, G., Rosengarten, D., Pereira, L., Amorim, A., et al. (2008). Counting the founders: the matrilineal genetic ancestry of the Jewish Diaspora. *PLoS ONE* 3, e2062.

69. Di Benedetto, G., Ergüven, A., Stenico, M., Castrì, L., Bertorelle, G., Togan, I., Barbujani, G. (2001). DNA diversity and population admixture in Anatolia. *Am. J. Phys. Anthropol.* 115, 144-156.
70. Stevanovitch, A., Gilles, A., Bouzaid, E., Kefi, R., Paris, F., Gayraud, R.P., Spadoni, J.L., El-Chenawi, F., Béraud-Colomb, E. (2004). Mitochondrial DNA sequence diversity in a sedentary population from Egypt. *Ann. Hum. Genet.* 68, 23-39.
71. Coudray, C., Olivier, A., Achilli, A., Pala, M., Melhaoui, M., Cherkaoui, M., El-Chennawi, F., Kossmann, M., Torroni, A., Dugoujon, J.M. (2009). The complex and diversified mitochondrial gene pool of Berber populations. *Ann. Hum. Genet.* 73, 196-214.
72. Rando, J.C., Pinto, F., González, A.M., Hernández, M., Larruga, J.M., Cabrera, V.M., Bandelt, H.-J. (1998). Mitochondrial DNA analysis of northwest African populations reveals genetic exchanges with European, near-eastern, and sub-Saharan populations. *Ann. Hum. Genet.* 62, 531-550.
73. Graven, L., Passarino, G., Semino, O., Boursot, P., Santachiara-Benerecetti, A.S., Langaney, A., Excoffier, L. (1995). Evolutionary correlation between control region sequence and restriction polymorphisms in the mitochondrial genome of a large Senegalese Mandenka sample. *Mol. Biol. Evol.* 12, 334-345.
74. Fadhlaoui-Zid, K., Plaza, S., Calafell, F., Ben Amor, M., Comas, D., Bennamar El gaaied, A (2004). Mitochondrial DNA heterogeneity in Tunisian Berbers. *Ann. Hum. Genet.* 68, 222-233.