

Supporting Information

Yamauchi et al. 10.1073/pnas.0810319106




Fig. S1. The $^{13}\text{C}\{^1\text{H}\}$ NMR (500 MHz, D_2O , 300 K) spectrum of $2\text{D}(1)_3$.




Fig. S2. The ^1H DOSY NMR (500 MHz, D_2O , 300 K) spectrum of $2\text{D}(1)_3$.




Fig. S3. The ^1H - ^1H COSY (500 MHz, D_2O , 300 K) spectrum of $2\text{Pd}(1)_3$.




Fig. S4. The ^1H - ^{13}C HSQC and HMBC (500 MHz, D_2O , 300 K) spectrum of $2\text{D}(1)_3$.




Fig. S5. The ¹H NMR (500 MHz, D₂O, 300 K) spectrum of 6-1.





Fig. S6. The $^{13}\text{C}\{^1\text{H}\}$ NMR (500 MHz, D_2O , 300 K) spectrum of $6\text{D}1$.




Fig. S7. The ¹H-¹³C HSQC (500 MHz, D₂O, 300 K) spectrum of 6D1.




Fig. S8. ORTEP drawing of 2'-(1)3 (50% probability ellipsoids).




Fig. S9. UV-vis spectrum of 1 in solid.

Other Supporting Information Files

[Table S1](#)

[Table S2](#)

[Table S3](#)

[Table S4](#)