

Table S17. One-way ANOVAs of *SW/RW* ratio in *Arabidopsis* accessions. Comparison between infected (I) and mock-inoculated (M) plants at each plant density.

N° Plants	Treatment	Boa-0				Cen-1				Ler			
		<i>n</i>	<i>df</i>	<i>F</i>	<i>P</i>	<i>n</i>	<i>df</i>	<i>F</i>	<i>P</i>	<i>n</i>	<i>df</i>	<i>F</i>	<i>P</i>
1 Plant													
	<i>I</i>	30	1	12.87	1x10 ⁻⁵	30	1	6.96	1x10 ⁻³	30	1	14.39	1x10 ⁻⁵
2 Plants													
	<i>M/I</i>	30	1	1.59	0.216	30	1	6.46	0.015	30	1	5.97	0.021
	<i>I/I</i>	30	1	0.16	0.696	30	1	0.02	0.897	30	1	1.32	0.260
4 Plants													
	<i>M/M/M/I</i>	30	1	7.76	0.002	30	1	5.88	0.024	30	1	7.25	0.009
	<i>M/M/I/I</i>	30	1	6.32	0.018	30	1	0.15	0.704	30	1	0.04	0.844
	<i>M/I/M/I</i>	30	1	4.62	0.035	30	1	0.67	0.420	30	1	0.00	0.954
	<i>M/I/I/I</i>	30	1	0.06	0.802	30	1	4.21	0.042	30	1	5.91	0.024
	<i>I/I/I/I</i>	30	1	18.01	1x10 ⁻⁵	30	1	0.80	0.498	30	1	6.10	0.019

Plant densities and treatments are listed on the left. ***n***: number of observations. ***df***: degrees of freedom. ***F***: *F*-value from the type III sum of squares ANOVA for each factor. ***P***: Estimated probability of obtaining this *F*-value under the null hypothesis.