

ERRATA

Evaluation of Three Commercial Agglutination Tests for the Identification of *Staphylococcus aureus*

DANIEL R. PENNELL, JEAN A. ROTT-PETRI, AND TERRENCE A. KURZYNSKI

Wisconsin State Laboratory of Hygiene, University of Wisconsin, Madison, Wisconsin 53706

Volume 20, no. 4, p. 616, Table 2, column 1: Under the entry "Tube coagulase," "6 h" should read "24 h."

Sensitivity and Specificity of Monoclonal Antibodies Directed Against Antigenic Determinants of *Treponema pallidum* Nichols in the Diagnosis of Syphilis

MICHAEL V. NORGDARD, COLEEN K. SELAND, JOHN R. KETTMAN, AND JAMES N. MILLER

Department of Microbiology, University of Texas Health Science Center at Dallas, Dallas, Texas 75235, and Treponemal Research Laboratory, Department of Microbiology and Immunology, University of California at Los Angeles School of Medicine, Los Angeles, California 90024

Volume 20, no. 4, p. 716, column 2, lines 32–40: These lines should read "Consequently, the use of current monoclonal antibody technology for the detection of *T. pallidum* may be as much as 10- to 100-fold more sensitive than that achieved through the use of dark-field microscopic examination. Further, the substitution of monoclonal antibody for polyclonal antibody in the direct fluorescent-antibody–dark-field microscopic technique (10) may result in a procedure with improved sensitivity, specificity, and reproducibility."

Improved Biotyping Schemes for *Campylobacter jejuni* and *Campylobacter coli*

R. MARTIN ROOP II, ROBERT M. SMIBERT, AND NOEL R. KRIEG

Departments of Biology (Microbiology Section) and Anaerobic Microbiology, Virginia Polytechnic Institute and State University, Blacksburg, Virginia 24061

Volume 20, no. 5, p. 991, Table 1: The results for *C. coli* under the heading "Hippurate hydrolysis" are for growth in strains minimal medium. (All *C. coli* biovariants were negative for hippurate hydrolysis, as indicated in Table 2.)