

**Table s1. Prevalence and intensity of injection and non-injection drug use among a cohort of Thai injection drug users  
(n = 252)**

<b>Type of Drug</b>	<b>Injection (times per day)*</b>				<b>Non-injection (times per day)*</b>			
	<b><u>Less than daily</u></b>	<b><u>1 - 2</u></b>	<b><u>&gt; 2 daily</u></b>	<b><u>Total</u></b>	<b><u>Less than Daily</u></b>	<b><u>1 - 2</u></b>	<b><u>&gt; 2 Daily</u></b>	<b><u>Total</u></b>
<b>Heroin</b>	74 (29.4%)	113 (44.8%)	42 (16.7%)	<b>229 (90.9%)</b>	1 (0.4%)	9 (3.6%)	9 (3.6%)	<b>19 (7.5%)</b>
<b>Yaba/Ice</b>	59 (23.4%)	66 (26.2%)	20 (7.9%)	<b>143 (56.7%)</b>	35 (13.9%)	32 (12.7%)	26 (10.3%)	<b>93 (36.9%)</b>
<b>Ecstasy</b>	--	--	--	--	5 (2.0%)	4 (1.6%)	--	<b>9 (3.6%)</b>
<b>Midazolam</b>	34 (13.5%)	98 (38.9%)	34 (13.5%)	<b>166 (65.9%)</b>	1 (0.4%)	7 (2.8%)	2 (0.8%)	<b>10 (4.0%)</b>
<b>Methadone</b>	3 (1.2%)	9 (3.6%)	2 (0.8%)	<b>14 (5.6%)</b>	3 (1.2%)	73 (29.0%)	--	<b>76 (30.2%)</b>
<b>Opiate<sup>†</sup></b>	7 (2.8%)	4 (1.6%)	--	<b>11 (4.4%)</b>	2 (0.8%)	3 (1.2%)	--	<b>5 (2.0%)</b>
<b>Morphine</b>	3 (1.2%)	1 (0.4%)	--	<b>4 (1.6%)</b>	--	--	--	--
<b>Benzodiazepine</b>	--	--	--	--	6 (2.4%)	14 (5.6%)	10 (4.0%)	<b>30 (11.9%)</b>
<b>Ketamine</b>	--	--	--	--	4 (1.6%)	1 (0.4%)	--	<b>5 (2.0%)</b>

*Note: Ecstasy, Benzodiazepine and Ketamine are only consumed through non-injection.*

*\*Percentages denote proportion of the total study sample*

*<sup>†</sup>Opiate refers to opiates other than heroin or methadone*