

Supplemental Table 1. Retinal thickness and visual acuity outcomes.

	Week				
	0	3-4	12-16	24-28	48-52
Patient 1					
OD: BCVA	83	87	79	85	73
OCT (µm)	237	253	249	241	287
OS: BCVA	90	92	88	85	82
OCT (µm)	223	223	254	260	325
Patient 2					
OD: BCVA	77	76	82	78	80
OCT (µm)	242	256	177	217	162
OS: BCVA	60	63	68	78	77
OCT (µm)	Unable	Unable	213	195	193
Patient 3					
OD: BCVA	65	65	79	--	84
OCT (µm)	Unable	Unable	Unable	--	205
OS: BCVA	68	82	79	--	85
OCT (µm)	297	233	205	--	331
Patient 4					
OD: BCVA	71	73	79	78	79
OCT (µm)	--	141	136	130	139
OS: BCVA	59	49	50	35	87
OCT (µm)	--	180	Unable	169	169
Patient 5*					
OD: BCVA	74	71	74	73	--
OCT (µm)	--	304	278	296	--
OS: BCVA	35	57	71	74###	--
OCT (µm)	--	--	285	245	--

* Patient 5 elected to withdraw from the study at the 28-week follow-up visit due to travel constraints. At the time of study withdrawal, he was improving from a flare of his uveitis, deemed likely secondary to his missed daclizumab infusion at week 20.

Note: Patient underwent cataract surgery in the left eye for a visually significant posterior subcapsular cataract. This eye was excluded from statistical analysis of pre- and post-study visual acuities.

Patient was given sub-Tenon's triamcinolone acetate (40 mg) for uveitic flare OS at week 24. Visual acuity at that time was 55 ETDRS letters. VA subsequently improved to 74 letters at his week 28 visit.