

**Table. Interaction of IAS<sub>5</sub> with Antithrombin in the presence of varying concentrations of NaCl at pH 7.4 and 25 °C.<sup>@</sup>**

<b>[NaCl]</b>	<b>Slope</b>	<b>Intercept</b>	<b>K<sub>D</sub><sup>@</sup></b>
<i>(mM)</i>	<i>(<math>\mu M^{-1}</math>)</i>	<i>(<math>cm^2 \mu M^{-1} min^{-1} kV^{-1}</math>)</i>	<i>(<math>\mu M</math>)</i>
0	-0.0270 ± 0.0040 <sup>§</sup>	-0.0850 ± 0.0040	37 ± 6
10	-0.0167 ± 0.0024	-0.0390 ± 0.0010	60 ± 8
20	-0.0075 ± 0.0006	-0.0190 ± 0.0030	133 ± 10
50	-0.0039 ± 0.0002	-0.0094 ± 0.0007	256 ± 12

<sup>@</sup> The binding affinities at 0, 10, 20 and 50 mM NaCl were measured as described in 'Materials and Methods' and shown in Figure 5. Scatchard analysis was performed to obtain the slope and intercept from which K<sub>D</sub> values were calculated. <sup>§</sup> Errors represent ±2 S. E.