

# **Structural insights into what glycan arrays tell us about how glycan-binding proteins interact with their ligands**

**Maureen E. Taylor and Kurt Drickamer**

Division of Molecular Biosciences, Department of Life Sciences, Imperial College, London SW7 2AZ, UK

**Screening of glycan arrays represent a powerful, high-throughput approach to defining the selectivity of glycan-binding receptors for oligosaccharide ligands. Correlating results from such arrays with structural analysis of receptor-ligand complexes provides one way to validate the arrays. Using examples drawn from the family of proteins that contain C-type carbohydrate-recognition domains, this review illustrates how information from the arrays reflects the way that selectivity and affinity for glycan ligands is achieved. A range of binding phenotypes is observed, from very restricted binding to a small set of structurally similar ligands to binding of broad classes of ligands with related terminal sugars and even to failure to bind any of the glycans on an array. These outcomes provide insights into the importance of multiple factors in defining the selectivity of these receptors, including the presence of conformationally defined units in some oligosaccharide ligands, local and extended interactions between glycans and the surfaces of receptors, and steric factors that exclude binding of some ligands.**

## **Key words**

glycan array / glycan-binding receptor / lectin / structure / modeling

## **Introduction**

Glycan-binding receptors are involved in biological processes that require recognition of selected sets of oligosaccharide ligands. Receptor-mediated glycan-binding events underlie selective adhesion between mammalian cells, sorting and trafficking of soluble glycoproteins, and detection of microbial pathogens based on their surface glycosylation (Taylor and Drickamer 2006, Varki et al. 2009). Glycan arrays are increasingly being used as a means of characterizing the glycan-binding specificity of these receptors (Feizi et al. 2003, Paulson et al. 2006).

There are important practical issues that have been addressed in the construction and probing of glycan arrays (Figure 1). Multiple different platforms have been described for the display of glycans using a range of immobilization strategies. These include both noncovalent adsorption, such as direct immobilization on nitrocellulose (Wang et al. 2002), binding of lipid-conjugated glycans to nitrocellulose surfaces (Fukui et al. 2002), and binding of biotinylated ligands to streptavidin-coated surfaces (Guo et al. 2004), as well as various types of covalent linkages to surfaces through the reducing ends of oligosaccharides (Blixt et al. 2004, Karamanska et al. 2008, Manimala et al. 2006). The nature of the surface and the spacer between the glycan and the surface can affect interaction with glycan-binding proteins, often in unpredictable ways. Data shown here were obtained with glycans covalently attached to polymer-coated glass surfaces through spacers of different lengths in a format developed by the Consortium for Functional Glycomics (Blixt et al. 2004). Arrays are generally probed with soluble lectins or soluble fragments of membrane receptors that have been fluorescently labeled. Because of the extensive washing steps needed to remove unbound receptors, binding to arrays is generally enhanced through avidity resulting from multivalent binding of naturally oligomeric receptor fragments or clusters of carbohydrate-binding domains generated by oligomerization motifs such as immunoglobulin Fc domains or biotinylation tags that can be clustered on streptavidin tetramers (Paulson et al. 2006, Powlesland et al. 2006).

Although it is clear that glycan arrays are powerful tools for highly parallel analysis of the interaction of receptors with a broad spectrum of potential ligands, an understanding of the significance of the positive and negative results obtained is still being developed. Multiple approaches have been employed for array validation, most commonly involving solid phase binding assays employing labeling strategies to detect bound receptors (van Vliet et al. 2005), quantitative binding competition assays (Guo et al. 2004), isothermal titration calorimetry (Gregg et al. 2008, Neu et al. 2008), or surface plasmon resonance (Bochner et al. 2005, van Liempt et al. 2006). The focus of this review is on structural features of the binding proteins and the glycan ligands that explain some of the behaviors observed on the arrays. Examples are drawn from the family of glycan-binding proteins that contain C-type carbohydrate-recognition domains (CRDs) (Figure 2). Many of these domains have been tested on the array created by the Consortium for Functional Glycomics (CFG), creating one of the largest collections of reasonably comparable datasets available. Importantly, both the binding data and the structural coordinates are publicly available through the CFG web site ([www.functionalglycomics.org](http://www.functionalglycomics.org)) and the Protein Data Bank (DPB) ([www.rcsb.org](http://www.rcsb.org)).

## Receptors that display highly selective ligand binding

The most readily interpreted results on the glycan array are those that reflect binding to a very limited set of glycans that clearly share a structural motif. The endothelial scavenger receptor C-type lectin illustrates this type of situation, since probing of the array even at relatively high protein concentrations reveals selective binding to only a few glycans, all of which contain the Lewis<sup>x</sup> or Lewis<sup>a</sup> trisaccharide motifs (Figure 3a) (Coombs et al. 2005, Feinberg et al. 2007). Plotting the data in rank order of the signal strength observed on the array reveals a sharp drop-off in binding, with relatively few structures that fall in a twilight zone between strong signals and background. In addition to the presence of the Lewis<sup>x</sup> and Lewis<sup>a</sup> motifs in the group of positive ligands, it is also important to note the absence of these motifs in most of the glycans that do not bind to the receptor.

The Lewis<sup>x</sup> trisaccharide is a commonly recognized structural motif in ligands for glycan-binding receptors. In addition to the scavenger receptor C-type lectin, DC-SIGN (discussed below) and the selectins bind to ligands that contain this unit (Guo et al. 2004, Rosen 2004, van Die et al. 2002, Vestweber and Blanks 1999). An important feature of these glycans is that the three sugar residues that make up the Lewis<sup>x</sup> epitope are consistently in the same conformation in all known structures that contain the motif, either in free glycans or glycans bound to receptors (Feinberg et al. 2007, Miller et al. 1992, Perez et al. 1996, Somers et al. 2000). This structure results from van der Waals packing of the fucose and galactose residues against each other, forming a rigid and stable structure (Figure 3b). In the complex with the scavenger receptor C-type lectin, the 3- and 4-hydroxyl groups of the galactose residue of the Lewis<sup>x</sup> ligand form coordination bonds with a bound calcium ion, an arrangement that is characteristic of the C-type lectins (Figure 2b). A further feature of the Lewis<sup>x</sup> conformation is that the relatively nonpolar B face of galactose remains exposed, allowing it to interact in a characteristic packing arrangement with the side chain of tryptophan and other aromatic amino acids. In addition to these interactions with the galactose residue, the fucose residue also makes contact with other residues projecting from the protein surface as can be seen in a surface representation of the bound complex (Figure 1c).

A key consequence of the preformed, rigid conformation of the glycan ligand is that there is little or no conformational entropy penalty associated with the binding interaction. Crystal structures of C-type CRDs in the absence of glycan ligands reveal that the binding site is also preformed, so that the only change needed to accommodate the glycan ligand is release of water molecules that occupy the coordination positions that will be replaced by the hydroxyl groups of the sugar. Thus, the enthalpy gain associated with the secondary contacts with the

surface of the protein is not negated by an extensive entropy penalty, resulting in an overall more favorable free energy of interaction that is reflected in a 70-fold enhanced affinity for Lewis<sup>x</sup> trisaccharide compared to galactose, as measured in binding competition assays (Coombs et al. 2005, Feinberg et al. 2007).

Although many oligosaccharides display conformational heterogeneity, the Lewis<sup>a</sup> and Lewis<sup>x</sup> trisaccharides are not the only examples of preferred local conformations that create preformed structural features that can interact with receptors with a minimal entropy penalty associated with glycan immobilization. In the case of another endothelial cell receptor, LSECtin, glycan array screening at high receptor concentration reveals binding to multiple groups of ligands with terminal fucose and GlcNAc residues, but at reduced concentrations much more selective binding is observed to glycans bearing terminal GlcNAc $\beta$ 1-2Man disaccharides (Figure 4a) (Powlesland et al. 2008). Analysis of the conformation of this disaccharide in multiple crystal structures reveals that in many cases it assumes a common preferred conformation in which the N-acetyl group of GlcNAc is positioned above the mannose residue, making van der Waals interactions with the B face of the hexose (Figure 4b,c) (Petrescu et al. 1999). This is one of two conformations of the disaccharide that have been observed, but the fact that it appears to be a preferred if not necessarily completely stable, preformed conformation would still reduce the entropy penalty associated with binding to a receptor. Although potential interactions with the binding site in LSECtin have been modeled, these remain to be examined experimentally.

In summary, glycan array results indicating highly selective binding to a group of glycans can reflect the presence of a preformed conformational feature in the ligands, which provides an extended surface for local interactions with the receptor surface at minimal entropy cost associated with glycan immobilization.

### **Receptors that bind to broad classes of glycan ligands**

One of the first receptors tested against glycan arrays was the receptor DC-SIGN, which appears to have a dual role in interaction of dendritic cells with pathogen surfaces as well as with T cells (Engering et al. 2002, Geijtenbeek et al. 2000). The glycan array analysis reveals binding to two classes of ligands: high mannose oligosaccharides and Lewis-type structures (Figure 5a), while structural analysis suggests that the mechanisms underlying interactions with the two classes of ligands differ significantly (Guo et al. 2004). The Lewis<sup>x</sup> trisaccharide sits in the binding site as a rigid unit, as in the case of the scavenger receptor C-type lectin, but placement of the fucose rather than the galactose residue in the primary binding site results in a more limited set of interactions (Figure 5b,c). Notably, the prominent phenylalanine residue on

the surface of the CRD is not positioned sufficiently close to the oligosaccharide to allow packing interactions with the galactose residue. Nevertheless, the limited contacts observed are probably sufficient to generate specificity because of the rigid structure of the oligosaccharide ligand.

In contrast, binding to the more flexible high mannose ligands involves a different energetic balance. Because high mannose oligosaccharides are known to assume multiple conformations, no one of which would exactly match the conformation needed to dock into the binding site, the entropy cost associated with freezing out one of these structures would be substantial. The structure of a ligand with five mannose residues in the binding site can be modeled on the available crystal structure of a mimic containing the core trimannose structure capped with two GlcNAc residues (Figure 5d,e) (Feinberg et al. 2007, Feinberg et al. 2001). The model reveals that by snaking through a groove on the surface of the CRD and wrapping around the surface phenylalanine residue, the oligosaccharide can make many favorable interactions. Thus, binding of this class of ligand must involve a significant trade-off between entropy costs for immobilization of the ligand and additional favorable contacts in a much more extended binding site. The binding of DC-SIGN to high mannose oligosaccharides thus provides a second paradigm for how relatively narrow specificity for a class of ligands can be achieved.

Other receptors display binding to a still broader spectrum of ligands on the glycan array, which structurally appears to correlate with binding largely to terminal residues in glycans. For example, ranking of ligands for the macrophage galactose lectin detected on the glycan array reveals preferential binding to glycans bearing GalNAc residues that are either unsubstituted or substituted only on the 6 position (Figure 6a), with simple GalNAc being one of the most prominent ligands. Although binding to galactose-terminated structures is also detected, these ligands generally give weaker signals. These results are consistent with other profiling methods that indicate a strong preference for GalNAc over galactose (van Vliet et al. 2005). Structural and mutagenesis studies indicate that GalNAc and galactose bind in the primary binding site on the CRD, ligated to the binding site calcium ion and packing against a tryptophan residue for increased affinity and specificity, with further selectivity for GalNAc resulting from favorable contacts with the 2-acetamido group (Figure 6b,c) (Iobst and Drickamer 1996, Kolatkar et al. 1998).

The overall trend in the types of glycans bound to the macrophage galactose receptor is evident from the array data and all of the ligands with high signals can be accounted for by interactions of a terminal sugar with the primary binding site. The fact that there are no other obvious common features of the most prominent ligands is consistent with the suggestion that specificity derives from interaction with the terminal sugar. However, many glycans that feature

apparently appropriate terminal residues do not rank amongst the best ligands and there are no obvious rules describing which ligands bind and which do not. Two factors are generally correlated with lower ranking in the binding: linkage in  $\alpha$  rather than  $\beta$  configuration and linkage adjacent to a fucose residue. These observations illustrate the importance of a further mechanism that has been invoked to explain binding specificity, which is exclusion of some ligands by steric hindrance. Surface features of the CRD near to the binding site may block binding of some ligands, particularly those with fixed geometry such as the Lewis<sup>x</sup> structure, rather than enhance binding through favorable interactions. In such cases, affinity must be generated largely through the interactions with the terminal monosaccharides, which makes the packing interaction of the galactose with the tryptophan residue particularly important. Selectivity then results more from exclusion of some classes of ligands. A similar principle of exclusion probably explains why DC-SIGN binds poorly to some fucose-containing ligands and well to others, particularly given the position of the surface phenylalanine residue near to the primary binding site.

Comparing the glycan array profiles for the scavenger receptor C-type lectin and LSECtin (Figures 3a and 4a) with those for DC-SIGN and the macrophage galactose receptor (Figures 5a and 6a) reveals differences in the shapes and color distributions of the profiles. The first set of receptors, in which the binding sites make multiple favorable interactions with a relatively rigid structural motif, give strong signals with a few structurally related glycans, after which there is a sharp drop-off in signal, reflecting the fact that binding is dependent on the presence of the specific binding epitope. In contrast, the profiles for the second set of receptors are more extended, with a continuous decline in signal and no strict segregation of structurally distinct ligands. Although binding to these receptors reflects a nucleating interaction with a mannose-type or a galactose-type monosaccharide, the affinity for oligosaccharides results from a complex balance between entropy costs of immobilizing larger oligosaccharides, with resulting favorable interactions, and steric exclusion.

### **Receptors with low selectivity binding sites**

At the most extreme end of the glycan array results are receptors that are known to bind sugar ligands but which interact poorly with all of the oligosaccharides on the array. Serum mannose-binding protein represents such a case, since very little binding is evident even with highly fluorescent probe at high receptor concentrations (Figure 7a). There are no obvious common features to the glycans which give the highest signals and the pattern is not reproduced in multiple screens of the array. The binding site in this protein is open, so that there are very few

constraints on what binds beyond the key hydroxyl groups that coordinate with the binding site calcium ion (Figure 2a) (Ng et al. 2002, Weis et al. 1992). However, as a result of this arrangement, there are very few possibilities for further favorable interactions to enhance affinity as in the case of DC-SIGN (Figure 7b). The binding of mannose, GlcNAc and fucose in the primary binding site is not enhanced by packing with an adjacent tryptophan residue as seen for the galactose- and GalNAc-binding receptors. Also, binding involves only a limited portion of the terminal residue of oligosaccharide ligands because other portions of the glycans are not in direct contact with the surface of the protein, which contrasts sharply with the arrangement in DC-SIGN.

The inherently weak and low specificity of the interaction of mannose-binding protein with terminal sugars means that biologically significant binding can only occur through multivalent interactions. Although multivalency plays a role in increasing avidity of interactions of many glycan-binding receptors, for mannose-binding protein the geometry of the placement of multiple binding sites facing a single direction is a determining feature for broad recognition of glycan-rich surfaces without requiring specific complex oligosaccharide structures (Figure 7c) (Weis and Drickamer 1994). The wide spacing of the binding site in the mannose-binding protein trimer and their rigid arrangement means that although they are able to engage with repetitive sugar structures on surfaces of bacteria and fungi, they probably cannot adapt well to the spacing of glycans on the arrays (Figure 7d). Thus, the lack of binding to the glycan array reflects an alternative structural mechanism of optimizing interactions with sugars, in this case with minimal local selectivity so that spatial distribution becomes the dominant factor in determining specificity.

## **Conclusions and perspective**

The description of some of the factors that lead to specificity and affinity in glycan-receptor interactions presented in this review illustrate how the power of both glycan arrays and structural analysis is enhanced by examining the results of the two approaches in combination. The fact that the interpretations are largely consistent with each other provides validation for the use of the arrays to define binding specificity of receptors. The array allows screening of far more ligands than could be undertaken in a structural analysis. Viewing the limited set of structures of receptor-ligand complexes in the light of information about the relative binding signals obtained with additional ligands makes it possible to suggest how the specificity is achieved. Combining such information may provide a realistic and reliable basis for predicting which ligands will bind to particular receptors as well as for modeling of how they bind. It may also provide a foundation

for prediction and modeling of glycan ligand binding to other receptors for which structural information is not available.

### **Funding**

This work was supported by the Wellcome Trust [Grant 075565] and the National Institute of General Medical Sciences [Grant GM62116 to the Consortium for Functional Glycomics].

### **Acknowledgements**

We thank Eliot Ward and Adrián Quintero-Martinez for providing the human macrophage galactose receptor for glycan array screening and David Smith of the Consortium for Functional Glycomics for the array screening experiments.

### **Conflict of interest statement**

None declared.

### **Abbreviations**

CRD, carbohydrate-recognition domain; CFG, Consortium for Functional Glycomics; PDB, Protein Data Bank.

### **References**

- Blixt, O, Head, S, Mondala, T, Scanlan, C, Huflejt, ME, Alvarez, R, Bryan, MC, Fazio, F, Calarese, D, Stevens, J, Razi, N, Stevens, DJ, Skehel, JJ, van Die, I, Burton, DR, Wilson, IA, Cummings, R, Bovin, N, Wong, C-H, Paulson, JC. 2004. Printed covalent glycan array for ligand profiling of diverse glycan binding proteins. *Proc Natl Acad Sci USA*. 101:17033-17038.
- Bochner, BS, Alvarez, RA, Mehta, P, Bovin, NV, Blixt, O, White, JR, Schnaar, RL. 2005. Glycan array screening reveals a candidate ligand for siglec-8. *J Biol Chem*. 280:4307-4312.
- Coombs, PJ, Graham, SA, Drickamer, K, Taylor, ME. 2005. Selective binding of the scavenger receptor C-type lectin to Lewis<sup>x</sup> trisaccharide and related glycan ligands. *J Biol Chem*. 280:22993-22999.
- Engering, A, Geijtenbeek, TBH, van Vliet, SJ, Wijers, M, van Liempt, E, Demaurex, N, Lanzacecchia, A, Franssen, J, Figdor, CG, Piguet, V, van Kooyk, Y. 2002. The dendritic cell-specific adhesion receptor DC-SIGN internalizes antigen for presentation to T cells. *J Immunol*. 168:2118-2126.



- Feinberg, H, Castelli, R, Drickamer, K, Seeberger, PH, Weis, WI. 2007. Multiple modes of binding enhance the affinity of DC-SIGN for high-mannose N-linked glycans found on viral glycoproteins. *J Biol Chem.* 282:4202-4209.
- Feinberg, H, Mitchell, DA, Drickamer, K, Weis, WI. 2001. Structural basis for selective recognition of oligosaccharides by DC-SIGN and DC-SIGNR. *Science.* 294:2163-2166.
- Feinberg, H, Taylor, ME, Weis, WI. 2007. Scavenger receptor C-type lectin binds to the leukocyte cell surface glycan Lewis x by a novel mechanism. *J Biol Chem.* 282:17250-17258.
- Feizi, T, Fazio, F, Chai, W, Wong, CH. 2003. Carbohydrate microarrays — a new set of technologies at the frontiers of glycomics. 13:637–645.
- Fukui, S, Feizi, T, Galustian, C, Lawson, AM, Chai, W. 2002. Oligosaccharide microarrays for high-throughput detection and specificity assignments of carbohydrate-protein interactions. *Nat Biotechnol.* 20:1011-1017.
- Geijtenbeek, TBH, Torensma, R, van Vliet, SJ, van Duijnhoven, GCF, Adema, GJ, van Kooyk, Y, Figdor, CG. 2000. Identification of DC-SIGN, a novel dendritic cell-specific ICAM-3 receptor that support primary immune responses. *Cell.* 100:575-585.
- Gregg, KJ, Finn, R, Abbott, DW, Boraston, AB. 2008. Divergent modes of glycan recognition by a new family of carbohydrate-binding modules. *J Biol Chem.* 283:12604-12613.
- Guo, Y, Feinberg, H, Conroy, E, Mitchell, DA, Alvarez, R, Blixt, O, Taylor, ME, Weis, WI, Drickamer, K. 2004. Structural basis for distinct ligand-binding and targeting properties of the receptors DC-SIGN and DC-SIGNR. *Nat Struct Mol Biol.* 11:591-598.
- Iobst, ST, Drickamer, K. 1996. Selective sugar binding to the carbohydrate-recognition domains of rat hepatic and macrophage asialoglycoprotein receptors. *J Biol Chem.* 271:6686-6693.
- Karamanska, R, Clarke, J, Blixt, O, MacRae, JI, Zhang, JQ, Crocker, PR, Laurent, N, Wright, A, Flitsch, SL, Russell, DA, Field, RA. 2008. Surface plasmon resonance imaging for real-time, label-free analysis of protein interactions with carbohydrate microarrays. *Glycoconjugate J.* 25:69-74.
- Kolatkar, AR, Leung, AK, Isecke, R, Brossmer, R, Drickamer, K, Weis, WI. 1998. Mechanism of N-acetylgalactosamine binding to a C-type animal lectin carbohydrate-recognition domain. *J Biol Chem.* 273:19502-19508.
- Manimala, JC, Roach, TA, Li, Z, Gildersleeve, JC. 2006. High-throughput carbohydrate microarray analysis of 24 lectins. *Angew Chem Int Ed.* 45:3607-3610.

- Miller, KE, Mukhopadhyay, C, Cagas, P, Bush, CA. 1992. Solution structure of the Lewis x oligosaccharide determined by NMR spectroscopy and molecular dynamics simulations. *Biochemistry*. 31:6703-6709.
- Neu, U, Woellner, K, Gauglitz, G, Stehle, T. 2008. Structural basis of GM1 ganglioside recognition by simian virus 40. *Proc Natl Acad Sci USA*. 105:5219-5224.
- Ng, KK-S, Kolatkar, AP, Park-Snyder, S, Feinberg, H, A., CD, Drickamer, K, Weis, WI. 2002. Orientation of bound ligands in mannose-binding proteins: implications of multivalent ligand recognition. *J Biol Chem*. 277:16088-16095.
- Paulson, JC, Blixt, O, Collins, BE. 2006. Sweet spots in functional glycomics. *Nat Chem Biol*. 2:238-248.
- Perez, S, Mouhous-Riou, N, Nifant'ev, NE, Tsvetkov, YE, Bachet, B, Imberty, A. 1996. Crystal and molecular structure of a histo-blood group antigen involved in cell adhesion: the Lewis x trisaccharide. *Glycobiology*. 6:537-542.
- Petrescu, AJ, Petrescu, SM, Dwek, RA, Wormald, MR. 1999. A statistical analysis of N- and O-glycan linkage conformations from crystallographic data. *Glycobiology*. 9:343-352.
- Powlesland, AS, Fisch, T, Taylor, ME, Smith, DF, Tissot, B, Dell, A, Pöhlmann, S, Drickamer, K. 2008. A novel mechanism for LSECTin binding to Ebola virus surface glycoprotein through truncated glycans. *J Biol Chem*. 283:593-602.
- Powlesland, AS, Ward, EM, Sadhu, SK, Guo, Y, Taylor, ME, Drickamer, K. 2006. Novel mouse homologs of human DC-SIGN: widely divergent biochemical properties of the complete set of mouse DC-SIGN-related proteins. *J Biol Chem*. 281:20440-20449.
- Rosen, SD. 2004. Ligands for L-selectin: homing, inflammation, and beyond. *Annu Rev Immunol*. 22:129-156.
- Somers, WS, Tang, J, Shaw, GD, Camphausen, RT. 2000. Insights into the molecular basis of leukocyte tethering and rolling revealed by structures of P- and E-selectin bound to SleX and PSGL-1. *Cell*. 103:467-479.
- Taylor, ME, Drickamer, K (2006) Introduction to Glycobiology. Oxford University Press, Oxford.
- van Die, I, van Vliet, SJ, Schiphorst, WECM, Bank, CMC, Appelmelk, B, Nyame, AK, Cummings, RD, Geijtenbeek, TBH, van Kooyk, Y. 2002. The dendritic cell specific C-type lectin DC-SIGN recognizes Lewis x, a major glycan epitope of *Schistosoma mansoni* egg antigen. *Glycobiology*. 12:641-642.
- van Liempt, E, Bank, CMC, Mehta, P, Garcia-Vallejo, JJ, Kawar, ZS, Geyere, R, Alvarez, RA, Cummings, RD, van Kooyk, Y, van Die, I. 2006. Specificity of DC-SIGN for mannose- and fucose-containing glycans. *FEBS Lett*. 580:6123-6131.

- van Vliet, SJ, van Liempt, E, Saeland, E, Aarnoudse, CA, Appelmeik, B, Irimura, T, Geijtenbeek, TB, Blixt, O, Alvarez, R, van Die, I, van Kooyk, Y. 2005. Carbohydrate profiling reveals a distinctive role for the C-type lectin MGL in the recognition of helminth parasites and tumor antigens by dendritic cells. *Int Immunol.* 17:661-669.
- Varki, A, Cummings, RD, Esko, JD, Freeze, HH, Stanley, P, Bertozzi, CR, Hart, GW, Etzler, ME (2009) *Essentials of Glycobiology*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York.
- Vestweber, D, Blanks, JE. 1999. Mechanisms that regulate the function of selectins and their ligands. *Physiol Rev.* 79:181-213.
- Wang, D, Liu, S, Trummer, BJ, Deng, C, Wang, A. 2002. Carbohydrate microarrays for the recognition of cross-reactive molecular markers of microbes and host cells. *Nat Biotechnol.* 20:275-281.
- Weis, WI, Drickamer, K. 1994. Trimeric structure of a C-type mannose-binding protein. *Structure.* 2:1227-1240.
- Weis, WI, Drickamer, K, Hendrickson, WA. 1992. Structure of a C-type mannose-binding protein complexed with an oligosaccharide. *Nature.* 360:127-134.

## Figure legends

**Figure 1. Probing glycan arrays and presenting the results.** Some of the issues associated with probing of glycan arrays are illustrated. (a) Almost no single glycan-CRD interactions are strong enough to be measured in a format requiring washing of the array surface, so avidity must be increased by using oligomeric forms of the receptors. These can be natural oligomers, in which stalk regions link two, three or four CRDs in a conformation like that seen in the extracellular domain of a receptor. Alternatively, oligomers can be formed through oligomerisation domains, such a biotinylated tag that allows formation of tetrameric complexes with streptavidin. Higher avidity is sometimes achieved through secondary complexing reagents such as antibodies. Differences in the way avidity is achieved make it very difficult to compare intrinsic affinities of different CRDs for glycans using the arrays. (b) Binding results are often presented in random order based on the position of glycans on the array or grouped by the nature of the oligosaccharides. As illustrated for the comparison of the related receptors DC-SIGN and DC-SIGNR, these approaches are particularly useful for comparing which ligands are preferred by different CRDs, since glycans occur at fixed positions on the x-axis in such graphs. Alternatively, data can be presented in rank order of the observed signals on the array, which can be useful for visualizing the degree of selectivity and demonstrating the effect of concentration of the labeled probe as illustrated in other figures in this review. Data based on Figure 1 in (Guo et al. 2004). (c) Symbol representation of glycan structures. Shapes and colors indicate monosaccharides and partial linkage information is indicated by their geometrical arrangement. Linkages are from the 1 position except for NeuAc, for which linkage is from the 2 position.

**Figure 2. Fundamentals of glycan recognition by C-type lectins.** Ligand binding to C-type CRDs is  $\text{Ca}^{2+}$ -dependent because the primary binding site interacts with monosaccharides through a conserved  $\text{Ca}^{2+}$  that coordinates hydroxyl groups 3 and 4 of the sugar. The arrangement of acid and amide groups on the amino acid side chains that bind this  $\text{Ca}^{2+}$  to the protein determines primary specificity. Mannose-type sugars, such as mannose, GlcNAc and glucose have equatorial 3 and 4 hydroxyl groups (a), while galactose type sugars, galactose and GalNAc, have an axial 4 hydroxyl group (b). In the latter case, affinity for galactose is usually enhanced by secondary interactions with an aromatic residue such as the tryptophan illustrated. Fucose, which has unusual stereochemistry because of its L configuration, sometimes binds to the mannose-type binding sites through its equatorial 2 and 3 hydroxyl groups and sometime in an alternative orientation involving the equatorial 3 hydroxyl group and the axial 4 hydroxyl.

Structures were drawn from entries 1kwu and 1afa in the PDB. This and subsequent molecular graphics figures were created with PyMol ([www.pymol.org](http://www.pymol.org)).

**Figure 3. Ligand binding by the scavenger receptor C-type lectin (SRCL).** (a) Glycan array data for the mouse receptor (200  $\mu\text{g/ml}$ ) plotted in rank order of ligand binding. The ten oligosaccharides given the highest signals are highlighted in red, along with other structurally similar glycans that are ranked lower. (b) Structure of Lewis<sup>x</sup> trisaccharide bound to SRCL. The surface of the protein is shown, with Ca<sup>2+</sup> highlighted in violet. (c) Structure of the complex in (b) with surface colored by underlying atom type and ligand presented as space-filling spheres. The figure was created from CFG primscreen \_GLYCAN\_v3\_72\_02172005 and structure 2OX9 in the PDB.




**Figure 4. Interpretation of glycan array results for LSEctin.** (a) Results for glycan arrays probed with labeled human LSEctin CRD-streptavidin tetramers (4.5  $\mu\text{g/ml}$ ). The structures of the five oligosaccharides giving the highest signals are shown. Bars are coded based on terminal sugars in the glycans: green for mannose and blue for GlcNAc. (b,c) Structure of a portion of a bi-antennary glycan terminating in GlcNAc- $\beta$ 1-2Man disaccharides showing packing of the N-acetyl group of GlcNAc on top of mannose. The figure is based on CFG primscreen\_1043 and structure 1SLA in the PDB.

**Figure 5. Binding of two classes of ligands to DC-SIGN.** (a) Data for screening of a recent version on the Consortium for Functional Glycomics glycan array with human DC-SIGN (200  $\mu\text{g/ml}$ ). Results are presented in rank order of binding, with bars for mannose-terminated glycans colored green and bars for glycans containing the Lewis<sup>a</sup> or Lewis<sup>x</sup> epitopes highlighted in red. The ten oligosaccharides giving the strongest signals are shown at the top and additional related glycans are indicated below. (b,c) Structure of the CRD from DC-SIGN in complex with a tetrasaccharide containing the Lewis<sup>x</sup> epitope. (d,e) Structure of the CRD with Man<sub>5</sub> modeled into the binding site based on the crystal structure with a GlcNAc<sub>2</sub>Man<sub>3</sub> oligosaccharide. The structures were created from entries 1sl5 and 1k9i in the PDB.


**Figure 6. Mode of saccharide binding by the macrophage galactose lectin.** (a) Binding data for the human receptor (4.5 mg/ml) on the CFG printed glycan array. The results are similar to previous studies with the human and rat receptors on earlier streptavidin-based arrays (Coombs et al. 2005, van Vliet et al. 2005). The data are color-coded to indicate glycans terminating in GalNAc with free 3 and 4 hydroxyl groups in pink and galactose with free 3 and 4 hydroxyl groups in orange, except galactose residues in  $\alpha$  linkage or adjacent to fucose residues, which are in yellow. (b,c) Model of the binding site in macrophage galactose lectin with a bound GalNAc residue, based on the structure of the galactose-binding mutant of mannose-binding

protein that was created by insertion of key binding site residues from the galactose-binding receptor. These residues include the marked tryptophan residue and the glycine-rich loop positioned just to the right of this residue, which serves to hold it in position for packing against galactose or GalNAc residues. For the model, additional surface residues were substituted into the structure using Insight software. Data are from CFG primscreen\_2010 and the model was built starting with structure 1afb in the PDB.

**Figure 7. Mechanisms of mannose-binding protein interaction with ligands.** (a) Screening of glycan array with the trimeric terminal fragment of rat serum mannose-binding protein at 250  $\mu\text{g/ml}$  concentration. The oligosaccharides giving the ten highest signals are illustrated at the top. Scale of the y axis is expanded compared to other figures. (b) Structure of a  $\text{Man}_6$  ligand complexed with the CRD of mannose-binding protein. (c,d) Diagrams illustrating the potential effect of glycan spacing on the avidity with which mannose-binding protein trimers bind to dense clusters of sugars on bacterial or fungal surfaces compared to their interaction with glycans on the array. The figure is based on data from CFG primscreen\_1618 and structure 1kx1 in the PDB.



Me- $\alpha$ -Man



Me- $\beta$ -Gal

