

**CHEMISTRY**   
**A EUROPEAN JOURNAL**

Supporting Information

© Copyright Wiley-VCH Verlag GmbH & Co. KGaA, 69451 Weinheim, 2008

# **Synthesis and Relaxometric Studies of a Dendrimer-based pH responsive MRI Contrast Agent**

M. Meser Ali,<sup>[a]</sup> Mark Woods,<sup>[a,b]</sup> Peter Caravan,<sup>[c]</sup> Ana C.L. Opina,<sup>[a]</sup> Marga Spiller,<sup>[d]</sup>  
James C. Fettinger,<sup>[e]</sup> A. Dean Sherry<sup>[a,f]\*</sup>


- a) *Department of Chemistry, University of Texas at Dallas, P.O. Box 830660, Richardson, Texas 75083*
- b) *Macrocyclics, 2110 Research Row, Suite 425, Dallas, Texas 75235*
- c) *A.A. Martinos Center for Biomedical Imaging, Department of Radiology, Massachusetts General Hospital and Harvard Medical School, Charlestown, MA 02129, (UNITED STATES).*
- d) *Department of Radiology, New York Medical College, Valhalla, New York 10595, (UNITED STATES).*
- e) *Department of Chemistry, University of California at Davis, One Shields Avenue, Davis, CA 95616-5298, (UNITED STATES).*
- f) *Advanced Imaging Research Center, University of Texas Southwestern Medical Center, 5323 Harry Hines Boulevard, Dallas, Texas 75390*

## Electronic Supplementary Information S1


**Table S1.** Crystallographic data for **5**

Identification code	jf1426ffmi	
Empirical formula	C <sub>18</sub> H <sub>30</sub> N <sub>2</sub> O <sub>8</sub> P <sub>2</sub>	
Formula weight	464.38	
Temperature	90(2) K	
Wavelength	0.71073 Å	
Crystal system	Orthorhombic	
Space group	P n a 2 <sub>1</sub>	
Unit cell dimensions	a = 19.2299(12) Å	α = 90°.
	b = 15.1795(9) Å	β = 90°.
	c = 15.6480(10) Å	γ = 90°.
Volume	4567.7(5) Å <sup>3</sup>	
Z	8	
Density (calculated)	1.351 Mg/m <sup>3</sup>	
Absorption coefficient	0.236 mm <sup>-1</sup>	
F(000)	1968	
Crystal size	0.46 x 0.41 x 0.25 mm <sup>3</sup>	
Crystal color and habit	Colorless Irregular Block	
Diffractometer	Bruker SMART1000	
Theta range for data collection	1.71 to 30.00°.	
Index ranges	-27 = h = 27, -21 = k = 21, -22 = l = 22	
Reflections collected	51060	
Independent reflections	13322 [R(int) = 0.0393]	
Observed reflections (I > 2σ(I))	11744	
Completeness to theta = 30.00°	100.0 %	
Absorption correction	SADABS, Analytical	
Max. and min. transmission	0.9430 and 0.8994	
Solution method	SHELXS-97 (Sheldrick, 1990)	
Refinement method	SHELXL-97 (Sheldrick, 1997) Full-matrix least-squares on F <sup>2</sup>	
Data / restraints / parameters	13322 / 27 / 645	
Goodness-of-fit on F <sup>2</sup>	1.118	
Final R indices [I > 2σ(I)]	R1 = 0.0572, wR2 = 0.1202	
R indices (all data)	R1 = 0.0655, wR2 = 0.1239	
Absolute structure parameter	0.38(9)	
Largest diff. peak and hole	0.450 and -0.458 e.Å <sup>-3</sup>	

Electronic Supplementary Information S2



**Figure S1.** Packing diagram of the crystal structure of **5** view down the c-axis



**Figure S2.** Packing diagram of the crystal structure of **5** view down the b-axis