

Synthesis of PNA FRET Probes via an Orthogonally Protected Building Block for Post-Synthetic Labeling of PNAs at the 5-Position of U.


Bereket Y. Oquare and John-Stephen Taylor*

Department of Chemistry, Washington University in St. Louis, One Brookings Drive,
St. Louis, Missouri 63130

*To whom correspondence should be addressed:

Department of Chemistry
Campus Box 1134
Washington University
One Brookings Drive
St. Louis, MO 63130
email: taylor@wustl.edu

	<u>Page</u>
Figure S1. ¹ H NMR spectrum of 3 .	S2
Figure S2. ¹³ C NMR spectrum of 3 .	S3
Figure S3. ¹ H NMR spectrum of 4 .	S4
Figure S4. ¹³ C NMR spectrum of 4 .	S5
Figure S5. ¹ H NMR spectrum of 5 .	S6
Figure S6. ¹³ C NMR spectrum of 5 .	S7
Figure S7. ¹ H NMR spectrum of 6 .	S8
Figure S8. ¹ H NMR spectrum of 7 .	S9
Figure S9. ¹³ C NMR spectrum of 7 .	S10
Figure S10. ¹ H NMR spectrum of 11 .	S11
Figure S11. ¹³ C NMR spectrum of 11 .	S12
Figure S12. ¹ H NMR spectrum of 13 .	S13
Figure S13. ¹³ C NMR spectrum of 13 .	S14
Figure S14. ¹ H NMR spectrum of 14 .	S15
Figure S15. ¹³ C NMR spectrum of 14 .	S16
Figure S16. ¹ H NMR spectrum of 16 .	S17
Figure S17. ¹³ C NMR spectrum of 16 .	S18
Figure S18. ¹ H NMR spectrum of 8 .	S19
Figure S19. ¹³ C NMR spectrum of 8 .	S20
Figure S20. ¹ H NMR spectrum of 10 .	S21
Figure S21. ¹³ C NMR spectrum of 10 .	S22
Figure S22. ¹ H NMR spectrum of 1 .	S23
Figure S23. ¹³ C NMR spectrum of 1 .	S24



S2




Figure S1. ^1H NMR (300MHz, DMSO-d_6) spectrum of tert-butyl 2-(5-iodo-2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)acetate (3).





Figure 3.3 ^1H NMR (300 MHz, DMSO-d_6) spectrum of (E)-3-(1-(2-tert-butoxy-2-oxoethyl)-2,4-dioxo-1,2,3,4-tetrahydropyrimidin-5-yl)acrylic acid (**4**).





Figure S4. ¹³C NMR (300 MHz, DMSO-d₆) spectrum of (E)-3-(1-(2-tert-butoxy-2-oxoethyl)-2,4-dioxo-1,2,3,4-tetrahydropyrimidin-5-yl)acrylic acid (4).



S6




Figure S5. ^1H NMR (300 MHz, DMSO-d_6) spectrum of (E)-2,5-dioxopyrrolidin-1-yl-3-(1-(2-tert-butoxy-2-oxoethyl)-2,4-dioxo-1,2,3,4-tetrahydropyrimidin-5-yl)acrylate (**5**).





Figure S6. ¹³C NMR (75 MHz, DMSO-d₆) spectrum of (E)-2,5-dioxopyrrolidin-1-yl-3-(1-(2-tert-butoxy-2-oxoethyl)-2,4-dioxo-1,2,3,4-tetrahydropyrimidin-5-yl)acrylate (5).



6




Figure S7. ¹H NMR (300 MHz, CD₃OD) spectrum of Synthesis of 2-(1-(2-aminoethylamino)ethylidene)-5,5-dimethyl-1,3-cyclohexanedione (6).




Figure S8. ^1H NMR (300 MHz, CD_3OD) spectrum of (E)-tert-butyl 2-(5-(3-(2-(1-(4,4-dimethyl-2,6-dioxocyclohexylidene)ethylamino)ethylamino)-3-oxoprop-1-enyl)-2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)acetate (**7**).




Figure S9. ^{13}C NMR (75 MHz, CD_3OD) spectrum of (E)-tert-butyl 2-(5-(3-(2-(1-(4,4-dimethyl-2,6-dioxocyclohexylidene)ethylamino)ethylamino)-3-oxoprop-1-enyl)-2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)acetate (7).




Figure S10. ^1H NMR(300 MHz, DMSO- d_6) spectrum of (E)-3-(2,4-dioxo-1,2,3,4-tetrahydropyrimidin-5-yl)acrylic acid (11).




Figure S11. ¹³C NMR(75 MHz, DMSO-d₆) spectrum of (E)-3-(2,4-dioxo-1,2,3,4-tetrahydropyrimidin-5-yl)acrylic acid (11).




Figure S12. ^1H NMR (300 MHz, DMSO-d_6) spectrum of (E)-tert-butyl-2-(3-(2,4-dioxo-1,2,3,4-tetrahydropyrimidin-5-yl)acrylamido)ethylcarbamate (**13**).




Figure S13. ¹³C NMR (75 MHz, DMSO-d₆) spectrum of (E)-tert-butyl 2-(3-(2,4-dioxo-1,2,3,4-tetrahydropyrimidin-5-yl)acrylamido)ethylcarbamate (**13**).




Figure S14. ¹H NMR (300 MHz, CD₃OD) spectrum of (E)-N-(2-aminoethyl)-3-(2,4-dioxo-1,2,3,4-tetrahydropyrimidin-5-yl)acrylamide (**14**).




Figure S15. ¹³C NMR (75 MHz, DMSO-d₆) spectrum of (E)-N-(2-aminoethyl)-3-(2,4-dioxo-1,2,3,4-tetrahydropyrimidin-5-yl)acrylamide (13).




Figure S16. ¹H NMR (300 MHz, DMSO-d₆) spectrum of (E)-N-(2-(1-(4,4-dimethyl-2,6-dioxocyclohexylidene)ethylamino)-ethyl)-3-(2,4-dioxo-1,2,3,4-tetrahydropyrimidin-5-yl)acrylamide (**16**).




Figure S17. ¹³C NMR (75 MHz, DMSO-d₆) spectrum of (E)-N-(2-(1-(4,4-dimethyl-2,6-dioxocyclohexylidene)ethylamino)-ethyl)-3-(2,4-dioxo-1,2,3,4-tetrahydropyrimidin-5-yl)acrylamide (**16**).




Figure S18. ^1H NMR (300 MHz, DMSO-d_6) spectrum of (E)-2-(5-(3-(2-(1-(4,4-dimethyl-2,6-dioxocyclohexylidene)ethylamino)ethylamino)-3-oxoprop-1-enyl)-2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl) acetic acid (**8**)




Figure S19. ¹³C NMR (75 MHz, DMSO-d₆) spectrum of (E)-2-(5-(3-(2-(1-(4,4-dimethyl-2,6-dioxocyclohexylidene)ethylamino)ethylamino)-3-oxoprop-1-enyl)-2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl) acetic acid (**8**).




Figure S20. ^1H NMR (300 MHz, CD_3COCD_3) spectrum of (E)-tert-butyl 2-(N-(2-(((9H-fluoren-9-yl)methoxy)carbonylamino)ethyl)-2-(5-(3-(2-(1-(4,4-dimethyl-2,6-dioxocyclohexylidene)ethylamino)ethylamino)-3-oxoprop-1-enyl)-2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)acetamido) acetate (**10**).




Figure S21. ^{13}C NMR (75 MHz, CD_3COCD_3) spectrum of (E)-tert-butyl 2-(N-(2-(((9H-fluoren-9-yl)methoxy)carbonyl)amino)ethyl)-2-(5-(3-(2-(1-(4,4-dimethyl-2,6-dioxocyclohexylidene)ethylamino)ethylamino)-3-oxoprop-1-enyl)-2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)acetamido) acetate (10).




Figure S22. ¹H NMR (300 MHz, DMSO-d₆) spectrum of (E)-2-(N-(2-(((9H-fluoren-9-yl)methoxy)carbonylamino)ethyl)-2-(5-(3-(2-(1-(4,4-dimethyl-2,6-dioxocyclohexylidene)ethylamino)ethylamino)-3-oxoprop-1-enyl)-2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)acetamido)acetic acid; Fmoc-U' (Dde)-OH (1).




Figure S23. ¹³C NMR (75 MHz, DMSO-d₆) spectrum of (E)-2-(N-(2-(((9H-fluoren-9-yl)methoxy)carbonylamino)ethyl)-2-(5-(3-(2-(1-(4,4-dimethyl-2,6-dioxocyclohexylidene)ethylamino)ethylamino)-3-oxoprop-1-enyl)-2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)acetamido)acetic acid; Fmoc-U' (Dde)-OH (1).