

Transplant	Organ	Genotype	B220+ cells(%)	CD3+ cells (%)	Mac1+ or Gr-1+ cells (%)
1 st T	PB	+/+	59.0±7.2	12.9±7.0	20.0±6.8
		-/-	64.3±9.6	11.6±5.5	15.7±2.1
1 st T	BM	+/+	27.5±7.0	4.0±2.1	68.2±7.5
		-/-	26.2±2.1	2.8±0.5	70.2±3.0
2 nd T	PB	+/+	52.9±4.8	28.4±5.0	18.8±1.6
		-/-	51.7±3.1	27.6±2.7	20.2±2.8
2 nd T	BM	+/+	28.6±11.4	6.0±4.3	61.8±4.8
		-/-	23.3±5.9	4.6±4.2	69.0±4.8
3 rd T	PB	-/-	42.2±16.6	8.1±2.9	40.8±15.1
3 rd T	BM	-/-	10.2±5.3	1.4±0.5	86.1±6.1

B (B220), T (CD3ε) and myeloid (Mac-1 and Gr-1) cells were analyzed by FACS on gated CD45.2+ PB or BM cells 4 months post transplantation. Data shown are the representative experiment from 3 independent experiments, and represent the percentage of each lineage in donor-derived cells (mean ± SD, n=6–10)