

	<i>P. elephas</i>	<i>P. mauritanicus</i>	<i>P. charlestoni</i>	<i>P. gilchristi</i>	<i>P. delagoae</i>	<i>P. barbarae</i>
Pael-10	0.351	0.166	0.533	0.050	0.097	-
Pael-11	0.634	0.494	0.378	0.789	0.645	0.733
Pael-12	0.937	0.882	0.933	0.867	0.883	0.870
Pael-14	0.926	0.757	0.911	0.918	0.869	0.902
Pael-20	0.865	0.954	0.956	0.959	0.883	0.918
Pael-21	0.918	0.765	0.711	0.722	0.774	0.743
Pael-22	0.520	0.568	0.714	0.603	0.520	0.542
Pael-28	0.821	0.770	0.778	0.701	0.500	0.703
Pael-31	0.899	0.803	0.929	0.868	0.864	0.517
Pael-44	0.964	0.938	0.933	0.969	0.956	0.960
Pael-48	0.794	0.570	0.889	0.879	0.881	0.694
Pael-49	0.882	0.775	0.511	0.776	0.709	0.661
Pael-53	0.782	0.405	0.733	0.765	0.323	0.223

All markers were shown polymorphic in every tested species with the exception of PE10 in *P. barbarae*.